

2018-01

Pliocene stratigraphic paleobiology in Tuscany and the fossil record of marine megafauna

Dominici, S

<http://hdl.handle.net/10026.1/10408>

10.1016/j.earscirev.2017.09.018

Earth-Science Reviews

Elsevier

All content in PEARL is protected by copyright law. Author manuscripts are made available in accordance with publisher policies. Please cite only the published version using the details provided on the item record or document. In the absence of an open licence (e.g. Creative Commons), permissions for further reuse of content should be sought from the publisher or author.

Manuscript Details

Manuscript number	EARTH_2017_214_R1
Title	Pliocene stratigraphic paleobiology in Tuscany and the fossil record of marine megafauna
Article type	Review Article

Abstract

Tuscany has a rich Pliocene record of marine megafauna (MM), including mysticetes, odontocetes, sirenians and seals among the mammals, and six orders of sharks among the elasmobranchs. This is reviewed with respect to paleogeography and sequence-stratigraphic occurrence in six different basins. Conditions at the ancient seafloor are explored by means of sedimentary facies analysis, taphonomy and multivariate techniques applied to a large quantitative dataset of benthic molluscs. MM is rare or absent in most basins during the Zanclean, except in one basin, and most abundant in Piacenzian deposits in all six basins. MM occurs preferentially in fine-grained shelfal highstand deposits of small-scale depositional sequences, or at condensed horizons of the maximum flooding interval. It is rare in shallow marine paleoenvironments and nearly absent in bathyal paleosettings. Paleogeographic and paleoecological evidence and a comparison with modern patterns of marine upwelling suggest that a wedge of nutrient-rich waters sustained in the offshore during the Pliocene a high biomass of primary producers and a community of apex consumers and mesopredators, similarly to the modern Northwestern Mediterranean Sea, with a higher species-richness than the modern. The highest MM diversity coincides with the mid-Piacenzian warm period, suggesting that facies control does not obscure a link between climate and diversity. We underline however that not all marine environments were suitable for marine mammal preservation. Buoyant carcasses were preferentially dismembered and destroyed in high-energy shallow waters, with the possible exception of delta front deposits, where sudden sediment input occasionally buried pristine carcasses. We hypothesise that carcasses sunken on the seafloor below the shelf break underwent destruction through the activity of a whale-fall biota of modern type, specialised in the consumption of decomposing tissues, both soft and mineralised. A taphonomic window was left between storm wave base and the shelf break. Here water pressure is high enough to prevent the formation of decomposing gases and the resurfacing of carcasses, while the lack of a specialised whale-fall biota slows down bone degradation with respect to deeper settings. Sedimentation rate was high enough to cover skeletal material before its complete destruction. An estimate of paleobathymetries based on multivariate techniques suggests that the preferential depth for the inclusion of MM in the fossil record was 30-300 m. The results are compared with major Mesozoic and Cenozoic MM records worldwide. Available evidence suggests that the late Neogene radiation of large whales, true ecosystem engineers, and their size increase, triggered the radiation of a bone-eating fauna that hampered, and hampers, MM preservation in the deep sea. Stratigraphic paleobiology and an ecosystem-level approach deliver useful insights in the nature of the fossil record.

Keywords Cetacea; Elasmobranchia; whales; dolphins; sharks; Pliocene; Sirenia; sequence stratigraphy; taphonomy; marine megafauna; Mollusca; open marine ecosystem structure; top-down control; shell bed; multivariate analysis; paleodepths; upwelling; Mediterranean; diversity; megabias

Corresponding Author Stefano Dominici

Order of Authors Stefano Dominici, Silvia Danise, Marco Benvenuti

Suggested reviewers Michelangelo Bisconti, Michael Benton, Shanan E. Peters, Steven Holland

Submission Files Included in this PDF

File Name [File Type]

Copy-of-Pliocene NWMS MM cover letter.docx [Cover Letter]

Pliocene NWMS MM cover letter.docx [Response to Reviewers]

Pliocene NWMS MM Submitted_September.docx [Revised Manuscript with Changes Marked]

Pliocene NWMS MM Revised Abstract.docx [Abstract]

Pliocene NWMS MM Submitted_September_untracked.docx [Manuscript File]

Fig. 1.tif [Figure]

Fig. 2.tif [Figure]

Fig. 3.tif [Figure]

Fig. 4.tif [Figure]

Fig. 5.tif [Figure]

Fig. 6.tif [Figure]

Fig. 7.tif [Figure]

Fig. 8.tif [Figure]

Fig. 9.tif [Figure]

Fig. 10.tif [Figure]

Fig. 11.tif [Figure]

Fig. S1.tif [Figure]

Submission Files Not Included in this PDF

File Name [File Type]

Tables 1-6, S1-S4 - Pliocene Tuscany REVISED2.xlsx [Table]

To view all the submission files, including those not included in the PDF, click on the manuscript title on your EVISE Homepage, then click 'Download zip file'.

Dear editor,

we submit a revised version of the manuscript now entitled “Pliocene stratigraphic paleobiology in Tuscany and the fossil record of marine megafauna”. We have carefully considered the comments of the first reviewer and your comments and revised the manuscript accordingly.

We look forward hearing from you.

For the authors,

Best regards,

Stefano Dominici

Dear editor,

we submit a revised version of the manuscript now entitled "Pliocene stratigraphic paleobiology in Tuscany and the fossil record of marine megafauna". We have carefully considered the comments of the first reviewer and your comments and revised the manuscript accordingly. In particular, we would like to explicit the following answers (in red) to review no. 1 (in blue):

Some suggestions that could improve the manuscript:

1) The ambitious paleoecological considerations intended to link the supposed high diversity/abundance of the marine vertebrate assemblage to the paleobiogeography/food availability of the northwestern Mediterranean are, in my opinion, not supported by solid data. In fact:

A) data about the systematic composition of the vertebrate fossil assemblage are only reported (for the marine mammals) in table 1 and not discussed in the text (note that I disagree with the authors' statement about "the systematics of Mediterranean Pliocene LMV have been recently reviewed"); data about the sharks are not reported;

B) also admitting that the systematic assignments reported in this list are sufficient for this analysis, no data are reported about, for example, the trophism and, more generally, the ecological behavior of the listed taxa;

C) diversity and abundance are cited for the vertebrates, but not quantified;

D) the link between diversity and primary productivity/food availability must be better analyzed considering the rich and often controversial bibliography on this topic;

E) baleen whales and other vertebrates from the Pliocene of Tuscany are here considered 'apex predator': the authors must define better this ecological term. In my opinion, baleen whales cannot be included among the top predators, as well as other 'large' marine vertebrates as they feed on plankton and small fishes with filtering techniques. Apex or top predators can be considered some (but not all) 'large' sharks as *Carcharodon carcharias* and *Cosmopolitodus hastalis*, and, among cetaceans, *Orcinus citoniensis*.

So, I suggest the authors to review the part focused on the paleoecology of the vertebrate assemblage, improving the data. Alternatively, it could be drastically shortened or totally removed.

1A: We enlarged the marine mammal dataset, from 39 to 64 reports, and carefully controlled existing literature on the systematics.

1B: We have greatly enlarged the literature on the ecology of modern Mediterranean marine mammals and introduced data on the ecology of sharks. Based on these data, we attributed a preferred prey and a feeding mechanism to each taxon of our list.

1C: We calculated minimum number of marine mammal specimens. The systematics were sufficient for a conservative estimate of standing diversity in terms of species richness of both marine mammals and sharks. Marine mammal abundance and richness have been calculated for six consecutive intervals and for the overall epoch and compared to modern values for the north-eastern Mediterranean. A Pliocene-modern comparison is also given for species richness in sharks. We introduced new tables, and two histograms (Fig. 9C-D), for an easy understanding of diversity trends during the Pliocene.

1D: In expanding the reference list, we have highlighted the overwhelming evidence for a relationship between marine megafauna diversity and primary productivity (bottom-up control on marine ecosystems), but also decoupled this from structural factors such as trophic cascade (top-down controls exerted by large predators).

1E: There's no formal definition of what "marine megafauna" is, but practice adopted also by palaeontologists includes all marine mammals, seabirds, sea turtles, sharks and rays (e.g., Lewison et al., 2004; Pimiento et al., 2017). We clearly stated what we mean by apex predator, using concepts derived from modern ecological literature. In particular we specify that the definitions of apex predators (or top predators) and mesopredators are relative, and to an extent context-dependent (species that in a contact are apex predators, in another are mesopredators). By any means, filter-feeding baleen whales are considered apex-predators of their community. To avoid misunderstanding between ecological and paleoecological usage, we informed on the trophic level of each Pliocene species by comparison with modern values, for both marine mammals and sharks.

2) The authors propose a link between the Pliocene-Pleistocene increase in whale size and in bone-eaters worms diversity: these considerations should be better supported by including more bibliographical data and also by an eventual analysis of the Pliocene Tuscany vertebrate assemblage. Are the marine vertebrates from Pliocene of Tuscany particularly large? If yes, please support this affirmation with data. My preliminary evaluation is that there are some large baleen whales (but not 'giant whales' as today), some medium-small sized baleen whales, and several relatively small marine mammals

(odontocetes, pinnipeds and sirenians). So, why also the 'small' vertebrates are not preserved in the deep sea sediments of the Pliocene of Tuscany?

We have given figures for size differences between Pliocene and modern marine megafauna. We are explicit in stating the complex nature of the comparison between MM at different times, but we are also clear in relating the radiation of bone-eating worms, and the importance of a depth-related taphonomic overprint in the Pliocene fossil record, to the emergence of large whales. That is to say, not all MM were large, but some were. We also see no need to limit this effect to the carcass of true giants, like the modern blue whale. That is to say, we think that large is large enough.

3) The analysis is focused on the 'large' marine vertebrates: in my opinion 'large' is rather vague as it is not well defined by the authors. It seems that the authors consider as 'large vertebrates' all cetaceans (including the smaller odontocetes), all pinnipeds and all sirenians, among the tetrapods (but not the marine turtles and birds), and Carcharhiniformes (note that several carchariformes, as the catsharks, are not longer than 50 cm) and Lamniformes among sharks: it is right?

As stated above, the ecological definition of "marine megafauna" includes smaller animals (sea birds, otters, turtles). We focus on the general structure of the MM association, not on the role of a particular species.

4) in the analysis are also included some 'large' sharks: their fossil record is exclusively based on isolated teeth and most of these are found associated to cetacean skeletons. This should be better outlined, since these remains have a different taphonomic significance than the bones of the other vertebrates

We have expanded the review of the Pliocene shark record of Tuscany in Tables 5-6 and discussed it in the text. Taphonomy has been expanded, also in relationship to evidence of interaction between different MM species.

5) the organization of the manuscript, including the presentation of data, results and discussion is often unclear and could be improved. Some suggestions:

- the numeration of the chapters/paragraphs must be revised (the number 3 is repeated three times)

- 178
179 - all the chapters from 'Distribution of large marine vertebrates', to 'the Biotic factors' (lines
180 950-2012) must be included in the 'Results'
181
182 - the paragraph '6.2 Taphonomic control' reports an interesting bibliographical review, but
183 no reference to the material here examined.
184
185
186

187 We have corrected the organization of the ms as suggested. Tables have been expanded
188 and augmented in number.
189
190
191

192 6) Table 2 appears as the more important source of this paper, since it condenses all the
193 data about the marine mammals discussed and elaborated in the work. By contrast, the
194 table does not seem to have been carefully prepared. For example, most of the specimens
195 lacks the description, although these data can be easily found in bibliography (e.g.: the
196 *Orcinus citoniensis* holotype is an articulated partial skeleton) and the few data reported
197 are often wrong (e.g. MOM2-2 and WSi2-1 are not articulated skeleton); moreover it is not
198 easy to identify the specimens reported without the catalogue number of the museum
199 where they are kept. For example it is not possible identify WFi4-3 from Orciano (no data
200 about the systematics and the bones preserved) considering that Lawley (1876) reported
201 several remains from the same locality; similarly, it is reported an articulated skeleton of
202 sirenian from Arcille (MOM2-2) based on Tinelli (2013): but Tinelli (2013) described 4
203 skeletons from Arcille (3 disarticulated and 1 articulated; see also: Sorbi et al. 2011, Tinelli
204 et al. 2012).
205
206
207
208
209
210
211
212
213

214 We have expanded Tab. 2 and included much more taphonomic information, making it
215 more homogenous and informative, also with respect to references. We have excluded
216 specimens cited, but no longer traceable (e.g. Lawley 1876).
217
218
219

220 7) All 'dugongs' in the text must be changed with 'sirenians' ('dugong' is referred only to
221 the extant *Dugong dugong* species)
222
223
224

225 Done
226
227

228 8) The fossil vertebrates from Tuscany are named in the text using a inhomogeneous
229 approach about their systematic assignation: I suggest to uniform (maybe using the more
230 detailed determination available from the literature for all specimens cited.
231
232
233
234
235
236

Done (see new Tab. 2)

9) It is unclear if the logs of Figure 4 have been measured by the authors or taken from bibliography. In the second case, I suggest to add the references. This is recommended particularly for the ones taken from the PhD thesis of Chiara Tinelli. This thesis is available online, but, since it has not yet been published, it is important to make clear when the authors used these data.

Done, references added in the caption of Fig. 2

10) In the log of Arcille locality (figure 4) the sirenian is placed at the base of the HST. I personally participated to the collection of the 4 skeletons discovered in this locality and all were found inside the TST, under the HST (see also Tinelli et al. 2012; Tinelli, 2013).

Done (we refer to a “maximum flooding interval”, instead of MFS)

Other more punctual corrections and comments are reported in the annotated manuscript.

We corrected the text following comments annotated by the reviewer on the ms

We would also like to answer your comments:

This is an interesting contribution about the fossil record of marine vertebrates and its relationship to depositional environments. Although it is focused on the Pliocene of Tuscany, the topic is wide enough to be of interest to the readership of Earth-Science Reviews.

Both reviewers liked the manuscript. Nevertheless, Reviewer 1 makes many valuable comments and suggestions on how to improve it. For example, he thinks that “The ambitious paleoecological considerations....are...not supported by solid data”, that more data are need to support the relationship between whale size and bone-eating worm diversity, that the definition of “large vertebrates” is quite ambiguous, and that the organisation of the manuscript and the tables need improvement.

Reviewer 2 is satisfied with the manuscript the way it is now. He says that the coordinates of the studied localities should be added, and that the data should be deposited in the Paleobiology Database.

We look forward uploading our data in PBDB

I have looked through the manuscript myself and have to agree with Reviewer 1: there is need for a moderate to major revision to make this basically nice and interesting study ready for publication. Reviewer 1 suggests to either improve or eliminate the palaeoecological analysis. I am strongly in favour of keeping it in (this makes the paper interesting also for a non-specialized scientific community) but in this case complement and improve the data. Consequently, the manuscript should be expanded to accommodate more data and more explanations. Also, please polish the tables (including the supplementary ones) and the general organization of the manuscript.

Additional suggestions from my side: In the legend to Figure 3, say “SCUBA diver” and not “man” for scale. In the legend to Figures 4 and 8, explain the sequence-stratigraphic abbreviations.

We kept the paleoecological analysis, as you suggested, and largely expanded the database and the reference list so as to sustain our point. We corrected Fig. 3 caption as suggested.

We look forward hearing from you.

For the authors,

Best regards,

Stefano Dominici

SPliocene stratigraphic paleobiology in ~~the Pliocene of~~

Tuscany and the fossil record of ~~large marine~~

~~vertebrates~~marine megafauna

Stefano Dominici (Università di Firenze, Museo di Storia Naturale, Firenze, Italy)

Silvia Danise (Plymouth University, School of Geography, Earth and Environmental Sciences,
Plymouth, United Kingdom)

Marco Benvenuti (Università di Firenze, Dipartimento di Scienze della Terra, Firenze, Italy)

Abstract

Tuscany has a rich Pliocene record of marine megafauna (MM), including mysticetes, odontocetes, sirenians and seals among the mammals, and six orders of sharks among the elasmobranchs. This is reviewed with respect to paleogeography and sequence-stratigraphic occurrence in six different basins. Conditions at the ancient seafloor are explored by means of sedimentary facies analysis, taphonomy and multivariate techniques applied to a large quantitative dataset of benthic molluscs. MM is rare or absent in most basins during the Zanclean, except in one basin, and most abundant in Piacenzian deposits in all six basins. MM occurs preferentially in fine-grained shelfal highstand deposits of small-scale depositional sequences, or at condensed horizons of the maximum flooding interval. It is rare in shallow marine paleoenvironments and nearly absent in bathyal paleosettings. Paleogeographic and paleoecologic evidence, and a comparison with modern patterns of marine upwelling suggest that a wedge of nutrient-rich waters sustained in the offshore during the Pliocene a high biomass of primary producers and a community of apex consumers and mesopredators, similarly to the modern Northwestern Mediterranean Sea, with a higher species-richness than the modern. The highest MM diversity coincides with

the mid-Piacenzian warm period, suggesting a link between climate and diversity. We propose that not all marine environments were suitable for marine mammal preservation. Buoyant carcasses were preferentially dismembered and destroyed in high-energy shallow waters, with the possible exception of delta front deposits, where sudden sediment input occasionally buried pristine carcasses. We hypothesise that carcasses sunken on the seafloor below the shelf break underwent destruction through the activity of a whale-fall biota of modern type, specialised in the consumption of decomposing tissues, both soft and mineralised. A taphonomic window was left between storm-wave base and the shelf break. Here water pressure is high enough to prevent the formation of decomposing gases and the resurfacing of carcasses, while the lack of a specialised whale-fall biota slows down bone degradation with respect to deeper settings. Sedimentation rate is high enough to cover skeletal material before its complete destruction. An estimate of paleobathymetries based on multivariate techniques suggests that the preferential depth for the inclusion of MM in the fossil record is 30-300 m. The results are compared with major Mesozoic and Cenozoic MM records worldwide. Available evidence suggests that the Neogene radiation of large whales, true ecosystem engineers, triggered the radiation of a bone-eating fauna that hampered, and hampers, MM preservation in the deep sea. Tuscany has a rich Pliocene record of marine megafauna (MM), including mysticetes, odontocetes, sirenians and seals among the mammals, and six orders of sharks among the elasmobranchs. This is reviewed with respect to paleogeography and sequence-stratigraphic occurrence in six different basins. Conditions at the ancient seafloor are explored by means of sedimentary facies analysis, taphonomy and multivariate techniques applied to a large quantitative dataset of benthic molluscs. MM is rare or absent in most basins during the Zanclean, except in one basin, and most abundant in Piacenzian deposits in all six basins. MM occurs preferentially in fine-grained shelfal highstand deposits of small-scale depositional sequences, or at condensed horizons of the maximum flooding interval. It is rare in shallow marine paleoenvironments and nearly

absent in bathyal paleosettings. Paleogeographic and paleoecological evidence and a comparison with modern patterns of marine upwelling suggest that a wedge of nutrient-rich waters sustained in the offshore during the Pliocene a high biomass of primary producers and a community of apex consumers and mesopredators, similarly to the modern Northwestern Mediterranean Sea, with a higher species-richness than the modern. The highest MM diversity coincides with the mid-Piacenzian warm period, suggesting that facies control does not obscure a link between climate and diversity. We underline however that not all marine environments were suitable for marine mammal preservation. Buoyant carcasses were preferentially dismembered and destroyed in high-energy shallow waters, with the possible exception of delta front deposits, where sudden sediment input occasionally buried pristine carcasses. We hypothesise that carcasses sunken on the seafloor below the shelf break underwent destruction through the activity of a whale-fall biota of modern type, specialised in the consumption of decomposing tissues, both soft and mineralised. A taphonomic window was left between storm wave base and the shelf break. Here water pressure is high enough to prevent the formation of decomposing gases and the resurfacing of carcasses, while the lack of a specialised whale-fall biota slows down bone degradation with respect to deeper settings. Sedimentation rate was high enough to cover skeletal material before its complete destruction. An estimate of paleobathymetries based on multivariate techniques suggests that the preferential depth for the inclusion of MM in the fossil record was 30-300 m. The results are compared with major Mesozoic and Cenozoic MM records worldwide. Available evidence suggests that the late Neogene radiation of large whales, true ecosystem engineers, and their size increase, triggered the radiation of a bone-eating fauna that hampered, and hampers, MM preservation in the deep sea. Stratigraphic paleobiology and an ecosystem-level approach deliver useful insights in the nature of the fossil record.

1. Introduction

The modern marine megafauna (MM) includes all marine mammals, seabirds, sea turtles and sharks, apex consumers that influence their associated ecosystems (Lewison et al., 2004), both pelagic and nearshore, through top-down forcing and trophic cascades, and now severely affected by human impact (Estes et al., 1998, 2011, 2016). Large vertebrates have played a key role in structuring marine ecosystems. On a macroevolutionary scale, predation pressure has shaped the evolution of marine preys, with feedbacks on predators, setting the stage for the Mesozoic marine revolution (Vermeij, 1977; Chen and Benton, 2012; Benton et al., 2013). The new ecosystem structure started in the the Early and Middle Triassic with several lineages of Actinopterygian fishes (Chen and Benton, 2012), continuing with marine reptiles possessing feeding styles (Fröbisch et al., 2013; Motani et al., 2015; but see also Motani et al., 2013) and reproductive adaptations (Motani et al., 2014) of modern type. Triassic and Jurassic novelties underwent a prolonged crisis during the Cretaceous, with the gradual extinction of plesiosaurs, and mosasaurs (Benson et al., 2010) and ichthyosaurs (Fischer et al., 2016), and a diversity drop of sharks (Guinot et al., 2012), but marine giants. A marine megafauna of comparable size of comparable size returned in the Paleogene, with the new diversification of neoselachian elasmobranchs (Kriwet and Benton, 2004) and the evolution of large marine mammals: Eocene archaeocetes (Uhen, 2008, Gingerich et al., 2009) and Oligocene odontocetes and mysticetes (Gingerich, 2005; Marx and Uhen 2010, Berta, 2012; Marx et al., 2016) empowered by high metabolic rates and new anatomic features (Armfield et al., 2013). Among the largest vertebrates of all times, after a dramatic size increase at the outset of glacial age (Marx et al., 2016; Bisconti et al., 2017; Slater et al., 2017), baleen and sperm whales are among today's ocean's ecosystem engineers (Roman et al., 2014) with which to compare their Mesozoic analogues (Smith et al., 2016). Notwithstanding an enormous

crucial ecological and evolutionary importance in ecology and evolution of these animals, the nature and distribution of the MM fossil record of large marine vertebrates (LMVs) has been relatively little explored, compared to that of marine invertebrates and terrestrial vertebrates. Available data suggest a strong correlation between taxic diversity and the number of marine fossiliferous formations, resulting in megabiases in the fossil record (e.g., Cretaceous: Benson et al., 2010). Within its vast history, studies on the geologically recent marine megafauna offer important insights, thanks to a considering our better knowledge of: 1) geological setting, in terms of outcrop extent and high-resolution stratigraphy; 2) ecologic role played by individual species, whether extant or extinct, in terms of habitat, trophic role, life histories and population structure, thanks to a comparison with extant descendants, or close relatives; 3) MM taphonomy, based on actuopaleontology. A recent global study revealed that MM extinction peaked in the late Pliocene, between 3.8 and 2.4Ma, linked with the sudden drop in the extension of nearshore environments due to a large sea level regression, confirming that the fossil record offers important insights on the vulnerability of keystone marine species to climate change (Pimienta et al., 2017).linked to the sudden drop in the extension of nearshore environments after a large sea level regression (Pimienta et al., 2017), confirming that the fossil record offers important clues on the vulnerability of keystone marine species to climate change. We contribute here to their a better understanding of the Pliocene fossil record by reviewing the rich and varied fossil record of Pliocene LMVsMM of Tuscany, in Italy. In particular, we consider all reports of Pliocene cetaceans, sharks, sirenians and pinnipedsmarine mammals and sharks and revise taphonomy and the sedimentary facies associated with all majorknown recent findings, framingsetting them within a sequence stratigraphic framework. At the same time we expand on We also expand knowledge on paleoenvironmental contextualisation by considering the paleoecology of fossil invertebrates on a regional basisWe also reconstruct the paleoenvironmental context and review data on a part of the marine ecosystem through

the paleoecology of fossil invertebrates on a regional basis, following a stratigraphic paleobiological approach that can be applied to both the recent and the distant geological past (Patzkowsky & Holland, 2012). Published studies that have taken this direction are still a few, examples concerning Jurassic ichthyosaurs, plesiosaurs and pliosaurs (McMullen et al., 2014), Cretaceous turtles, plesiosaurs, bony fish and sharks (Schemisser McKean & Gillette, 2015), Eocene archaeocetes, dugongs/sea cows and sharks (Peters et al., 2009), and Neogene whales, pinnipeds/marine mammals and sharks (Boessenecker et al., 2014). All of these papers record the co-occurrence of shelly faunas, only one undertaking quantitative studies of the distribution of fossil invertebrates associated with sedimentary facies (Jurassic of the Sundance Formation: McMullen et al., 2014; see also Danise & Holland, 2017). The benefits of an outcrop-scale sequence stratigraphic approach include: (1) an independent record of relative sea-level change to test paleobiological hypotheses (see also Pyenson & Lindberg, 2011; Noakes et al., 2013); (2) a chronostratigraphic scheme for high-resolution correlations; (3) a means to recognise minor and major breaks of the record; (4) an ecological and sedimentary framework for taphofacies distribution (Patzkowsky & Holland, 2012); and (5) an independent control of onshore-offshore patterns of fossil assemblages (e.g., Tomašových et al., 2014).

~~From a top-down approach that emphasises the role of large marine vertebrates (LMVs) as apex predators of their trophic web, r~~ Researchers that study the geologic history of marine ecosystems have focused on patterns of ecological restructuring based on the taxonomy of selected groups (e.g., Thorne et al., 2011; Benton et al., 2013; Scheyer et al., 2014; Fischer et al., 2016), at the expenses of a more holistic approach that includes functional diversity and embraces as many ecosystem components as possible (Dineen et al., 2014). ~~This attitude should be reversed, similarly to~~ By analogy with ecologists working on extant ecosystems who shift focus from models based on single groups (e.g., Steeman et al., 2009) to an all-embracing vision of marine life (Lawton, 1994; Sergio et al., 2014).

connecting food web ecology with landscape ecology (Polis et al., 1997; Estes et al., 2011), stratigraphic paleobiology can draw from the fossil record and offer multidimensional insights on the complex geological history of modern marine ecosystems.

After revising ~~LMVs~~fossil MM hosted in major museums of Tuscany, both isolated and articulated remains, we ~~then~~ focus on all fossil ~~skeletons~~bones that ~~could~~can be stratigraphically (e.g., Bianucci et al., ~~1995~~1995~~1998~~2001; Tinelli ~~2013~~2013) and taphonomically framed (e.g., Dominici et al. ~~2009~~2009; Bianucci ~~2010~~2010; Danise ~~&and~~ Dominici, 2014). ~~The systematics of MM lists for the~~ Mediterranean Pliocene ~~LMV~~ have been recently ~~reviewed~~updated (~~cetaceans~~marine mammals: Landini et al., 2005; Bianucci et al., 2009a; Sorbi et al., 2012; Bianucci ~~&and~~ Vomero, 2015~~4~~4; ~~elasmobranchs~~sharks: Marsili, 2006; ~~sirenians~~Sorbi et al. 2012). Species-level ecological data, ~~are~~ available ~~from~~on modern ~~Mediterranean studies of apex predators~~consumers and mesocarnivores (Pauly et al., 1998; Cortés, 1999), with detailed information made available for Mediterranean species following conservation concerns ~~(e.g., baleen and sperm whales~~marine mammals: Notarbartolo di Sciara, et al., 2003, 2008, ~~2016~~2016; Pirota et al. 2011), with ~~species-level data~~ (~~;~~ sharks: Cavanagh ~~&and~~ Gibson, 2007), allowing for a ~~significant~~detailed paleoecological evaluation of the Tuscan fossil record. The actualistic approach is also viable for species of benthic molluscs, about half of which are still extant in modern Mediterranean sea floors~~bottoms~~ (55% of extant species of Mediterranean and North Sea bivalves, excluding strictly brackish and bathyal forms, i.e., 202 out of 367 species, survives from the Zanclean: Raffi et al., 1985). The regional quantitative study of molluscan assemblages was the basis for an independent assessment of paleoenvironments, paleoecology and paleobathymetry. A revisione of abundance distributions of marine molluscs, the largest contributors to Mediterranean Pliocene shell beds and a key component of Mesozoic and Cenozoic marine ecosystems (Stanley, 1975~~17~~17; Vermeij, 1977), allowed to further explore the structure and composition of ~~the benthic component of~~ Pliocene marine ecosystems, and reconstructing

a paleobathymetric gradient (e.g., Scarponi ~~&and~~ Kowaleski, 2004) along which to frame ~~LMVMM taphonomy occurrence~~. The present work must necessarily start with a review of the chronostratigraphy and physical stratigraphy of the Tuscan marine Pliocene.

1. Geological setting

The Pliocene succession of Tuscany was deposited in a complex setting characterized ~~sed~~ by continental collision related to the later evolution of the Northern Apennines chain. According to a well-established hypothesis, the region, affected by shortening before the Middle-Late Miocene, accommodated by NE-verging thrust and fold systems, underwent crustal extension during the late Neogene and the Quaternary (DeCelles, 2012; Fig. 1). Crustal extension generated differential subsidence in a series of normal-fault controlled hinterland sedimentary basins, filled throughout by continental and shallow marine, mostly clastic successions (Martini ~~&and~~ Sagri, 1993; Pascucci et al., 2006; Brogi, 2011). An alternative hypothesis places the late orogenic hinterland basins in a more complex tectonic setting characterized ~~sed~~ by the alternation of compressive, extensional and transcurrent stress fields (Benvenuti et al., 2014; Bonini et al., 2014).

The Neogene Tuscan basins considered in this work include, from West to East, and from North to South, the Fine Basin (FB; Bossio et al., 1997), the Volterra-Era Basin (VEB; Bossio et al., 1994), The Elsa Basin (EB; Benvenuti et al., 2014), the Ombrone-Orcia Basin (OOB; Bossio et al., 1991; Nalin et al., 2010), the Siena-Radicofani Basin (SRB; Ghinassi ~~&and~~ Lazzarotto, 2005; Martini et al., 2011, 2016), and the Chiana Basin (CB; Fig. 1: Pesa Basin not considered here). With one exception (OOB, see below), these basins show a shape conditioned by the structural and physiographic features of the inner portion of the Northern Apennines. Their NW-SE general elongation reflects the trend of the thrust-related anticline ridges developed during earlier collisional stages. These compressive structures have

bounded most basins through their infilling, only to be obliterated by younger parallel normal fault systems, leaving an invariant stratigraphic onlap of the Pliocene successions onto the basin margins. Despite a NW-SE distribution of the hinterland basins, the structural setting is responsible for a NE-trending physiographic and paleogeographic gradient, where the FB is closest, and the CB furthest, from offshore settings throughout the late orogenic phase, with important implications for the facies architecture and the distribution of marine vertebrates and shell beds. Differential active uplift of the basin shoulders during the Pliocene, coupled with important erosional phases, resulted in a different preservation of the original stratigraphic architecture. The infill during the Zanclean is generally characterized by relatively continuous open marine successions, the correlative fluvial-coastal systems missing due to uplift and erosion of basin margins. On the other hand, the Piacenzian infill is characterized by genetically-connected, fluvial, coastal and shallow marine facies tracts, particularly well-preserved in the EB, hinting to reduced uplift of the marginal areas. The modern physiography of OOB, escaping the structurally-controlled geometry of the other basins, mimics instead an original fluvial network developed during the latest Messinian, flooded after the Salinity Crisis (Bossio et al., 1991; Benvenuti et al., 2015), filled during the Pliocene, and finally disrupted by post-Pliocene uplift and erosion. This difference in the structural history also justifies the preservation in OOB of Zanclean fluvial and shallow marine facies (Fig. 2).

1.1 Pliocene Stratigraphy

The Neogene succession of Tuscany is up to 2000 m-thick, about half of which belongs to the uppermost Miocene-Pleistocene interval (Bossio et al., 2004; Benvenuti et al., 2014). The Pliocene has been traditionally subdivided into three main informal lithostratigraphic units: continental conglomerates and sandstones at the base, overlain by

the “Blue Clay Formation” (*Argille Azzurre*: Zanclean-lower Piacenzian), marking the post-Messinian Mediterranean marine transgression and forming the thickest part of the basin infill (e.g., Bossio et al., 1994, 1997; Ghinassi & Lazzarotto, 2005), and the “Upper Sands” (*Sabbie superiori*: Piacenzian-Gelasian) and conglomerates, deposited during the ensuing regression. Several finer lithostratigraphic units have been introduced to define the local stratigraphy, resulting in a complex and largely informal lithostratigraphic terminology which includes Zanclean lower “Blue Clays” and Zanclean-Piacenzian upper “Blue Clays” (Capezzuoli et al., 2015), the latter eventually further separated by the widespread occurrence of Piacenzian carbonates (Nalin et al., 2016). The lower Zanclean (OOB: Ghinassi, 2007; Nalin et al., 2010; Dominici et al., 2012) and the Piacenzian, are characterised by the high-frequency alternation of coarse-grained and fine-grained facies, ranging from fluvial to marine shelf settings (Benvenuti et al., 1995, 2007, 2014; Martini et al., 2011, Fig. 2). The dynamics of the Pliocene infilling are better-understood in the EB, where six synthemms have been defined, each up to more than 200 m-thick, further subdivided in a number of elementary and composite depositional sequences and chronologically calibrated through marine biostratigraphy and continental vertebrate biochronology (Benvenuti & Del Conte, 2013; Benvenuti et al., 2014, with references).

2. Materials and methods

Stratigraphic sections were measured and described at several localities (Fig. 1). Siliciclastic and carbonate facies were described, subdivided into groups of facies based on lithology, sedimentary structures and ichnology, and interpreted in terms of process and depositional environment (Tab. 1). Each group represents a set of individual facies forming monogenic associations (in the sense of Mutti et al., 1994), i.e., the meter-scale stacking of facies which express the autocyclic behaviour of specific depositional systems within a given

accommodation space (Benvenuti ~~&and~~ Del Conte^{et al.}, 2013). Sequence stratigraphic concepts have been applied to reconstruct the dynamics of basin infills at a hierarchy of scales, advancing hypotheses on controlling factors. The chronostratigraphic subdivision of Benvenuti et al. (2014), which divides the Pliocene into six synthem, S1-S6 from older to younger, was extended to all six Tuscan basins by referring to available biostratigraphic schemes (Fig. 2). The sequence stratigraphic interpretation of S2 in OOB is based on Tinelli (2013). Other parts of the S1-S3 succession were drawn based on available lithostratigraphic literature (see below). Studies integrating sedimentary facies analysis, biostratigraphy and magnetostratigraphy (Nalin et al.^{et al.}, 2016) have helped correlating ~~several discontinuous~~ carbonate bodies typical of S4. The reader is referred to Benvenuti et al. (2007~~;7;~~ 2014) for details on facies analysis and sequence stratigraphic interpretation of synthem S3-S6.

First, data on the geographic distribution of fossil marine mammals~~, -and~~ large sharks ~~and~~ sirenians were ~~collected largely based on collections housed~~ at the Natural History museums of the University of Florence (UFMSN), University of Pisa (UPMSN), and Accademia de' Fisiocritici of Siena (AFMSN), the three largest collections of Tuscany, ~~and at the Geological Museum Giovanni Capellini, Bologna University (MGGC)~~. In particular, counts of cetaceans were based on UFMSN collections (Mysticeti and Odontoceti, ~~n~~N = 142), ~~and those of~~ large shark on the sum of UFMSN, UPMSN and AFMSN collections (Carcharhiniformes and Lamniformes, ~~n~~N = 337, data synthesized~~sed~~ from Marsili, 2006), sirenians from all reports in Tab. 2 (N = 10). Each record is formed either by a single element (e.g., whale bone, shark tooth), by a few elements of the same individual, or by a whole, quasi-articulated skeletons. A large proportion of this dataset lacks precise location, allowing only for some crude stratigraphic attribution (Fig. 3).

On a second step, all fossil Tuscan Pliocene cetaceans ~~and other marine mammals (Sirenia, Carnivora)~~, sirenians and pinnipeds that could be framed within the available high-resolution stratigraphic framework and associated with taphonomic data, were selected. At this step,

after excluding ~~isolated and smaller~~ unidentified MM remains, a dataset of ~~39 cases~~ 64 specimens (cetaceans $n = 3250$; ~~dugongs~~ sirenians $n = 410$; pinnipeds $n = 24$) was assembled. ~~Many of them were a~~ Associationed with shark teeth is frequent (55% of 25 cases according to Danise ~~&and~~ Dominici, 2014 for the Italian Pliocene; see also Bianucci et al., 2002, 2010). The majority of the ~~39~~ 64 specimens are ~~part of~~ included in the catalogue of UFMSN, UPMSN, ~~and~~ AFMSN and MGGC, whereas a few are stored in smaller collections of the municipalities of Montaione, Scandicci (Florence province), and Certaldo (Pisa province), ~~and one in a private property (Castello di Villa Banfi, near Montalcino, Siena province), and one in the Museum National d'Histoire Naturelle in Paris (France).~~ ~~Each one of the 39 LMVs~~ Whenever possible, large marine vertebrates were coded by synthem ($N = 60$) and depositional environment ($N = 54$). We analysed abundance distributions among marine mammals, and species richness of marine mammals and sharks. To infer Pliocene paleoecology, fossil taxa recognised in Tuscany were compared with their closest descendants, focusing on the species today living in the North-Western Mediterranean Sea (NWMS).

~~Since a~~ All known ~~LMV~~ MM-bearing sedimentary facies are associated with a mollusc-dominated benthic fauna. ~~T,~~ the third step of the analysis concerned a quantitative study of shell beds, allowing: 1) to interpret the regional evolution from a perspective independent from the sedimentary facies, ~~and~~ 2) to ~~characterize~~ re-ise the ~~MM~~ paleoenvironmental and bathymetric distribution ~~of LMVs. Molluscs are a key component of Pliocene communities,~~ and 3) allowing to explore the structure of the benthic component of marine ecosystems and to identify underlying environmental ~~factors~~ controls ~~that underline the paleoenvironmental distribution of LMVs.~~ 72 Bulk samples were collected at major shell beds at bed resolution throughout the succession and sieved with 1 mm mesh size. Fossils of bivalves, gastropods and scaphopods were identified to species level. The minimum number of individuals was calculated following standard approaches (see Patzkowsky ~~&and~~ Holland, 2012), resulting

in a richness of 525 species (S) and a total abundance of 64206 individuals (N). We coded each fossil assemblage by synthem, tract of small-scale depositional sequence, and depositional environment. Most samples belonged to facies types F2-F5 (Tab. 1; see also Tomašových et al., 2014). Facies F6 usually lacks macrofossils and allowed for the collection of only one sample. No samples were collected in facies F1, lacking marine shells, facies F7, mostly devoid shells, and facies F8, which is richly fossiliferous, but lacks aragonite shells and is associated with specimens hardly extractable from the rock. The resulting quantitative dataset served for statistical analyses on the distribution of species-level abundances on a siliciclastic shelf depositional system, performed with the software Primer 6.0 (Clarke and Gorley, 2006). Analyses included clustering and nMDS ordination techniques on a Bray-Curtis similarity matrix, of standardised, square-root transformed data (72 samples; S = 333 and N = 63518 after the exclusion of singletons). To test statistically whether there is a significant difference between two or more groups of sampling units based on sedimentary facies, we performed an analysis of similarity (ANOSIM). To interpret the outcomes of the quantitative study and the significance of clusters we used species-level autoecologic information available for the most abundant species, based on the distribution of extant forms. This information, retrieved from the Marine Biodiversity and Ecosystem Functioning EU website (MARBEF: www.marbef.org), included the average life depth of 23 modern species that in our dataset had an overall abundance > 0.15%.

3. Results

3.1. Unconformity-bounded units

3.1.1 Synthem S1: the early Zanclean transgression

The Miocene-Pliocene transition, marking the return to marine conditions after the Messinian salinity crisis (Krijgsman et al., 1999), is recorded in limited exposures of earliest Zanclean, open marine mudstones resting both unconformably or conformably onto latest Messinian non-marine deposits (*Lago-Mare*), an isochronous boundary being dated in the Mediterranean at 5.33 Ma (Roveri et al., 2014). Differences depend on the specific structurally-controlled distribution of hinterland basins, where an uplifting chain determined the presence of thresholds delaying the early Zanclean marine flooding from inner (EB, OOB: Benvenuti et al., 2015a) to outer hinterland basins (FB, VEB). In inner basins such as EB, where S1 has been defined, continental deposition continued into the earliest Zanclean, marine flooding occurring within the MPL1 biozone (references in Benvenuti et al., 2015a). A chronostratigraphical equivalent of S1 is represented in FB and VEB (Bossio et al., 1978) by an open-shelf mudstone, conformably resting on latest Messinian *Lago-Mare* deposits (Roveri et al., 2014). An apparently analogous situation is documented in OOB, where MPL1 shelfal mudstones of the early Zanclean (normal chron C3n) rest on a Messinian to basal Pliocene paleovalley fill (Benvenuti et al., 2015a).

3.1.2 Synthem S2: Zanclean differential preservation

Synthem S2, as recognized in the EB (Benvenuti et al. 2014), is represented by relatively thin fluvial conglomerates unconformably resting on S1 (biozones MPL1-MPL2), capped by S3 (biozone MPL4a: Bossio et al., 1993; 2001), comprising important stratigraphic gaps at its base and top. On the other hand, in FB (Bossio et al., 1997), VEB (Bossio et al., 1994) and SRB (Ghinassi & Lazzarotto, 2005), the same chronostratigraphic interval is recorded by monotonous epibathyal mudstones several hundred meters thick, locally intercalated with delta-front hyperpycnal sandstones and conglomerates. In a very broad

sense, the concept of S2 is extended to these basins, by assuming that erosional unconformities in EB pass to into correlative conformities in rapidly subsiding adjacent basins, where thick successions could be accomodated. Apart from exceptions, no shells were found in bathyal mudstone or in deltaic sandstone. In OOB, the same time span is marked by a N-S facies gradient characterized ~~sed~~ by a single deepening-upward succession, from fluvio-deltaic sandstone to shelfal mudstone, replaced by a succession made of four distinct regressive-transgressive units in the Orcia valley to the north (Ghinassi, 2005; Benvenuti et al., 2015b). A laterally-continuous shell bed, with sharks remains and skeletons and articulated bones of whales, ~~lugongsirenians~~ and ~~large bonyteleost~~ fishes (Danise, 2010; Sorbi et al., 2012; Tinelli, 2013), marks a major transgressive surface overlain by open shelf mudstones (Sorbi et al., 2012; Tinelli, 2013: biozone MPL2) that is hypothetically traced along a NE-SW profile (Figs. 4, 5).

3.1.3 Synthem S3: Zanclean-Piacenzian transition

Synthem S3 is subdivided in EB into a lower and an upper interval (Benvenuti et al., 2014). The lower division is represented by deepening-upward, coarse-grained delta front system, overlain by an upper mudstone division from an open shelf setting. The upper part is rich with shell beds, and occasional articulated whale skeletons, associated with shark teeth, have been recovered (Danise ~~&and~~ Dominici, 2014). Similar shelf mudstones of the MPL4 biozone crop out in VEB and SRB (Bossio et al., 1993; Riforgiato et al., 2005) whereas biostratigraphically equivalent mudstones in FB testify to an upper epibathyal paleoenvironment.

3.1.4 Synthem S4: early Piacenzian warm climate and high sea-level

Synthem S4 (Benvenuti et al., 2014) has been recognized in FB and VEB by facies similarities and chronostratigraphic correlation. In EB, S4 comprises a lower interval dominated by richly fossiliferous, massive mudstone or very-fine-grained sandstone (prodelta-inner shelf), overlain by bioclast-rich sandstones recording prograding mixed carbonate-clastic ramp, outcropping in the southeastern part of the Elsa valley. Equivalent deposits, also comprised in biozone MPL4b, are patchily distributed in FB, VEB, SRB, OOB and other basins of southern Tuscany (Ghinassi & Nalin, 2010; Ghinassi, personal communication, 2015). The upper interval of S4 is formed by a succession of delta front sandstones, passing in EB eastern margin to a few tens of m-thick fluvial succession, hinting at an original depositional gradient. S4 is apparently missing due to erosion north of San Gimignano (EB), and around Lajatico (VEB). Biostratigraphic data allow to refer S4 to the upper part of biozone MPL4b and the lower part of MPL5a, thus comprising the mid-lower part of the Piacenzian, globally characterized between 3.264-3.025 Ma by warm climate and relatively high sea level (Raymo et al., 2009; Dowsett et al., 2013; Prista et al., 2015).

3.1.5 Synthem S5: mid-Piacenzian high-frequency sea level variation

Synthem S5, recognized in EB and VEB, is bounded below by an erosional unconformity that cuts deeply into underlying units, bringing S5 directly on top of S3 (EB: log 15; VEB: logs 5-6 in Fig. 1). S5 is up to about 200 m in EB, where it has been subdivided into a hierarchy of small-scale depositional sequences (Benvenuti et al., 2007; Dominici et al., 2008: see following paragraphs). Each composite depositional sequence forms a tens-of-m-thick asymmetric sedimentary cycle, composed by a deepening-upward part, from fluvial or coastal coarse-grained sediments, to open shelf mudstones, sometimes topped by a regressive shoreface or delta sandstone, other times directly overlain by the next sequence through a sharp contact. Fluvial, lagoonal brackish-water, and other intertidal deposits mark

the lower part of each composite sequence, usually topped by a laterally-continuous shell bed, from a few cm to a few dm-thick, representing a surface of transgression. Shell beds are particularly well-developed around the middle part of sequences, where they separate shoreface and delta sandstones from overlying open shelf mudstones, marking the time of maximum flooding (MFS). Large marine vertebrates, including articulated whale skeletons and large sharks (Danise ~~&and~~ Dominici, 2014), are often recovered both at MFS and overlying mudstone (Fig. 1). Towards the north-eastern margin of EB, cyclothemic fluvial conglomerates, sandstones and mudstones replace coastal and fully marine deposits, testifying to an original facies gradient. In the central part of EB, composite sequences are stacked to form a deepening-upward succession, with a topmost thick and laterally-continuous open shelf mudstone interval, directly onlapping the S4-S5 basal unconformity on the eastern EB (log 15, Fiano: Fig. 1).

3.1.6 Synthem S6: Piacenzian-Gelasian climate change and regression

As S5, synthem S6 is also built through a hierarchy of small-scale depositional sequences, better expressed in EB, but also documented in SRB and CB. In EB, fluvial coarse-grained sandstones fill a deep valley incised in S5 deposits (logs 8-10 in Fig. 1), resting on the basal unconformity of S6. Intertidal or coastal lagoon deposits form the transgressive systems tract of the composite depositional sequence. A laterally-continuous shell bed testifies to the MFS of S6, topped by highstand shoreface and delta front sandstones. The Piacenzian-Gelasian boundary, corresponding to a major climatic transition from warmer-moister to colder-drier conditions (Benvenuti et al., 1995b; 2007), is marked by the Gauss-Matuyama reversal detected at Montopoli (Lindsay et al., 1980). Highstand marine sandstones are characterized by the recovery of two mysticete skeletons, at Montopoli (EB, see Capellini, 1905) and Sinalunga (CB, Fig. 7). Fully continental environments were established

throughout the Gelasian in all basins here under study (e.g., Benvenuti [& Del Conte](#), 2013; Benvenuti et al., 2014; Bianchi et al., 2015).

3.2 Sedimentary facies and facies associations

Seven groups of siliciclastic facies and one group of carbonate facies, with very different fossil content, have been recognized (Tab. 1). Siliciclastic facies form a paleoenvironmental gradient from terrestrial to marine and, in the case of marine facies, from shallowest to deepest (Fig. 8a). Facies types are fluvial conglomerate and sandstone, and alluvial mudstone (F1); intertidal to very shallow subtidal mudstone and muddy sandstone (F2); shallow subtidal coarse- and medium-grained sandstone with sparse conglomerate (F3); deep subtidal muddy fine-grained sandstone (F4); open shelf sandy mudstone (F5); outer shelf to upper bathyal mudstone (F6); outer shelf and bathyal turbidite sandstone and conglomerate (F7). An eighth group is formed by facies deposited subtidally in limited mixed carbonatic-siliciclastic ramps (F8), irrespective of depth (Nalin et al., 2016). Facies F6-F7 are restricted to Zanclean deposits (synthems S1-S3); facies F5 and F8 characterise the upper Zanclean-middle Piacenzian interval (synthems S3-S4); facies F1-F4 characterise the upper Piacenzian (synthems S5-S6). The OOB succession, characterised by facies F1-F4, is an exception within the Zanclean.

3.3 Elementary depositional sequences (EDS)

Both fluvial (Benvenuti [& Del Conte](#), 2013) and marine facies groups (siliciclastics: Benvenuti et al., 2007; 2014; carbonates: Nalin et al., 2016) are stacked to form facies associations which record cyclic variations of depositional and environmental conditions in response to a change in accommodation space. Physical surfaces and the intervening

deposits allowed to subdivide depositional sequences in systems tracts (Benvenuti et al., 2007; Dominici et al., 2008). At the simplest scale, these hybrid facies associations form elementary depositional sequences, up to 10-20 m thick, in their turn stacked to form composite sequences (original concepts from Mutti et al., 1994). This hierarchy is particularly evident in synthem S5-S6, formed at a time of pronounced glacio-eustatic oscillations and expressed around coastal settings, where maximum facies contrast allows for the expression of subtle cycle of sea level variation (e.g., Benvenuti ~~&and~~ Dominici, 1992; Benvenuti et al., 2007; Dominici et al., 2008). Analogue sharp facies contrast within Zanclean EDS in OBB (Tinelli, 2013), but is otherwise absent in deeper sediments (facies F5-F7). EDSs have different expressions depending on the time interval and the sedimentary basin.

3.3.1 Zanclean EDS (synthems S1-S3)

In most basins, deposition of synthems S1-S2-S3 takes place at outer-shelf or bathyal depths, well below the point on a depositional profile where the rate of relative sea level change is zero (equilibrium point). Here the sediment supply is not sufficient to fill the available accommodation space and an aggradational style of deposition prevails, with the result that in most Zanclean settings smaller cycles of sea level variations are not marked by a facies change. The sharp facies change recorded where the monotonous muddy deposition is interrupted by turbidite sandstone and conglomerate, is connected with synthem boundaries and major tectonic phases of restructuring of the region. ~~LMVs~~MM and shell beds are practically absent. This situation reverses in the Orcia-Ombrone basin, where depths of deposition are shallower and EDS are expressed. At Arcille deltaic sandy conglomerates and sandstones (facies F3) are overlain by a fluvial cross-bedded sandstone (facies F1a), separated by a transgressive surface (TS) from an overlying bioturbated shallow marine sandstone. A *Haustator vermicularis* shell bed (Danise, 2010; Tinelli, 2013)

forms the MFS separating the shoreface sandstone from an open marine mudstone with scattered shells (facies F5), marking a sudden and prolonged deepening of the basin (biozone MPL2: Sorbi et al., 2012; Tinelli, 2013). The succession is topped by deposits from shallower depths, expression of the falling-stage (FSST), below the upper SB (Fig. 4). Similar small-scale depositional sequences, expressed through fining-upward cycles no less than 40 m-thick, are also present in synthem S3 at Case al Poggio, near Siena (biozones MPL3-MPL4a: Bianucci et al., 2001) and at Castelfiorentino (biozone MPL4b: Benvenuti et al., 2014).

3.3.2 Piacenzian EDS (synthems S4-S6)

As depth of deposition shallows during the Piacenzian, and cycles of sea level variation widen, the cyclic stacking of EDS becomes the typical depositional theme (Benvenuti et al., 2007, 2014; Dominici et al., 2008). In FB, the northwesternmost basin, depths remain considerable and facies change is more subtle. Pliocene at Orciano Pisano is traditionally assigned to the “Blue Clays” formation (Bossio et al., 1997), but two distinct bodies were evidenced since the late nineteenth century (D’Ancona, 1867). The lower one is formed by grey claystone with very rare shells, the upper one by muddy, very-fine grained gray sandstone richly fossiliferous. The lower part of the latter interval outcrops at the foothill of the small town of Orciano Pisano, around the locality Case Nuove (Bianucci & Landini, 2005; Berta et al., 2015). Here a laterally persistent shell bed is dominated by the turritellid *Archimediella spirata*, overlain by a 25 m-thick monotonous sandstone interval with intercalated shell beds or sparse shells, becoming muddier upward. The *Archimediella* shell bed is interpreted as the TS of an EDS, coinciding with the lower SB of a lower Piacenzian EDS. The overlying muddy fine-grained sandstone is the TST and HST (MPL5a: Dominici et al., 2009; Fig. 6). At shallower settings, in all basins to the East and South of FB, EDS of synthem S4 take the form of an alternation of mudstone and carbonate (facies F5 and F8:

Nalin et al., 2016), or mudstone and sandstone (facies F5 and F3-F4: Benvenuti et al., 2014; biozone MPL5a). Middle and upper Piacenzian EDS form and alternation of coastal mudstone and sandstone (facies F1-F2-F3: Benvenuti ~~&and~~ Dominici, 1992; Dominici, 1994), with MFS and HST marked by a shell bed topped by a lower shoreface sandstone, or a shelf mudstone (facies F4-F5: Benvenuti et al., 2007; Dominici et al., 2008).

43. Distribution of large marine vertebrates

3.4.1 Geographic distribution

The MM geographic distribution ~~of all LMVs that could be assigned to a given locality and sedimentary facies~~ is listed in Table 2. All LMVsMM that ~~werewas~~ geographically located, irrespective of stratigraphy, ~~werewas~~ plotted in Fig. 3. The largest number were recovered in FB in Northwestern Tuscany, with a peak at a few sites around the small town of Orciano Pisano, in the Pisa province. This coincides with the highest number of known species, including mammals, elasmobranchs, turtles, and large bony fishes ~~(see Bianucci & Landini 2005, Marsili 2006)~~. LMVMM ~~abundance is also high~~ is also abundant in the province of Siena, around Volterra (VEB; Bianucci ~~&and~~ Landini 2005), and around San Quirico, particularly rich with elasmobranchs according to the available data (SRB; Marsili, 2006). A fourth basin with a consistent number of findings is OOB, where cetaceans, sharks, ~~dugongsirenians~~ and large ~~bonyteleost~~ fishes have been unearthed.

3.4.2 SMM stratigraphic distribution ~~of LMVs~~

3.4.2.1 Synthems S1-S2

In the vicinity of Saline di Volterra, ~~provenance of a mandible with teeth of one of the sites~~ with remains of *Pliophoca etrusca* (Berta et al., 2015) and sperm whale (Tab. 2), S1 is represented by a bathyal mudstone (facies F6, biozone MPL1). Lower-middle Zanclean ~~LMVsMM are~~ is otherwise absent, with the exception of S2 in OOB, where ~~they~~ large marine vertebrates are concentrated ~~along the MFS at the maximum flooding interval~~, outcropping at Poggio alle Mura (Danise, 2010), Camigliano (Sorbi et al., 2012) and Arcille (Tinelli et al., 2012; Tinelli, 2013), in biozone MPL2. At Poggio alle Mura a slightly disarticulated ~~balaenopterid~~ balaenid whale skeleton (WOM1-1 in Tab. 2) has been excavated in contact with the laterally-persistent *Haustator* shell bed at the maximum flooding interval (Figs. 4, 5; Danise, 2010; Tinelli, 2013). At Camigliano and Arcille, tens of km from Poggio alle Mura, the *Haustator* shell bed is associated with other articulated skeletons and isolated ~~LMVMM~~ remains, including several specimens of the sirenian *Metaxytherium subapenninum* (MOM21-1, MOM1-2-5), large bony fishes, rays, and sharks *Carcharias taurus*, *Carcharhinus* sp., *Galeocerdo cuvieri*, and *Squatina* sp. sharks and rays (Sorbi et al., 2012; Tinelli et al., 2012; Tinelli, 2013). A partial skull of the delphinid *Etruridelphis giulii*, with right and left dentaries fractured, but nearly complete and with most teeth still in their alveoli (DSi2-1 in Tab. 2), was recovered southeast of Siena, near Chianciano Terme (SRB) at the top of a mudstone (facies F6) intercalated with turbiditic sandstone beds (Facies F7), topped by a monotonous mudstone interval (upper Zanclean, uppermost part of biozone MPL3: Bianucci et al., 2009b). In the same basin, two undetermined beaked whales and bones of four different specimens attributed to *Metaxytherium subapenninum* were found in the middle of a fining-upward succession, at the boundary between biozones MPL3 and MPL4a (Bianucci et al., 2001; WSi2-1-2 and MSi2-1-4 in Tab. 2), suggesting that findings at the two SRB localities belong to the same MFS, in the upper part of S2. ~~Four~~ Three specimens of marine mammals were identified in Synthem S1, ~~13~~ seven in Synthem S2.

3.4.2.2 Synthem S3

A partial delphinid skull, with right and left dentaries fractured, but nearly complete and with most teeth still in their alveoli (WSi2-1 in Tab. 2), was recovered southeast of Siena, near Chianciano Terme (SRB) at the top of a mudstone (facies F6) intercalated with turbiditic sandstone beds (Facies F7), topped by a monotonous mudstone interval (upper Zanclean, uppermost part of biozone MPL3: Bianucci et al., 2009). In the same basin, a sirenian skull and mandible were found in the middle of a fining-upward succession, also at the boundary between biozones MPL3 and MPL4a (Bianucci et al., 2001; MSi3-2 in Tab. 2), suggesting that these two findings belong to the same MFS, in the lower part of S3.

The upper part of ~~the latter synthem S3~~ yielded several other LMVs, including an articulated and well-preserved balenopterid skeleton found in a 30 m-thick mudstone succession at Castelfiorentino, in the Elsa basin (EB, WEI3-12). These strata are richly fossiliferous, with several shell beds with epifaunal cemented taxa, such as vermetid gastropods, oysters and corals (Facies F5), within a normal-polarity magnetostratigraphic interval (Gauss chron: Andrea Albianelli, personal communication, 1999). Based on physical stratigraphic correlation, this can be assigned to a lower Piacenzian HST. ~~Finally, aA dolphin skull of~~ Hemisyntachelus sp. (Aldinucci et al., 2011) and an incomplete, almost articulated partly articulated vertebrae and costae of a dolphin skeleton (Arbeid et al., 2015) were recently excavated a few hundred meters apart one from the other, along a monotonous mudstone S3 succession intercalated with several *Ostrea* and *Serpulorbis* shell beds, near Certaldo (facies F5, DWEI3-1-2 in Tab. 24). Overall, specimens of marine mammals attributed to Synthem S3 were sixeleven (some are uncertain and may come from the upper part of S2: Tab. 2).

3.4.2.3 Synthem S4

Synthem S4 yielded the highest ~~number of the reviewed large marine vertebrates~~abundance ~~and species-richness of the Tuscan MM~~. In the Fine basin, the locality of Orciano Pisano is represented in Table 2 by ~~nine~~18 records of whales, dolphins (a partial skeleton: Bianucci, 1996; Bianucci et al., 2009), seals (Berta et al., 2015), tens of other ~~cetacean specimens~~unidentified cetacean elements, hundreds of shark teeth and vertebrae (Fig. 3), and sea turtles, large bony fishes, and sea birds hosted in museum collections (Bianucci ~~& Landini, 2005; Marsili, 2007b; Cioppi and Dominici, 2011~~). In locality Case Nuove, a single middle Piacenzian ~~T~~Transgressive surface is known to have~~has~~ yielded a whole, articulated whale skeleton (Figs. 4, 6), teeth of blue and white sharks and bones of sea birds (Dominici et al., 2009). From the same site come ~~ss a n-articulated seal~~of the monk seal *Pliophoca etrusca* (Berta et al., 2015~~4~~), and possibly many other museum specimens labelled "Orciano Pisano" ~~(e.g., Higgs et al., 2012)~~, suggesting that this ~~surface~~interval ~~is forms~~ the most prolific bonebed of the region (Tab. 2). A few cm above the *Archimediella* shell bed, glauconitic and deeply bioeroded whale bones (Danise, 2010), associated with ichnological evidence of the activity of *Osedax* bone-eating worms (see Higgs et al., 2012), lie in a bioturbated muddy, fine-grained sandstone with a complex boxwork of *Ophiomorpha* and *Thalassinoides* trace fossils (Fig. 6), associated with a diverse paleocommunity of molluscs and other benthic invertebrates ~~from several trophic groups with complex trophic connections~~ (Dominici et al., 2009; Danise et al., 2010). ~~In the adjacent~~At the boundary between FB and VEB, an incomplete ~~dolphin~~ skull and skeleton of the dolphin *Etruridelphis giulii* was recovered near Lorenzana (Lawley, 1876; Bianucci, 1996; Bianucci et al., 2009~~b~~), at a locality associated with a muddy sandstone interval (facies F4) in synthem S4. An incomplete and ~~disarticulated mysticete~~mysticete ~~balenopterid whale~~ was recovered in a sandstone at San Gimignano, associated with pectinid bivalves (Fig. 7; facies F3; Elsa basin, EB: Danise ~~& Dominici, 2014~~), here tentatively assigned to the uppermost part of the synthem

(WEI4-1 in Tab. 2). Overall, specimens of marine mammals attributed to Synthem S4 were 2728.

3.4.2.4 Synthem S5

The next MM richer stratigraphic interval are Piacenzian deposits of synthem S5. An incomplete and disarticulated balaenid skeleton was found in intertidal deposits of the lower part of the synthem, at La Farfalla Casenuove (facies F2, EB; Bianucci et al., 1995; Collareta et al., 2016; WEI5-1 in Tab. 2). A large ~~bowhead whale~~balaenid (Balaena) was recovered higher up section, a few meters above a laterally-continuous very thick and complex *Haustator vermicularis* shell bed, up to 2 m-thick (Benvenuti et al., 1995a), traced laterally for 2 km to the east of San Miniato ("*Turritella strata*", De Stefani, 1874), and forming a surface of transgression within the TST of ~~synthem~~ S5 (Benvenuti et al., 2007; 2014; Dominici et al., 2008). The MFS is formed by a *Glycymeris insubricus* shell bed, separating around San Miniato shoreface sandstone (facies F3 or F4) from offshore mudstone (F5; Benvenuti et al., 2007). The ~~Balaena sbalaenid~~ skeleton was almost articulated, ~~complete~~ and bioeroded, closely associated with teeth of the great white shark and other scavengers (Borselli ~~&and~~ Cozzini, 1992; Bianucci et al., 2002; Danise and Dominici, 2014), in the early HST of synthem S5 (Benvenuti et al., ~~2007~~; ~~2014~~; Dominici et al., 2008; WEI5-2 in Tab. 2). The *Glycymeris* shell bed can be traced laterally for several km. In the vicinity of Fiano, it includes bioeroded and encrusted gravels inherited from underlying successions, interpreted as revinement deposits. The interval of maximum flooding ~~and early HST~~ is marked by the stacking of at least three distinct shell beds, all including a high-diversity association with bioeroded and encrusted shells. This situation suggests that balaenid WEI5-2, ~~in the middle of EB~~, lies in correspondence of an interval of low rates of sedimentation. In SRB, near

Castelnuovo Berardenga, shelfal mudstones (facies type F5) have yielded MM remains at a few localities. Delphinid remains were found at the “I Sodi” quarry and at Troiola (DSi5-1-2, in Tab. 2). Bones of a beaked whale and undetermined mysticetes are reported from Guistrigona (Manganelli and Benocci, 2014) and a fragmentary specimen of the monk seal *Pliophoca etrusca* from Castelnuovo Berardenga (Berta et al., 2015). A very rich shark fauna, including sawsharks, thresher, frilled, bluntnose sixgill, bramble, gulper, kitten, sand tiger, shorten mako, basking and requiem sharks, an association suggesting an upper slope paleoenvironment for the surroundings of Castelnuovo Berardenga Scalo (Cigala-Fulgosi et al., 2009; the same mudstone interval at the same locality has been interpreted as a shelfal deposit: Martini et al., 2016). Marine mammals identified in Synthem S5 were eight. Judging from historical accounts (Capellini, 1883), the killer whale *Orcinus citonensis* (DCh4-1) was recovered in a sandy unit lying on top of a thick mudstone interval (S3 or S4) and is tentatively assigned to S5. Specimens of marine mammals attributed to Synthem S5 were 10.

3.4.2.5 Synthem S6

A fairly complete skeleton of a large balaenid whale was recovered in the second half of the 19th Century in the EB near Montopoli Valdarno (Capellini 1905), in open shelf strata attributable to the interval of maximum flooding of ~~Synthem~~ S6 (WEI6-1 in Tab. 2). The MFS of S6, of uppermost Piacenzian age, is marked in EB by a laterally continuous *Pteria phalenacea* shell bed, with a high-diversity association of macroinvertebrates, including a rich decapod paleocommunity (Garassino et al., 2012). In the ~~same locality~~ vicinity, near Palaia, a right whale (*Eubalena* sp.) was recovered in 1974 in sandy mudstones, in association with mollusc shells and teeth of the great white shark (*Carcharodon carcharias*: Bisconti, 2002; Sorbini et al., 2014). A tightly articulated balaenopterid skeleton was found

at Sinalunga (WCh4-1 in Tab. 2), in deltaic sandstones and conglomerates (Fig. 8). Marine mammal specimens attributed to Synthetium S6 with some confidence were three.

4.3 MM facies type distribution

In an ideal deepening-up gradient, multi-element findings of marine mammals are very rare in intertidal and very shallow subtidal paleoenvironments (facies type F2, 1,5%: Fig. 9A), moderately represented in delta or shoreface sandstones (facies type F3, 4,6%), most abundant in sandy mudstone of open shelf settings (facies type F5, 87,70%), rare in outer shelf and bathyal sediments (facies type F6-F7, 6,21,5%). The most pristine and complete skeletons are associated with gravelly well-sorted sands from event sedimentation, suggesting a negative relationship between taphonomic loss and sedimentary processes at delta fronts. In the tightly articulated and pristine Sinalunga balaenopterid (WCh4-1 in S4-CB), the cortical surface of the tightly-connected vertebrae is practically intact, and carpal, metacarpal and phalanges of the flipper are in perfect anatomical connection, as if a sudden depositional event buried a fresh carcass (Fig. 8; similar pristine skeleton are found in deep water turbiditic succession: Stinnesbeck et al., 2014). Another pristine and tightly articulated skeleton, belonging to a killer whale (WCh3-1 in CB), was collected at Cetona in the second half of the 19th century, in a locality associated with sandstones, also possibly of deltaic origin. Large vertebrates embedded in fine-grained, muddy matrix (shelf deposit formed below storm wave base) and those associated with laterally-persistent shell beds (condensed deposits) are slightly disarticulated and fairly complete, showing signs of long permanence in a low energy, well-oxygenated seafloor before the final burial. Bioerosion of bones caused by phototrophic cyanobacteria and algae, heterotrophic fungi and bacteria (Orciano balaenopterid WFi4-1 in S4-FB), and eventually by whalebone-eating siboglinid worm of genus *Osedax* (on a ziphiid humerus, WFi4-14: Higgs et al., 2012), occurred at

condensed intervals, in association with glauconite (Danise, 2010). In one instance, a condensed shelly interval is traced for a few kilometers, connecting bioturbated shoreface sandstones yielding slightly disarticulated sirenian skeletons (MOM2-1, MOM2-2) and other MM (Tinelli, 2013), with open shelf deposits yielding slightly disarticulated whale remains (WOM2-1 in S2-OOB).

Regarding a sequence stratigraphic interpretation, pristine skeletons from delta front sediments can be part of the FSST (Fig. 4; or in an alternative interpretation it may belong to, of the early TST, when incised valleys are filled with coastal deposits). TST deposits account for 126,2% of cases (Fig. 9B). Much more frequently, articulated or slightly disarticulated skeletons are associated with the HST of the corresponding depositional sequence, lying above the MFS (6956,9% of cases), or within the maximum flooding interval directly in coincidence with it, above a laterally-persistent shell bed (10,8%). Bone bioerosion is less pronounced in the late HST, when sedimentation rate increases (Castelfiorentino whale WEI3-2 in S3-EB).

4.4 MM abundance and species-richness

Some taxa need revision, but a conservative estimate of the different morphologies suggest that at least 17 marine mammal species lived in the NWMS during the Pliocene (possibly more than 20, an estimate for the whole epoch, i.e., 5,332-2,588 Ma), against only 9 nine presently living in the same area (plus two occasional visitors). Among the cetaceans, six families were present, against only five presently living in the Ligurian Sea (Tab. 3). The most abundant Pliocene species of Tuscany are the sirenian *Metaxytherium subapenninum* (N = 10), the dolphin *Etruridelphis giulii* (N = 7) and the delphinid *Hemisyntrachelus cortesii* (N = 5; Tab. 4). Overall abundance and species richness are not randomly distributed, but are maximum in Piacenzian strata of synthem S4, dated at 3.2-3.0 Ma (Tab. 2, Figs Fig-2,

9C-D2), particularly in the FB and VEB (Tab. 4). A species list of marine mammals summing up findingsfossils found around Orciano Pisano and at La Rocca, near Volterra, yields a species richness (S) of 11-1213. This Piacenzian peak in marine mammal diversity is matched by the fossil record of sharks, also maximum near Orciano Pisano (S = 27), with a second peak in S5, around Castelnuovo Berardenga (SRB, S = 16). Differently from the marine mammals, showing a complete turnover from the Pliocene to the recent (Tab. 3), 34% of Pliocene shark species are still extant in the NWMS (Tabs. 5-6). Marine mammal frequency and diversity gradually decreases in S5-S6.

35. Paleoenvironment-fauna relations

The paleoecology of Pliocene MM of Tuscany can be reconstructed by comparison with the ecology of their modern relatives. This approach can be applied at the family or genus level for marine mammals (Tab. 3), at the genus or species level for sharks (Tab. 5). With the only exception of the sirenian *Metaxytherium subapenninum*, feeding on seagrasses (Domining, 2001), and small demersal sharks (e.g., catsharks, frilled sharks), all MM studied here are pelagic forms that had no direct connection to conditions at the seafloor (Tabs. 4 and 6). ~~With the exception of the few benthopelagic marine vertebrates (seals, dugongs), many LMVs of the Pliocene of Tuscany were epipelagic (e.g., mysticetes, larger sharks) or mesopelagic (e.g., sperm whales, ziphiids), with no direct connection with the ecological conditions at the seafloor.~~ The paleoecology of benthic habitats informs however on the situation of the overlying water column in terms of ~~ecological~~ factors that matter to the distribution of ~~pelagic~~~~epipelagic and mesopelagic~~ organisms. ~~These factors include, such as~~ water depth, salinity and nutrient levels. In the second place, since ~~all LMV~~~~all MM~~ remains after death ultimately sink to the seafloor, benthic paleoecology is also a means to

understand taphonomic controls on LMVMM distribution. Cluster analysis based on the distribution of mollusc species in 72 samples resulted in the identification of four main groups of ~~mollusc~~-samples, roughly corresponding to the four main facies types recognized ~~sed~~ based on lithology and sedimentary structures (F2-F5 in Tab. 1). Clusters are formed by samples from outer shelf and upper slope (three samples), open shelf (34 samples), shoreface (23 samples), and transitional settings, such as brackish-water coastal lagoons and tidal flats (12 samples: see Supplement Material, Fig. S1). ANOSIM confirms that sedimentary facies type can broadly predict what benthic assemblage it will yield (Tab. 73; general $R = 0,632$). The difficulty to discriminate between upper and lower shoreface facies, and between shoreface and open shelf facies is confirmed by overlaps in sample distribution in the NMDS ordination diagram (Fig. 109A). Samples AG1, MON1 and MON2 allow to re-interpret the associated sandstones, originally included in upper shoreface facies type, as offshore deposits. The presence of gravels and cobbles intensely bioeroded by bivalves and polychaete (ichnofossils include *Gastrochaenolithes*, *Meandripolydora* and *Caulostrepsis*) and encrusted by balanids, oysters, serpulids, and bryozoans, suggests they are part of condensed beds resulting from transgressive pulses (hiatal concentrations). The relationship between facies type and mollusc association is broadly summarized in the following paragraphs (see online Supplement Information for a list of characterizing species).

35.1 Coastal lagoon, tidal flat and embayment

Intertidal faunas ~~are~~ always associated with facies type F2 and are characterised by low-diversity associations, sometimes with less than 10 taxa and dominated by one or two species, including species today living in brackish waters of the Mediterranean, at intertidal or very shallow subtidal depths (e.g. *Cerastoderma edule*, *Nassarius reticulatus*,

Scrobicularia plana: Pérès ~~&and~~ Picard, 1964). Facies type F2 is also associated with samples having a species richness higher than the preceding and including species typical of seagrass bottoms and known to withstand moderate changes of salinity. In only one instance a ~~LMV~~large vertebrate was associated with intertidal deposits. ~~Interestingly, this was (not a sirenian, but a balaenid, lying on top of large wood fragments-(:~~ Bianucci, 1995; Collareta et al., 2016; Fig. 98A).

35.2 Upper shoreface

Facies type F3 is associated with a high-diversity assemblage representing a paleocommunity dominated by suspension feeders adapted to shifting sandy bottoms, with bivalves typical of modern shoreface sandy bottoms (e.g., families Glycymeridae, Tellinidae, Donacidae and Veneridae). Among extinct species of this recurring assemblage, some are large-sized or have very thick shells. Some species of this group indicate the presence of vegetated bottoms. Small pyramidellid gastropods are parasitic on echinoderms, also typical of the upper shoreface.

35.3 Lower shoreface

Species richness further increases in collections associated with facies type F4 (lower shoreface). Species typical of this recurring assemblage include both suspension-feeding and detritus-feeding bivalves and gastropods. The following gastropod families are usually represented by several species: Trochidae, Rissoidae, Cerithiidae (from vegetated bottoms), Naticidae, Muricidae, Turridae, Conidae, Terebridae, Bullidae, Cylichnidae (carnivores), Pyramidellidae (echinoderm parasites). Many bivalve species occur in both facies types F2 and F3. At three different sites and at different stratigraphic units ~~LMVs~~

werelarge marine vertebrates, including mysticetes, sirenians and sharks, were recovered in association with shell beds dominated by the ~~turritellid and~~ gregarious turritellid gastropod *Haustator vermicularis* (Fig. 6C-D).

35.4 Offshore and upper slope

Another important set of species recurred in facies type F5 (mudstone deposited in offshore bottoms at shelf depths). Among characterizing gastropods are the suspension feeders (*Turritella tricarinata*, *Archimediella spirata* and *Petalochoncus intortus*), deposit feeders (*Aporrhais uttingeriana*) and carnivores or scavengers (*Epitonium frondiculoides*, *Nassarius semistriatus*, *Mitrella nassoides*). Also the bivalves occupy many different ecological niches (e.g., infaunal detritus feeders, epifaunal suspension feeders, either free-living, byssate, or cemented). Outer shelf and uppermost bathyal ~~samplessediments~~ from F6 mudstones, studied at only one location, are characterized ~~sed~~ by a separate set of carnivorous gastropods and by a few small bivalve species. Ubiquitous molluscs include species found from intertidal to outer shelf depths (e.g., *Corbula gibba*), and those preferential of open marine waters, from lower shoreface to outer shelf. Most multi-element findings of marine mammals are associated with sandy mudstones from open shelf settings, below storm wave base (~~F5, 75-80% of occurrences; e.g., Orciano whale: Fig. 8A~~), also in association with an *Archimediella spirata* shell bed (Fig. 6C-D). No molluscs were recovered in facies type F7, with the exception of bathyal mudstone in the lowermost Pliocene of FB, associated with sparse specimens of the gryphaeid epifaunal bivalve *Neopycnodonte navicularis* -(not sampled).

35.5 Carbonate platform

A recurring benthic assemblage associated with the highly fossiliferous facies type F8, not included in the quantitative analysis, consists of the large pectinid bivalves *Gigantopecten latissima* and *Hinnites crispus* and by a mixture of photozoans (large benthic foraminifera), hard-substrata dwellers (brachiopods, echinoderms), encrusters (red algae, bryozoa) and bioeroders (clionid sponges). All fossil-rich carbonates are associated with synthem S4 , in the middle part of the Piacenzian (Fig. 2, see also Nalin et al., 2016).

46. Paleodepths

Multivariate techniques ~~can be~~are usefully applied to stratigraphic and paleobiologic analysis (Scarponi & Kowalewski, 2004). We used the results of the ordination analysis to estimate absolute depths of the final resting place of some ~~LMV~~large vertebrates listed in Tab. 2. Samples in the NMDS ordination plot following a water depth gradient, with shallower samples to the left (low values of NMDS axis 1) and deeper samples to the right (high values of axis 1). Therefore, NMDS values of axis 1 can be used as a proxy for relative water depth. We calculated absolute palaeodepths by fitting a logarithmic regression curve between absolute paleodepth of 23 modern species (data from MARBEF database), common in our dataset, and the values on NMDS axis 1 (Supplement Material: Tabs. S1-S4). The regression analysis, with $R^2 = 0.813$, indicates that scores along the nMDS main axis are a good predictor of the preferred depth for the 23 modern species (Fig. 910B, inset), thus supporting the bathymetric interpretation. This allowed to estimate the absolute depth of the 72 ~~mollusc~~ samples, which ranged from 0.4 m to 365 m, i.e., from intertidal to upper slope depths (Fig. 109B). ~~These results are,~~ in accordance with a previous estimate of absolute paleodepths, ~~carried out~~ in the upper part of the Pliocene ~~succession~~ of EB, ~~and~~ based on counts of foraminifera. ~~(Dominici et al., 2007). According to the present estimates,~~ F Facies type F2 is deposited at 0-105 m depth ~~(mostly 0-5 m)~~, F3 at 3-30 m, F4 at 10-100

m, and F5 at 40-300 m, F5 at around 350 m depth (only one sample: Fig. 910B). The position of fossil cetaceans, dugongsirenians, pinnipeds and large sharks (respectively W, M, P and S, N = 13) was plotted near the corresponding shell bed in the NMDS ordination. The resulting pattern shows that all large mammalsMM considered, associated with open shelf settings and with facies types F4-F5, cluster around a depth of about 100 m, spanning 30-300 m (Fig. 9B).

5. Interpretation

~~Large marine vertebrates are nearly unrecorded in synthem S1-2 (Zanclean, mostly bathyal mudstone and turbiditic sandstone), notwithstanding these successions are the thickest, but they are recovered in S2 in OOB (shoreface and open shelf deposits). LMVs are abundant in synthems S3-5, peaking in synthem S4 (middle part of the Piacenzian), with intermediate abundances in synthem S6 (upper Piacenzian-lowermost Gelasian). In an ideal deepening-up gradient, multi-element findings of marine mammals are very rare in intertidal and very shallow subtidal paleoenvironments (facies type F2: one instance in Bianucci, 1995, out of 39 cases explored), moderately represented in delta or shoreface sandstones (facies type F3), most abundant in sandy mudstone of open shelf settings (facies type F5, 77% of occurrences: Fig. 8A; e.g., Orciano whale: Figs. 4, 6), nearly absent in outer shelf and bathyal sediments (Saline di Volterra seal, PER1-1 in S1-VB; Lucciolabella dolphin, WSi2-1 in S3-SB). Regarding their taphonomy, the most pristine and complete skeletons are associated with gravelly well-sorted sands from event sedimentation, suggesting a negative relationship between taphonomic loss and sedimentary processes at delta fronts. In the tightly articulated and pristine Sinalunga balaenopterid (WCh4-1 in S4-CB), the cortical surface of the tightly-connected vertebrae is practically intact, and carpal, metacarpal and phalanges of the flipper are in perfect anatomical connection, as if a sudden depositional event buried a fresh carcass (Fig. 7; similar pristine skeleton are found in deep water~~

turbiditic succession: Stinnesbeck et al. 2014). Another pristine and tightly articulated skeleton, belonging to a killer whale (WCh3-1 in S4-CB), was collected at Cetona in the second half of the 19th century, in a locality associated with sandstones, also possibly of deltaic origin. LMVs embedded in a fine-grained, muddy matrix (shelf deposit formed below storm wave base) and those associated with laterally-persistent shell beds (condensed deposits) are slightly disarticulated and fairly complete, showing signs of long permanence in a low energy, well-oxygenated seafloor before the final burial. Bioerosion of bones caused by phototrophic cyanobacteria and algae, heterotrophic fungi and bacteria (Orciano balaenopterid WFi4-1 in S4-FB), and eventually by whalebone-eating siboglinid worm of genus *Osedax* (on a ziphiid humerus: Higgs et al. 2012), occurred at condensed intervals, in association with glauconite (Danise 2010). In one instance, a condensed shelly interval is traced for a few kilometers, connecting bioturbated shoreface sandstones yielding slightly disarticulated dugong skeletons (MOm2-1, MOm2-2) and other LMVs (Tinelli, 2013), with open shelf deposits yielding slightly disarticulated whale remains (WOm2-1 in S2-OOB). Regarding a sequence stratigraphic interpretation, pristine skeletons from delta front sediments can be part of the FSST (Fig. 4; or in alternative interpretation, of the early TST, when incised valleys are filled with coastal deposits). TST deposits account for 12% of cases. Much more frequently, slightly disarticulated skeletons are associated with the HST of the corresponding depositional sequence, lying above the MFS (69% of cases), or directly in coincidence with it, above a laterally-persistent shell bed (10%). Bone bioerosion is less pronounced in the late HST, when sedimentation rate increases (Castelfiorentino whale WEI3-2 in S3-EB).

67. Discussion**Factors of the Pliocene NWMS marine megafauna fossil record**

The detailed sequence-stratigraphic framework and the abundant shell beds, offering an independent check on sedimentary facies distribution by benthic paleoecology, allows also to explore factors behind MM geographic and stratigraphic distribution, and to sort out evolutionary, ecological, and taphonomic drivers of this particular fossil record. NWMS Pliocene distribution can be compared with similar studies in different settings and at different times, to draw conclusions on the general quality of MM fossil record.

7.1 Evolutionary control

Available data allow a meaningful comparison of NWMS MM diversity across the Pliocene, particularly detailed for marine mammals (Fig. 9D). Although the study suggests a facies control on MM distribution, with large vertebrate MM remains being generally connected associated with late TST-HST muddy sandstones and mudstones from lower shoreface and offshore shelf paleosettings, the temporal pattern of biodiversity recorded on a regional basis should in part likely represent reflects also a paleobiologic phenomenon, since lower shoreface and offshore shelf sediments are represented in all synthem. Marine mammals are unrecorded in the thickest part of synthem S1-2 (Zanclean, mostly bathyal mudstone and turbiditic sandstone), but they are present in S2, in OOB (Zanclean shoreface and open shelf deposits) and SRB (Zanclean upper slope deposits). MM is uncommon gradually rises in S3, at the upper Zanclean-lower Piacenzian, whereas it suddenly peaks in synthem S4, where species richness of both marine mammals and sharks is highest. S5 reflects a lower diversity of marine mammals, but still a high diversity of sharks, while values of both groups drop to the lowest abundance and species richness in S6, during the upper Piacenzian-lowermost Gelasian (Fig. 9D).

The S4 diversity peak coincides with the middle part (3.264–3.025 Ma) of the Piacenzian, a time interval in which the earth experienced global average temperature 1.84 °C-3.60 °C

warmer than the pre-industrial period (Dowsett et al., 2013). Climatic impact is testified by the widespread occurrence of carbonate deposits in S4 (Fig. 2), with sedimentary facies indicative of warm-temperate to subtropical conditions, with summer sea-surface temperature considerably warmer than 20°C and winter temperatures colder than 20°C (Nalin et al., 2016). This suggests a causative link between global climate and biodiversity, S4 diversity peak recording a global phenomenon, possibly an increase of speciation rate connected with global warming. Similarly, we propose that the lower diversities recorded at S5-S6 are the regional expression of an increase in extinction rate related to climatic cooling and global sea-level drop, ultimately leading to the global MM extinction event recorded on a coarser scale at the Pliocene-Pleistocene boundary (Pimienta et al., 2017): the finer stratigraphic resolution adopted here suggests a stepwise extinction event. NWMS data also point to a selective effect, extinction being recorded by marine mammals, with a 100% regional turnover between Pliocene and Recent (Tab. 3), but not as much by the shark fauna, with 34% of the species still living in the Mediterranean, while still others have shrank their distribution to subtropical latitudes. Also the Piacenzian teleost fish fauna (Cigala Fulgosi et al., 2009) and the benthic mollusks (Raffi et al., 1985) show a high percentage of holdovers, suggesting that marine mammals have been particularly prone to climatic change (see also Steeman et al., 2009).

Estimates of body size in Pliocene mysticetes of Tuscany, with several specimens reaching 10 m (Danise and Dominici, 2014: seven specimens in Tab. 1), are comparable to global values derived from the literature for this time interval (Lambert et al., 2010), confirming that NWMS baleen whales were larger than their Miocene analogues, and smaller than modern forms. As with regard to odontocetes, the most common Pliocene delphinid, *Etruridelphis*, was larger than the modern analogue *Stenella* (Bianucci et al., 2009b). The same is true for *Hemisyntrachelus cortesii*, larger than modern *Tursiops* (Bianucci, 1997a). On the other hand, the largest extant delphinid *Orcinus orca*, reaching 9 m, is about twice as long as

Orcinus citonensis (Heyning and Dahlheim, 1988). The high percent of holdover points to a more conservative figure for larger sharks (Tab. 5), but the presence in the Pliocene of the gigantic *Carcharocles megalodon* and some large thermophilic species today restricted to lower latitudes suggests that impoverishment of the fauna is coupled with an average decrease in size (Marsili, 2008). Comparing sizes suggests an overall restructuring of NWMS MM during the last three million years (see also Bisconti, 2009).

The geographic and stratigraphic distribution of large marine vertebrates in the fossil record, as that of any other key elements of marine ecosystems, depends on ecological (i.e., the production of organic parts) and taphonomic factors (their preservation). The detailed sequence-stratigraphic framework available for the very thick Pliocene succession of Tuscany, and the abundant shell beds that offer an independent check on sedimentary facies distribution by benthic paleoecology, allow to explore ecological and taphonomic drivers on the distribution of Northwestern Mediterranean LMVs. In its turn, this can be compared with similar studies in different settings and at different times, to draw conclusions on the general quality of LMV fossil record.

67.21 Ecological control

Marine

The composition of the Pliocene NWMS MM is affected in the first place by the availability of food. At the lowest trophic level, inferring from the ecology of the modern MM (Tabs. 3, 5), we found herbivore sirenians feeding on seagrasses (trophic level, TL = 2,0), all others being carnivores, thus having TL > 3,0 (Fig. 1211). The lowermost levels among the carnivores were occupied by baleen whales and whale sharks (TL = 3,2-3,4), filter-feeding on pelagic crustaceans and fishes (krill). Roughsharks, catsharks, and houndsharks, with many species living in the modern Mediterranean, have a relatively species-poor Pliocene record, probably due to a preservation bias related to their small size. At TL = 3,7-4,0 were

one species of catshark and a monk seal, both feeding near the seafloor on crustaceans, teleost fishes and cephalopods. At TL = 4,1 were sandbar, tiger and blue sharks, feeding on teleost fishes, cephalopods and on marine mammals. At the same level, in slope environments, kitefin sharks mainly fed on other sharks. The majority of MM species were found at TL = 4,2, with smaller odontocetes (three species) and 11 species of sharks, including several requiem sharks, a houndshark and a hammerhead. Larger dephinids, sperm whales, beaked whales, together with mackerel, sand and sand tiger sharks, occupied high trophic levels (TL = 4,3), followed at the top of the global NWMS food web by white shark, megalodon shark, one species of sevengill shark (genus *Notorynchus*), and killer whales, all feeding on marine mammals and smaller sharks (TL = 4,4-4,7). With no exception, all highest levels encountered in modern NWMS offshore pelagic and nearshore communities were occupied during the Pliocene by an analogous MM, often by the same species (large sharks), or by congeneric or con-familial species (marine mammals: Tab. 3, 5). The Pliocene pelagic ecosystem, typified by the mid-Piacenzian S4 association, must have been however trophically more diversified (Fig. 11), including aquatic megaherbivores, several balaenid filter feeders, larger and more diverse dolphins, and sharks species today extinct (e.g., *Carcharocles megalodon*) or restricted to tropical seas (e.g., *Galeocerdo cuvier*). With the exception of TL = 2, all Pliocene NWMS MM were either apex predators of their community, or mesopredators, occupying trophic positions below apex predators. The definitions of apex predators (or top predators) and mesopredators are relative and to an extent context-dependent (species that in a context are apex predators, in another are mesopredators, e.g., Estes et al., 1998). Since predation is a trophic interaction in which one animal (predator) consumes another (prey) as a source of energy (food), irrespective of the means by which this is accomplished (Lourenço et al., 2013), filter-feeding baleen whales can be considered apex-predators of their community (e.g., Lewiston et al., 2004; Notarbartolo di Sciara et al., 2016). Among sharks, 68% of living Mediterranean

elasmobranches are ranked as top predators, with a trophic level of 4 or more (Goffredo and Dubinsky, 2014, including superorder Batoidea, against 22% of teleost fishes), an estimate that can be extended to Pliocene NWMS MM. Relationships between apex predators and mesopredators are complex and hard to define in ecology, involving predation on other predators (intraguild predation, combining competition and predation: Polis et al., 1989), where consumption and competition need to be proved (Lourenço et al., 2013). In the Mediterranean Pliocene, paleontological evidences of carnivores serving as food to MM include killing of prey (Bianucci et al., 2010) and scavenging (Cigala Fulgosi, 1990; Bianucci et al., 2002; Dominici et al., 2009). The occurrence of intraguild predation must have been far more extended than what taphonomy can prove, however, given a Pliocene diversity of NWMS very large raptorial feeders higher than the modern, including the killer whale *Orcinus citonensis*, the large delphinid *Hemisyntrachelus cortesii* (phylogenetically related to the modern killer whale: Murakami et al., 2014), the white shark *Carcharodon carcharias*, the largest shark of all times *Carcharocles magalodon* (Marsili, 2008), and a diverse association of smaller carnivores, including monk seal, delphinids and sharks with $4,0 < TL < 4,2$ (Tab. 3, 5), candidate prey for larger raptorial feeders. This interaction likely exerted in its turn a control on community structure at lower trophic levels through processes like “mesopredator release” and trophic cascades (Roemer et al., 2009), eventually linking pelagic and nearshore communities, including benthic animals and plants, like in many modern ecosystems (Estes et al., 2011, 2016), down to slope depths (e.g., Parrish, 2009). Large raptorial feeders could exert a control on the diversity of the filter-feeding MM, like it has been suggested on a global scale since the Miocene (Lambert et al., 2010), and for the Pliocene by Bisconti (2003), when baleen whales were more diversified, both in terms of species richness, size range, and feeding strategies, including both skim- and lunge-filter feeding (respectively balaenid and balaenopterid whales: Berta et al., 2016; Hocking et al., 2017), minimising competition for food and diversifying spatial niches (see also Marx et al.,

2017, for the upper Miocene). A further important top-down control on community structure is suggested by modern studies on the role of baleen and sperm whales as nutrient vectors, both in horizontal (during seasonal migration) and vertical direction (during daily feeding migration: Roman and McCarthy, 2010), a process particularly important in oligotrophic seas (Alleger et al., 2017), like large sectors of the Mediterranean. Finally (literally, after death), MM becomes a detrital sources of energy and habitat for deep sea whale-fall communities (Roman et al., 2014; Smith et al., 2015), with paleontological evidence available for the NWMS (Dominici et al., 2009; Higgs et al., 2012; Baldanza et al., 2013).

Much evidence suggests that a bottom-up control on the structure of NWMS MM community was exerted by wind-driven upwelling currents, through enhanced oceanic productivity and concentration of preys. Over geologic time, these factors may have exerted a selection favoring large size, triggering the Plio-Pleistocene emergence of whale gigantism in several lineages (Slater et al., 2017). The largest among marine top predators can travel long distances and cross oceans, but tend to congregate in shallow waters with abundant prey. Baleen whales, abundant in boreal eutrophic waters (Woodley & Gaskin, 1996), with population size under the control of food availability (Croll et al., 2005), always require high prey density for efficient bulk filter feeding (Goldbogen et al., 2011). Six different species of mysticetes currently foraging in the Southern ocean, among which the largest animals that have ever lived in world oceans, exploit the high biomass of Antarctic krill, their main food resource. The Great white shark is abundant in the offshore of California (Jorgensen et al., 2010), Australia-New Zealand, South Africa, and formerly in the Mediterranean (Compagno, 2001; Bonfil et al., 2005), clustering in proximity of seal colonies. Marine mammals are in fact an important food resource in some areas, in addition to fish and cephalopods including monk seals. The distribution in the modern Mediterranean is no exception to general rule, only in a smaller scale, with large sharks and marine mammals congregating in productive areas, such as the Strait of Sicily and the Balearic Archipelago

(Tabs. 3, 5), with the second largest animal on earth, the fin whale, showing a movement pattern that parallels seasonal variability in available feeding habitat (Notarbartolo di Sciara, 2016), contributing to the horizontal transfer of nutrients. Large sharks can also prey or scavenge on cetaceans (Compagno, 2002), whereas foraging may also take place in the offshore, diving at 100-200 m on average (Jorgensen et al., 2010). Among the pinnipeds, grey seals live from estuaries to open shelves (Boehme et al. 2012) and elephant seals are abundant in upwelling zones (Biuw et al. 2007). Like whales, also pinnipeds are diverse and abundant in the nutrient-rich Southern Ocean, where they feed on krill, cephalopods, and fish. The diverse association of Pliocene pinnipeds, sharks and whales in Tuscany points therefore to the presence of nutrient-rich waters in the Northwestern sector of the Mediterranean (NWMed). A further and independent paleoecological evidence of a nutrient-rich regime is the association of many LMVs with “*Turritella* beds”, turritelline gastropods showing gregarious habit and high abundance in areas of high nutrient content (Allmon, 1988). The modern Tuscan shoreline faces Tuscany faces the Ligurian Sea, the northernmost reach of the Mediterranean, where a deep-water upwelling current coming from Southeast convects nutrients to the water surface, leading to high levels of primary productivity in its western sectors, extending westward to the Provençal and Balearic Seas, with spring algal blooms. These waters host different trophic regimes with respect to other parts of the Mediterranean, in an otherwise oligotrophic Mediterranean Sea (Lazzari et al., 2012; Melanotte-Rizzoli et al., 2014; Stambler, 2014). Supporting a conspicuous biomass of zooplankton (Cuzin-Roudy 2011), the modern NWMed Ligurian Sea sustains large populations of fin whales (*Balaenoptera physalis*) and striped dolphins (*Stenella coeruleoalba*), together with other large cetaceans (Notarbartolo di Sciara et al., 2008). Mediterranean-resident fin whales have adapted to exploit localised mesoscale hotspots of productivity that are highly variable in space and time (Notarbartolo di Sciara et al., 2016), feeding behaviors possibly mediated by the depth of prey and species-specific behaviours.

allowing to minimise competition with other large filter feeders (see Friedlaender et al., 2014). ~~Among the odontoceti,~~The sperm whale *Physeter macrocephalus* uses habitat across a range of depths and a specialised diet (Rendell and Frantzis, 2016), gathering along ~~NWMed~~NWMS steeper slopes, where water currents allows for higher trophic level biomass (Pirotta et al., 2011). ~~Beaked whales are particularly abundant in the Ligurian Sea and central Tyrrhenian Sea, preferring submarine canyons at slope depths (Podestà et al., 2016). The largest among common Mediterranean delphinids, *Grampus griseus*, is frequent in the Ligurian and Thyrrhenian seas where it forages on cephalopods at depths 300-1500, where upwelling currents are most effective (Azzellino et al., 2016).~~

The ~~NWMed~~NWMS existed as a Liguro-Provençal back-arc oceanic basin since the upper Miocene, when Sardinia rotated to its present position (Gattacceca et al., 2007), and in coincidence with the formation of the hinterland basins here under study (Muttoni et al., 2001), justifying the assumption that the modern ~~NWMed~~NWMS trophic regime is a feature that dates back at least to the upper Miocene. ~~Past eutrophic conditions are testified in Tuscany by d~~Consistently, diatomites deposited in Tuscany during the Messinian, before the salinity crisis (Bossio et al., 1997; Roveri et al., 2014), ~~indicate that high productivity was a primitive feature of the Ligurian Sea and by t.~~ Paleontological evidences include the very high diversity and abundance of Pliocene ~~LMVs~~MM in FB, the closest to modern upwelling areas (Fig. 3), ~~and the association of several MM with “*Turritella* beds”, turritelline gastropods showing gregarious habit and high abundance in areas of high primary productivity (Allmon, 1988).~~ Both a comparison with the modern and paleoecologic data strongly suggest that, ~~particularly at times of transgression, maximum flooding and highstand,~~ a wedge of ~~NWMed~~NWMS nutrient-rich waters intersecting the Tuscan shelf ~~would have positively affected the general biomass~~exerted a strong bottom-up control ~~on community structure through mixing and upwelling of nutrients, stimulation of phytoplankton blooms, followed by zooplankton increase, while nekton and vertebrates tracked plankton~~

concentrations, as is typical of modern upwelling systems throughout the world (Polis et al., 1997). High surface primary productivity in the study area during the Pliocene would have caused a concentration of ~~food~~detritus-falls, supporting a diverse community of deep-sea scavengers.

67.32 Taphonomic control

Taphonomic data on articulation and completeness of MM specimens are available for large mammals, the shark record being formed mainly by isolated teeth collected through superficial picking during the years (Cigala Fulgosi et al., 2009), with only a few contextualised studies (e.g., Bianucci et al., 2002; Dominici et al., 2009). Our record includes several marine mammals with a high degree of articulation (14%) and completeness (27%), or with at least cranial and post-cranial elements of the same individual (38%). All articulated specimens and the most complete skeletons are associated with shelf mudstones, usually forming the HST of the relative depositional sequence (Tab. 2), with two exceptions, one associated with delta-front, coarse-grained beds (WCh6-1, Fig. 6), another from outer shelf or upper slope sediments (DSi2-1). These data point to a strong environmental control on the quality of the MM fossil record, at least concerning articulation and completeness of skeletons, very shallow and very deep (bathyal) depths being generally unfavourable to the preservation of MM bones (Fig. 9A). To explain these results, hypotheses are based on the available data on carcasses of MM in modern marine environments.

The biostratinomy of lung-breathing marine mammals depends on water depth (Allison et al. 1991; Smith et al., 2005). The vast majority of mammals are negatively buoyant and sink after death, but in shallow waters the low hydrostatic pressure allows putrefaction gases to develop and carcasses to resurface. Skeletons are scavenged (Dicken, 2008) and disintegrate while floating, leading to the preferential deposition of isolated bones. At deeper

settings, where high hydrostatic pressure allows the carcass to lay relatively undisturbed on the seafloor, the skeleton becomes only slightly disarticulated after soft tissue removal (Reisdorf et al., 2012). The subsequent fate of deep-water, disarticulated skeletons depends on the nature of the scavenging fauna and other elements of the whale-fall community, and on the time of exposure on the seafloor (Boessenecker et al. 2014). At depths deeper than the slope breaks, where sedimentation rate is very low, carcasses are exposed for a long time and the skeleton is rapidly disintegrated. Time-series analyses carried out at modern whale-fall communities at slope depths (range 382-2893 m: Lundsden et al., 2010) suggests that carcasses up to 17 m are rapidly degraded, with the deepest whale carcasses disappearing after only seven years of exposure on the sea floor. Larger skeletons may persist on deeper settings for decades, but if not buried, they also ultimately undergo complete destruction. Although environmental forcing triggered by higher temperatures, active currents and sediment transport, clearly plays a role, a specialized whale-fall fauna rapidly consumes both soft and mineralized tissues. This fauna is characterized by low diversity and high abundance of microorganisms, most likely as a result of both specialization to nutrient enrichment and high growth rates. Whale-fall habitats likely undergo a temporal microbial succession from primarily heterotrophic to more heterotrophic/chemosynthetic metabolisms until the whale biomass is completely exploited (Smith et al., 2015). Of all the specialized taxa, bone-eating polychaetes of genus *Osedax*, with their soft root-like tissues that erode the bones to access nutrients (Tresguerres et al., 2013; Minamoto et al., 2017), are the primary cause of bone disintegration, particularly of denser bones (Higgs et al., 2011). Found also at shelf depths, but invariably in low abundance (Huusgaard et al., 2012; Higgs et al., 2014b), bone-eating worms occur in high numbers in the deep sea (Smith et al., 2015) where they act as biodiversity regulators (Alfaro-Lucas et al., 2017). The general paucity of novel taxa on shallow-water whale falls suggests that species-rich, specialized whale-fall communities

develop only in the food-poor deep sea. Accordingly, among new animal species described in the recent literature from whale falls, only about 10% have been found on whale remains at depths of less than 260 m (12 out of 129 new species: Smith et al., 2015). Of all ~~large marine vertebrates~~MM investigated so far, only whales are known to host a whale-fall community, but also carcasses of large elasmobranchs undergo rapid destruction ~~on the deep-sea~~at bathyal depths, teeth being all that eventually remains (Higgs et al., 2014a; teeth are also lost by sharks during feeding: Pokines and Symes, 2013). As a consequence, over geological time no large marine vertebrate is expected to be recovered at depths greater than the shelf break. On the opposite side, the lack of a biota specialized ~~in~~sed in exploiting large organic falls, coupled with higher rates of deposition in proximity of sediment sources, make it more probable that ~~articulated, or only partially disarticulated~~the most articulated and complete whale skeletons become part of the fossil record of shelf settings below storm wave base.

78. Comparison with other studies

The study of sedimentary facies uncovers some environmental factors that directly control the taphonomy of large marine vertebrates. Such abiotic drivers include water pressure, wave energy and sedimentation rate — three factors summarized ~~by~~sed by water depth, and upwelling, bringing to the surface deep sea nutrients and concentrating preys. The taphonomic pathway of large marine carcasses is also driven by biotic factors that change in geological time in response to coevolution between bacteria, scavengers and their substrates. We now explore the multifaceted nature of ~~LMVMM~~ taphonomy by reviewing Mesozoic and Cenozoic studies ~~comparable to ours~~where sufficient data for stratigraphic paleobiology are available.

7.8.1 Abiotic factors

According to one of the few previous studies on the sequence stratigraphic distribution of ~~large marine vertebrates~~MM, Jurassic ichthyosaurs, plesiosaurs, and pliosaurs of the Sundance Seaway, in North America, display facies control and are found primarily in offshore mudstone and at condensed intervals at the maximum flooding surface, rather than shoreface and estuarine sandstone (McMullen et al.,¹⁷ 2014, ~~in a study that included isolated LMV remains~~). Taphonomic data on Upper Cretaceous marine reptiles and large fishes suggest that partially articulated and disarticulated skeletons are associated with little biological activity and relatively rapid burial by muddy sediments, deposited in the North American seaway during an interval of maximum flooding (Schemisser McKean ~~&and~~ Gillette,¹⁷ 2015). Upper Cretaceous mosasaur remains are particularly concentrated in fine-grained shelf deposits in Europe (Jagt ~~&and~~ Jagt-Yazykova,¹ 2016). Complete, partially articulated whale skeletons of archaeocetes, together with bones and teeth of ~~dugongsirenians~~ and sharks, are abundant in offshore marine flooding surfaces (MFS) in the Priabonian late TST of Egypt. In a further parallelism with the Pliocene of Tuscany, Late Eocene well-articulated whales are associated with rapidly accumulating shoreface sediments of the FSST, comprising “*Turritella* shell beds” (Peters et al.,¹⁷ 2009). During the Oligocene, eomysticete whale bones were deposited at shelf depth below storm wave base. The occurrence of sparse traces attributed to *Osedax* and the association with a glauconitic limestone testify to the exposure of bones on the seafloor without undergoing complete destruction (Boessenecker ~~&and~~ Fordyce,¹⁷ 2014), in a manner similar to ~~LMVs~~MM at Orciano (see Danise et al.,¹⁷ 2010;¹⁷ Higgs et al.,¹⁷ 2012). Burdigalian/Langhian MM recovered in the Antwerpen Sands, in Belgium (including isolated remains of a baleen whale, several odontocetes and a pinniped) are fragmented, worn and associated with clayey sandstone

rich in glauconite, suggesting long exposure on the sea-floor (Louwye et al., 2010). Bones are concentrated at the base of a coarsening-upward succession, on top of shallow marine, coarse-grained sandstone, suggesting this is a surface of maximum flooding. An association of LMVsMM taxonomically comparable to ~~those that~~ here under study is encountered in the Mio-Pliocene of the Purisima Formation, in Central California. Taphofacies differ in some aspects. The Pliocene of California yields laterally persistent bonebeds with polished and phosphatised bones, and abundant phosphate nodules that are absent in Tuscany, indicating times of higher sediment starvation during transgressive pulses, in an area of much stronger nutrient content (the California Current system is a northern-hemisphere analogue of the Peruvian upwelling system, associated with the economically most important fish stocks in the world: Mann &and Lazier₁₇, 2006). Shoreface deposits indicate stronger wave energy, and the preferential absence of molluscs in bonebeds indicates chemical destruction of carbonate shells (Boessenecker et al.₁₇, 2014, in a quantitative MM taphonomic study ~~that includes isolated LMV fragments~~). Episodic sedimentation, however, causes the preferential preservation of articulated remains in the Californian offshore as in the Pliocene of NWMedNWMS. The late Miocene Pisco Formation in Perú offers another, more extreme example of LMVMM taphonomy in a eutrophic setting. Here almost two hundred cetaceans, pinnipeds, and sharks were described in an exceptional state of preservation (Bianucci et al.₁, 2015), within a monotonous succession of finely laminated white diatomites (Di Celma et al.₁, 2015), suggesting very high primary productivity in an area of intense upwelling and volcanic activity. Algal blooms sustained high biomass of apex predators (see Marx &and Uhen₁, 2010), triggering at the same time anoxic conditions at the seafloor where LMVMM carcasses remained intact (Brand et al.₁, 2004₁₇; see also Gioncada et al.₁, 2016; Marx et al., 2017, also in offshore paleosettings). Finally, the stratigraphic distribution of Pliocene cetaceans in western Emilia (Italy) shows an uneven distribution of findings (N = 24, dolphins and baleen whales) and a strong positive correlation with offshore

mudstones (no findings in shoreface sandstones, rare occurrences in epibathyal mudstones: Freschi and Cau, 2016), paralleling the distribution of Tuscan Pliocene MM.

78.2 Biotic factors

Many reviews of Triassic (Camp₁ 1980₁₇; Hogler₁ 1992; Motani et al.₁ 2008; Hu et al.₁ 2011; Liu et al.₁ 2014) and Early Jurassic marine reptiles (Benton &and Taylor₁ 1984) report mostly good preservation and a high degree of completeness and articulation of skeletal material. This record might be partially controlled by prevailing anoxic or dysoxic conditions in the bottom waters of many Mesozoic fossiliferous deposits (e.g., Middle Triassic Besano and Guangling Formations, Lower Jurassic Blue Lias and Posidonia Shale Formations), which precluded organism activity within the sediment, and prevented predation or scavenging of the carcasses on the sea bottom (Beardmore &and Furrer₁ 2016). Evidence of advanced levels of disarticulation or bone degradation (Martill₁ 1985; sauropterygians, crocodilians, ichthyosaurs and fishes from the Middle Jurassic Oxford Clay Formation), is mostly attributed to physical factors (e.g., weathering on the sea floor). Up to the early-Late Cretaceous, biological activity is testified by circumstantial evidence of scavenging (*Hybodus* teeth associated with marine reptile skeletons, Martill et al.₁ 1994), and by the more common occurrence of microbial mats, grazers and encrusters (Martill₁ 1987₁₇; Meyer₁ 2011₁₇; Danise et al.₁ 2014₁₇; Reolid et al.₁ 2015), but lack traces of bone-eating worms and sulphophilic fauna typical of modern whale falls. The siboglinid *Osedax* is an evolutionary novelty in possessing a root system that hosts heterotrophic mutualists and secretes bone-dissolving acids (Tresguerres et al.₁ 2013₁₇; Miyamoto et al.₁ 2017), and an ecosystem engineer (Alfaro-Lucas et al.₁ 2017). Genus *Osedax* is today associated with whale falls worldwide (Taboada et al.₁ 2015), but its impact on LMVMM has changed in time. The oldest trace fossils attributable to a bone-eating fauna is found on early-Late Cretaceous plesiosaur

and sea turtle bones (about 100 Ma: Danise ~~&~~ Higgs, 2015). Time estimates suggest that *Osedax* diverged from other siboglinids in the Middle Cretaceous (around 108 Ma: Taboada et al., 2015). However, if the bone-eating worm lives also on the bones of birds and terrestrial mammals (Rouse et al., 2011), its global nature and high species diversity in modern oceans suggest that whale falls, as complex and species-rich habitats, have been the most important biodiversity generators (Higgs et al., 2014b; Smith et al., 2015). And although the first appearance of *Osedax* is concomitant with the occurrence of large marine reptiles and teleost fishes in the oceans, long before whale evolution, and although their distribution is not limited to large carcasses (Pyenson and Haas, 2007), the radiation of ocean-going mysticetes at the Oligocene onset of the Antarctic Circumpolar Current (Fordyce, 2003) clearly increased available substrata worldwide. The steady increase of cetacean size during the Neogene, with a dramatic pulse in the last five million years, when Neoceti surpassed 10 m length and reached 30 m in the Pleistocene (Lambert et al., 2010; Slater et al., 2017), would have thus triggered a second and more massive radiation of bone-eating worms (Kiel and Goedert, 2006). Consistently with this hypothesis, the preferential distribution of modern *Osedax* in high latitude settings worldwide (Taboada et al., 2015) suggests that biodiversity hotspots coincide with the feeding grounds of larger cetaceans. Other bone-eaters of modern deep water whale-fall ecosystem belong to the group of abyssochrysid snails, with fossils found on Late Cretaceous plesiosaur ~~bones~~ (Kaim et al. 2008) and sea turtle bones (within a chemosynthesis-based association: Jenkins et al., 2017). Modern abyssochrysid whalebone-eaters of genus *Rubyspira*, hosting a specific and exclusive microbiome (Aronson et al., 2017), split during the upper Eocene/lower Oligocene (Johnson et al., 2010). ~~Like for *Osedax*, s~~Species of *Rubyspira* ~~likely~~ benefited too from the radiation of ocean-going whales. Although scanty, available evidence on the geological history of bone-eaters thus makes the ephemeral nature of large carcasses in

modern deep seas — and their absence in bathyal deposits of the Pliocene of ~~NWMed~~NWMS — a larger-than-life model for the Mesozoic and the early Paleogene.

9. Conclusions

- 1) Sedimentary facies in the Pliocene of Tuscany are vertically stacked to form small-scale depositional sequences particularly in the upper half, Piacenzian part of the succession, with laterally-continuous shell beds marking transgressive surfaces and ~~the surface~~intervals of maximum flooding. Small-scale sedimentary sequences are stacked to form six major, unconformity-bounded stratigraphic units (synthems) of regional extension, forming a high-resolution framework to study the chronostratigraphic distribution of marine megafauna (MM).
- 2) ~~Sedimentary facies~~Benthic biotopes, identified through a quantitative study of a large mollusk~~mollusc dataset~~, can be arranged to form an ideal onshore-offshore, bathymetric gradient, connecting terrestrial environments with deep sea epibathyal bottoms, consistently with the distribution of sedimentary facies. Bones~~MM remains of large marine vertebrates~~ and shell beds are present in all marine paleoenvironments, but their distribution is uneven.
- 2) The fossil record of large vertebrates, including marine mammals (abundance data) and ~~larger~~ sharks (presence-absence data), is particularly rich in sediments deposited between storm wave base and ~~the~~ shelf break, moderately rich in shoreface sediments, very poor in epibathyal sediments. Paleobathymetric estimates suggest that a peak of abundance of better preserved skeletons occurs around 100 m depth, within the 30-300 m range.
- 3) Species-rich~~ness~~ of MM and abundance of marine mammal remains steadily increase during the Zanclean, reaching a peak in synthem 4, possibly as a response to the mid-

Piacenzian warm period, and gradually decrease until reaching a minimum in proximity of the Plio-Pleistocene boundary.

4) Comparison with ecological data on a global scale, paleogeographic and topographic considerations on the North-Western Mediterranean Sea (NWMS) Mediterranean paleogeography₁ and the frequentoccasional association of Pliocene large marine vertebrates^{MM} with *Turritella* beds₁ suggest that upwelling and high-nutrient conditions in the Northwestern Mediterranean offshore sustained during the Pliocene a rich community of apex predators and mesopredators. Higher taxonomic MM diversity in the NWMS during the Pliocene, suggests higher niche-partitioning, with respect to the modern NWMS and a top-down control on community structure.

5) A comparison with studies on the biota exploiting tissues of large food particles sunken on the seafloor, both modern and ancient, suggests that the poor epibathyal record of Pliocene larger vertebrates of Tuscany may be caused by the destructive action of bone-eating invertebrates. This biotic driver of the marine vertebrate fossil record was less efficient before the radiation of ocean-going whales in the late Eocene-lower Oligocene. On the other hand, bone-eaters played a major taphonomic role on a global scale after the Pliocene-Pleistocene increase in whale size.

6) Due to resurfacing of carcasses in shallow waters, the taphonomy of large marine vertebrates^{MM} fossil record through the Mesozoic-Cenozoic is reviewed through stratigraphic paleobiology shows a strong facies control, being more continuous comparable through the Mesozoic-Cenozoic in rocks formeddeposited below storm wave base and above the shelf-slope break during sea-level transgressions and highstands.

Acknowledgements

We thank Michelangelo Bisconti (SCNHM) for helpful discussion on mysticete taxonomy and paleoecology, Giovanni Bianucci (DSTUP) for his insightful review on the first draft of

3121
3122 this paper, and an anonymous reviewer. Maurizio Forli helped by sharing his thorough
3123
3124 knowledge of fossiliferous Pliocene outcrops of Tuscany.
3125
3126
3127

3128 **References**

3130
3131
3132
3133 Alfaro-Lucas, J.M., Shimabukuro, M., Ferreira, G.D., Kitazato, H., Fujiwara, Y. &and
3134 Sumida, P.Y.G. 2017. Bone-eating Osedax worms (Annelida: Siboglinidae) regulate
3135 biodiversity of deep-sea whale-fall communities. Deep Sea Research II
3136 <http://dx.doi.org/10.1016/j.dsr2.2017.04.011>.
3137
3138
3139

3140
3141
3142
3143 Barría, C., Coll, M., Navarro, J., 2015. Unravelling the ecological role and trophic
3144 relationships of uncommon and threatened elasmobranchs in the western Mediterranean
3145 Sea. Mar. Ecol. Prog. Ser. 539, 225-240, doi: 0.3354/meps11494.
3146
3147
3148
3149

3150
3151
3152 Barría, C., Navarro, J., Coll, M., 2017. Trophic habits of an abundant shark in the
3153 northwestern Mediterranean Sea using an isotopic non-lethal approach, Estuarine, Coastal
3154 and Shelf Science, doi: 10.1016/j.ecss.2017.08.021.
3155
3156
3157
3158

3159
3160
3161 Alderighi, L., Bianucci, G., Tinelli, C., Petri, A., 2011. Certaldo (FI). Località Canonica:
3162 scoperta e recupero di delfino fossile. Notiziario della Soprintendenza per i Beni Archeologici
3163 della Toscana 7, 250-252.
3164
3165
3166
3167

3168
3169
3170 Almond, W. 1988. Armfield, B.A., Zheng, Z., Bajpai, S., Vinyard, C.J., Thewissen, J.G.M.,
3171 2013. Development and evolution of the unique cetacean dentition. PeerJ 1:e24; DOI
3172 10.7717/peerj.24.
3173
3174
3175
3176
3177
3178
3179
3180

Arbeits, B., Cioppi, E., Dominici, S., Wierer, U., 2015. Certaldo (FI). Località San Giorsole: recupero di resti paleontologici. Notiziario della Soprintendenza per i Beni Archeologici della Toscana 11, 154.

Aronson, H.S., Zellmer, A.J. & Goffredi, S.K., 2017. The specific and exclusive microbiome of the deep-sea bone-eating snail, *Rubyspira osteovora*. FEMS Microbiology Ecology, 93, fiw250, doi: 10.1093/femsec/fiw250.

Azzellino, A., Aioldi, S., Gaspari, S., Lanfredi, C., Moulins, A., Podestà, M., Rosso, M., Tepsich, P., 2016. Risso's dolphin, *Grampus griseus*, in the Western Ligurian Sea: trends in population size and habitat use. In: Notarbartolo di Sciara, G., Podestà, M., Curry, B.E. (Eds.), Mediterranean Marine Mammals Ecology and Conservation. Advances in Marine Biology 75, 207-234, doi: 10.1016/bs.amb.2016.08.003.

Baldanza, A., Bizzarri, R., Famiani, F., Monaco, P., Pellegrino, R., Sassi, P., 2013. Enigmatic, biogenically induced structures in Pleistocene marine deposits: a first record of fossil ambergris. Geology 41, 1075-1078, doi: 10.1130/G34731.1.

Beardmore, S. R., Furrer, H., 2016. Evidence of a preservational gradient in the skeletal taphonomy of Ichthyopterygia (Reptilia) from Europe. Palaeogeography, Palaeoclimatology, Palaeoecology 443, 131-144.

Benson, R.B.J., Butler, R.J., Lindgren, J., Smith, A.S., 2010. Mesozoic marine tetrapod diversity: mass extinctions and temporal heterogeneity in geological megabiases affecting vertebrates. Proceedings of the Royal Society B: Biological Sciences, 277 (1683), 829-834.

Benvenuti, M., Dominici, S., 1992. Facies analysis, paleoecology and sequence stratigraphy in a Pliocene siliciclastic succession, San Miniato (Pisa, Italy). *Bollettino della Società Paleontologica Italiana* 31, 241-259.

Benvenuti, M., Bertini, A., Conti, C., Dominici, S., 2007. Integrated analyses of litho- and biofacies in a Pliocene cyclothemic, alluvial to shallow marine succession (Tuscany, Italy). *Geobios* 40, 143-158.

Benvenuti, M., Bertini, A., Conti, C., Dominici, S., Falcone, D., 1995a. Analisi stratigrafica paleoambientale integrata del Pliocene dei dintorni di San Miniato. *Quaderni del Museo di Storia Naturale di Livorno, Supplemento*, 29-49.

~~Benvenuti, M., Dominici, S., Rook, L., 1995b. Inquadramento stratigrafico-deposizionale delle faune a mammiferi villafranchiane (unità faunistiche Triversa e Montopoli) del Valdarno Inferiore nella zona a Sud dell'Arno. *Il Quaternario* 8, 457-464.~~

Benvenuti, M., Degli Innocenti, D., 2001. The Pliocene deposits in the Central-Eastern Valdelsa Basin (Florence, Italy), revised through facies analysis and unconformity-bounded stratigraphic units. *Rivista Italiana di Paleontologia e Stratigrafia* 107, 265-286.

~~Benvenuti, M., Bertini, A., Conti, C., Dominici, S., 2007. Integrated analyses of litho- and biofacies in a Pliocene cyclothemic, alluvial to shallow marine succession (Tuscany, Italy). *Geobios* 40, 143-158.~~

Benvenuti, M., Del Conte, S., 2013. Facies and sequence stratigraphic modeling of a Upper Pliocene–Lower Pleistocene fluvial succession (Valdelsa Basin, central Italy). *Sedimentary Geology* 294, 303-314.

Benvenuti, M., Del Conte, S., Scarselli, N., Dominici, S., 2014. Hinterland basin development and infilling through tectonic and eustatic processes: latest Messinian-Gelasian Valdelsa Basin, Northern Apennines, Italy. *Basin Research* 26, 387-402.

Benvenuti, M., Dominici, S., 1992. Facies analysis, paleoecology and sequence stratigraphy in a Pliocene siliciclastic succession, San Miniato (Pisa, Italy). *Bollettino della Società Paleontologica Italiana* 31, 241-259.

Benvenuti, M., Dominici, S., Rook, L., 1995b. Inquadramento stratigrafico- deposizionale delle faune a mammiferi villafranchiane (unita` faunistiche Triversa e Montopoli) del Valdarno Inferiore nella zona a Sud dell'Arno. *Il Quaternario* 8, 457-464.

Benvenuti, M., Moratti, G., Sani, F., Bonini, M., Oriol, L.O., Papini, M., Rook, L., Cavallina, C., Cavini, L., 2015. Messinian-earliest Zanclean tectonic-depositional dynamics of the Cinigiano-Baccinello and Velona basins (Tuscany, Italy). *Italian Journal of Geosciences*, 134, 237-254.

Berta, A., 2012. *Return to the sea: the life and evolutionary times of marine mammals*. University of California Press, Berkeley, California, 205 pp.

Berta, A., Kienle, S., Bianucci, G., Sorbi, S., 2015. A Reevaluation of *Pliophoca etrusca* (Pinnipedia, Phocidae) from the Pliocene of Italy: phylogenetic and biogeographic implications. Journal of Vertebrate Paleontology, DOI: 10.1080/02724634.2014.889144 e889144, doi: 10.1080/02724634.2014.889144.

Berta, A., Lanzetti, A., Ekdale, E.G., and Deméré. T.A., 2016. From teeth to baleen and raptorial to bulk filter feeding in Mysticete cetaceans: The role of paleontological, genetic, and geochemical data in feeding evolution and ecology. Integrative and Comparative Biology 56, 1271-1284, doi: 10.1093/icb/icw128.

Bianchi, V., Ghinassi, M., Aldinucci, M., Boaga, J., Brogi, A., Deiana, R., 2015. Tectonically driven deposition and landscape evolution within upland incised valleys: Ambra Valley fill, Pliocene–Pleistocene, Tuscany, Italy. Sedimentology— 62, 897-927, doi: 10.1080/17445647.2013.829412.

Bianucci, G., 1995. A new record of baleen whale from the Pliocene of Tuscany (Italy). Atti della Società Toscana di Scienze Naturali 102, 101-104.

Bianucci, G., 1996. The Odontoceti (Mammalia, Cetacea) from Italian Pliocene. Systematics and phylogenesis of Delphinidae. Palaeontographia Italica, 83: 73-167.

Bianucci, G., 1997. Bianucci, G., 1997a. Hemisyntrachelus cortesii (Cetacea, Delphinidae) from Pliocene sediments of Campore Quarry (Salsomaggiore Terme, Italy). Bollettino della Società Paleontologica Italiana 36, 75-83.

Bianucci, G., 1997b. The Odontoceti (Mammalia Cetacea) from Italian Pliocene. The Ziphiidae. *Palaeontographia Italica* 84, 163–192.

Bianucci, G., 1999. I cetacei fossili del Museo di Storia Naturale dell'Università di Pisa. *Atti della Società Toscana di Scienze naturali, Memorie, Serie A* 103, 63-68.

Bianucci, G., Bisconti, M., Landini, W., Storai, T., Zuffa, M., Giuliani, S., Mojetta, A., 2002. Trophic interactions between white sharks (*Carcharodon carcharias*) and cetaceans: a comparison between Pliocene and recent data. 33-48. In Vacchi, M., La Mesa, G., Serena, F., Sèret, B., (eds). *Proceedings 4ht Meeting of the European Elasmobranch Association, Livorno (Italy) 27–30 September 2000*. Imprimerie F. Paillart, Abbeville, France, 209 pp.

Bianucci, G., Di Celma, C., Landini, W., Post, K., Tinelli, C., de Muizon, C., Gariboldi, K., Malinverno, E., Cantalamessa, G., Gioncada, A., Collareta, A., Gismondi, R.-S., Varas-Malca, R., Urbina, M., Lambert, O., 2016. Distribution of fossil marine vertebrates in Cerro Colorado, the type locality of the giant raptorial sperm whale *Livyatan melvillei* (Miocene, Pisco Formation, Peru). *Journal of Maps*, 12, 543-557, doi: 10.1080/17445647.2015.1048315.

Bianucci, G., Landini, W., 2005. I paleositi a vertebrati fossili della Provincia di Pisa. *Atti della Società Toscana di Scienze Naturali, Memorie serie A*, 110, 1-22.

Bianucci, G., Mazza, P., Merola, D., Sarti, G., Cascella, A., 2001. The early Pliocene mammal assemblage of Val di Pugna (Tuscany, Italy) in the light of calcareous nannoplankton biostratigraphical data and paleoecological observations. *Rivista Italiana di Paleontologia e Stratigrafia* 107, 425-438.

Bianucci, G., Bisconti, M., Landini, W., Storai, T., Zuffa, M., Giuliani, S., Mojetta, A., 2002. Trophic interactions between white sharks (*Carcharodon carcharias*) and cetaceans: a comparison between Pliocene and recent data. 33-48. In Vacchi, M., La Mesa, G., Serena, F., Sèret, B., (eds). Proceedings 4ht Meeting of the European Elasmobranch Association, Livorno (Italy) 27–30 September 2000. Imprimerie F. Paillart, Abbeville, France, 209 pp.

Bianucci, G., Sarti, G., Catanzariti, R., Santini, U., 1998. Middle Pliocene cetaceans from Monte Voltraio (Tuscany, Italy). Biostratigraphical, paleoecological and paleoclimatic observations. *Rivista Italiana di Paleontologia e Stratigrafia* 104, 123-130.

Bianucci, G., Mazza, P., Merola, D., Sarti, G., Cascella, A., 2001. The early Pliocene mammal assemblage of Val di Pugna (Tuscany, Italy) in the light of calcareous nannoplankton biostratigraphical data and paleoecological observations. *Rivista Italiana di Paleontologia e Stratigrafia* 107, 425-438.

Bianucci, G., Bisconti, M., Landini, W., Storai, T., Zuffa, M., Giuliani, S., Mojetta, A., 2002. Trophic interactions between white sharks (*Carcharodon carcharias*) and cetaceans: a comparison between Pliocene and recent data. 33-48. In Vacchi, M., La Mesa, G., Serena, F., Sèret, B., (eds). Proceedings 4ht Meeting of the European Elasmobranch Association, Livorno (Italy) 27–30 September 2000. Imprimerie F. Paillart, Abbeville, France, 209 pp.

Bianucci, G., Sorbini, C., 2014. Le collezioni a cetacei fossili del Museo di Storia Naturale dell'Università di Pisa. *Museologia Scientifica Memorie* 13, 93-102.

Bianucci, G., Sorbi, S., Vaiani, S.C., Landini, W., 2009a. Pliocene marine mammals from Italy: A systematic and stratigraphic overview. In: International conference on vertebrate

palaeobiogeography and continental bridges across Tethys, Mesogea, and Mediterranean Sea, Bologna, Abstract book, 9-12.

Bianucci, G., Sorce, B., Storai, T., Landini, W., 2010. Killing in the Pliocene: shark attack on a dolphin from Italy. Palaeontology 53, 457-470, doi: 10.1111/j.1475-4983.2010.00945.x.

Bianucci, G., Vaiani, S.C., Casati, S., 2009**b**. A new delphinid record (Odontoceti, Cetacea) from the Early Pliocene of Tuscany (Central Italy): systematics and biostratigraphic considerations. Neues Jahrbuch für Geologie und Paläontologie Abhandlungen 254, 275-292.

Bianucci, G., Sorce, B., Storai, T., Landini, W., 2010. Killing in the Pliocene: shark attack on a dolphin from Italy. Palaeontology 53, 457-470, doi: 10.1111/j.1475-4983.2010.00945.x.

Bianucci, G., Sorbini, C., 2014. Le collezioni a cetacei fossili del Museo di Storia Naturale dell'Università di Pisa. Museologia Scientifica Memorie 13, 93-102.

Bianucci, G., Source, B., Storai, T., Landini, W., 2010. Killing in the Pliocene: shark attack on a dolphin from Italy. Palaeontology 53, 457-470.

Bianucci, G., Di Celma, C., Landini, W., Post, K., Tinelli, C., de Muizon, C., Gariboldi, K., Malinverno, E., Cantalamessa, G., Gioncada, A., Collareta, A., Gismondi, R.-S., Varas-Malca, R., Urbina, M., Lambert, O., 2016**5**. Distribution of fossil marine vertebrates in Cerro Colorado, the type locality of the giant raptorial sperm whale *Livyatan melvillei* (Miocene, Pisco Formation, Peru). Journal of Maps, 12, 543-557, doi: 10.1080/17445647.2015.1048315.

Bianucci, G., Vomero, V., 2014. The cetacean collections of Italian museums. *Museologia Scientifica*, Memorie 13, Volume 2, 135 pp.

Bisconti, M., 2002. An early Late Pliocene right whale (genus *Eubalaena*) from Tuscany (Central Italy). *Bollettino della Società Paleontologica Italiana* 41, 83-91.

Bisconti, M., 2003. Evolutionary history of Balaenidae. *Cranium* 20, 9-50.

Bisconti, M., Lambert, O., Bosselaers, M., 2017. Revision of “*Balaena*” *belgica* reveals a new right whale species, the possible ancestry of the northern right whale, *Eubalaena glacialis*, and the ages of divergence for the living right whale species. *PeerJ* 5, e3464, doi: [10.7717/peerj.3464](https://doi.org/10.7717/peerj.3464).

~~Biuw, M., Boehme, L., Guinet, C., Hindell, M., Costa, D., et al., 2007. Variations in behavior and condition of a Southern Ocean top predator in relation to in situ oceanographic conditions. PNAS 21, 13705-13710.~~

~~Boehme, L., Thompson, D., Fedak, M., Bowen, D., Hammill, M.O., et al., 2012. How many seals were there? The global shelf loss during the Last Glacial Maximum and its effect on the size and distribution of grey seal populations. PLoS ONE 7(12): e53000. doi:10.1371/journal.pone.0053000~~

Boessenecker, R.W., Perry, F.A., Schmitt, J.G., 2014. Comparative taphonomy, taphofacies, and bonebeds of the Mio-Pliocene Purisima Formation, Central California: strong physical control on marine vertebrate preservation in shallow marine settings. PLoS ONE 9(3): e91419. doi:10.1371/journal.pone.0091419.

Boessenecker, R.W., Fordyce, R.E., 2014. Trace fossil evidence of predation upon bone-eating worms on a baleen whale skeleton from the Oligocene of New Zealand. Lethaia 48, 326-331.

[Boldrocchi, G., Kiszka, J., Purkis, S., Storai, T., Zinzula, L., Burkholder, D. 2017. Distribution, ecology, and status of the white shark, *Carcharodon carcharias*, in the Mediterranean Sea. Reviews in Fish Biology and Fisheries, doi: 10.1007/s11160-017-9470-5.](#)

Commented [1]:

Bonfil, R., Meyer, M., Scholl, M.C., Johnson, R., O'Brien S., et al., 2005. Transoceanic migration, spatial dynamics, and population linkages of white sharks. Science 310, 100-103.

Borselli, V., Cozzini, F., 1992. Il recupero di un cetaceo fossile in località Ponte a Elsa (Pisa). Museologia Scientifica 8, 9-22.

Bossio, A., Mazzei, R., Salvatorini, G., Sandrelli, F., 1993. Geologia dell'area compresa tra Siena e Poggibonsi ("Bacino del Casino"). Atti della Società Toscana di Scienze Naturali 107, 69-85.

Bossio, A., Mazzanti, R., Mazzei, R., Pascucci, V., Salvatorini, G.F., Sandrelli, F., 1994. Il Bacino di Volterra: notizie preliminari sull'evoluzione della sua area centro meridionale durante il Pliocene. Studi Geologici Camerti, volume speciale 1, 19-31.

Bossio, A., Foresi, L.M., Mazzanti, R., Mazzei, R., Salvatorini, G.F., 1997. Note micropaleontologiche sulla successione miocenica del Torrente Morra e su quella Pliocenica del bacino dei fiumi Tora e Fine (Provincia di Livorno e Pisa). Atti della Società Toscana di Scienze Naturali 104, 85-134.

Brogi, A., 2011. Bowl-shaped basin related to low-angle detachment during continental extension: the case of the controversial Neogene Siena Basin (central Italy, Northern Apennines). Tectonophysics, 499, 54-76.

Camp, C.L., 1980. Large ichthyosaurs from the Upper Triassic of Nevada. Palaeontographica Abteilung A, A170 (4-6), 139- 200.

Canese, S., Cardinali, A., Fortuna, C.M., Giusti, M., Lauriano, G., Salvati, E., Greco, S., 20067. The First identified winter feeding ground of fin whale (*Balaenoptera physalus*) in the Mediterranean Sea. Journal of the Marine Biological Association of the U.K. 86, 903-907.

Capellini, G., 1876. Sulle balene fossili toscane. Atti della Reale Accademia dei Lincei 3, 9-14.

Capellini, G., 1883. Di un'Orca fossile scoperta a Cetona in Toscana. Memorie dell'Accademia delle Scienze dell'Istituto di Bologna 4, serie 4, 1-25.

Capellini, G. 1905. Balene fossili toscane - III. *Idiocetus guicciardinii*. Memorie dell'Accademia delle Scienze dell'Istituto di Bologna 2, serie 6, 71-80.

Cavanagh, R.D., Gibson, C., 2007. Overview of the conservation status of cartilaginous fishes (Chondrichthyans) in the Mediterranean Sea. Gland, Switzerland and Malaga, Spain: The World Conservation Union (IUCN), vi + 42 pp.

Chen, Z.-Q., Benton, M.J., 2012. The timing and pattern of biotic recovery following the end-Permian mass extinction. *Nature Geoscience* 5, 375-383.

Cigala Fulgosi, F., 1990. Predation (or possible scavenging) by a great white shark on an extinct species of bottle-nosed dolphin in the Italian Pliocene. *Tertiary Research*, 12, 17-6.

Cigala Fulgosi, F., Casati, S., Orlandini, A., Persico, D., 2009. A small fossil fish fauna, rich in *Chlamydoselachus* teeth, from the Late Pliocene of Tuscany (Siena, central Italy). *Cainozoic Research* 6, 3-23.

Cioppi, E., 2014. I cetacei fossili a Firenze, una storia lunga più di 250 anni. *Museologia Scientifica Memorie* 13, 81-89.

Cioppi, E., Dominici, S., 2011. Pliocene marine vertebrates, in: Monechi, S., Rook, L. (Eds.), *The Museum of Natural History of the University of Florence, Volume 3, The geological and paleontological collections: Firenze, Firenze University Press*, 224-235.

Clarke, K.R., Gorley, R.N., 2006. *PRIMER v6: User Manual/Tutorial*. Plymouth: PRIMER-E., 190 p.

Cockcroft, V.G., Cliff, C., Ross, G.J.B., 1989. Shark predation on Indian Ocean bottlenose dolphins *Tursiops truncatus* off Natal, South Africa. South African Journal of Zoology 24, 305-310.

Collareta, A., Bosselaers, M. & Bianucci, G., 2016. Jumping from turtles to whales: a Pliocene fossil record depicts an ancient dispersal of *Chelonibia* on mysticetes. Rivista Italiana di Paleontologia e Stratigrafia 122, 35-44.

Compagno, L.J.V., 2002. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Volume 2. Bullhead, mackerel and carpet sharks (Heterodontiformes, Lamniformes and Orectolobiformes). FAO Species Catalogue for Fishery Purposes 1 (2) Rome, 269 p.

Cortés, E., 1999. Standardized diet compositions and trophic levels of sharks. ICES Journal of Marine Science 56, 707–717, doi: 10.1006/jmsc.1999.0489.

Croll, D.A., Marinovic, B., Benson, S., Chavez, F.P., Black, N., Ternullo, R., Tershy, B.R., 2005. From wind to whales: trophic links in a coastal upwelling system. Marine Ecology Progress Series 289, 117-130.

Cuzin-Roudy, J., 2011. Reproductive strategies of the Mediterranean krill, *Meganyctiphanes norvegica* and the Antarctic krill, *Euphausia superba* (Crustacea: Euphausiacea). Invertebrate Reproduction & Development 23, 105-114, doi: 10.1080/07924259.1993.9672301.

Danise, S., 2010. Modern and fossil shallow water whale fall communities. [FirenzeUniversity of Florence](#), PhD Thesis, 178 pp.

Danise, S., Dominici, S., 2014. A record of fossil shallow-water whale falls from Italy. *Lethaia*, 47, 229–243.

Danise, S., Higgs, N.D., 2015. Bone-eating *Osedax* worms lived on Mesozoic marine reptile deadfalls. *Biology Letters* 11: 20150072.

Danise, S., Holland, S.M., 2017. Faunal response to sea-level and climate change in a short-lived seaway: Jurassic of the Western Interior, USA. *Palaeontology*, doi: 10.1111/pala.12278.

Danise, S., Twitchett, R.J., Matts, K., 2014. Ecological succession of a Jurassic shallow-water ichthyosaur fall. *Nature Communications* 5: 4789, doi: 10.1038/ncomms5789 | www.nature.com/naturecommunications.

DeCelles, P.G., 2012. Foreland basin systems revisited: variations in response to tectonic settings. In: C., Busby, & A., Azor (eds.), *Tectonics of Sedimentary Basins: Recent Advances*. Hoboken: Wiley-Blackwell, 405-426.

De Stefani, C., 1874. I terreni subapennini dei dintorni di S. Miniato al Tedesco. *Bollettino Malacologico Italiano* 7, 5-88.

Di Celma, C., Malinverno, E., Gariboldi, K., Gioncada, A., Rustichelli, A., Pierantoni, P.P., Bianucci, G., 2015. Stratigraphic framework of the late Miocene to Pliocene Pisco formation

at Cerro Colorado (Ica Desert, Peru). Journal of Maps.
doi:10.1080/17445647.2015.1047906.

Dicken, M.L., 2008. First observations of young of the year and juvenile great white sharks (*Carcharodon carcharias*) scavenging from a whale carcass. Marine and Freshwater Research 59, 596-602.

Dineen, A.A., Fraiser, M.L., Sheehan, P.M., 2014. Quantifying functional diversity in pre- and post-extinction paleocommunities: A test of ecological restructuring after the end-Permian mass extinction. Earth-Science Reviews 136, 339-349.

Dominici, S., 1994. Regressive-transgressive cycles from the Pliocene of the San Miniato area (Tuscany, Italy): paleoecology and sequence stratigraphy. Bollettino della Società Paleontologica Italiana Supplement 2, 117-126.

Dominici, S., Rook, L., Benvenuti, M., Abbazzi, L., 1995. Tapir remains in paralic deposits of Pliocene age in lower Valdarno (Tuscany, Italy): Facies analysis and taphonomy. Geobios 28, 131-135.

Dominici, S., Conti, C., Benvenuti, M., 2008. Foraminifer communities and environmental change in marginal marine sequences (Pliocene, Tuscany, Italy). Lethaia 41, 447-460.

Dominici, S., Cioppi, E., Danise, S., Betocchi, U., Gallai, G., Tangocci, F., Valleri, G., Monechi, S., 2009. Mediterranean fossil whale falls and the adaptation of mollusks to extreme habitats. Geology 37, 815–818.

Domning, D.P., 2001, Sirenians, seagrasses, and Cenozoic ecological change in the Caribbean. Palaeogeography, Palaeoclimatology, Palaeoecology 166, 27-50.

Dowsett, H.J., Robinson, M.M., Stoll, D.K., Foley, K.M., Johnson, A.L.A., Williams, M., Riesselman, C.R., 2013. The PRISM (Pliocene Palaeoclimate) reconstruction: Time for a paradigm shift, Philosophical Transactions of the Royal Society, 371(2001): doi:10.1098/rsta.2012.0524.

Ellis, J.A., Musick, J.A., 2007. Ontogenetic changes in the diet of the sandbar shark, Carcharhinus plumbeus, in lower Chesapeake Bay and Virginia (USA) coastal waters. Environmental Biology and Fishing 80, 51–67, doi: 10.1007/s10641-006-9116-2.

Estes, J.A., Burdin, A., Doak, D.F., 2016. Sea otters, kelp forests, and the extinction of Steller's sea cow. PNAS 113, 880–885, doi: 10.1073/pnas.1502552112.

Estes, J.A., Terborgh, J., Brashares, J.S., Power, M.E., Berger, J., Bond, W.J., Carpenter, S.R., Essington, T.E., Holt, R.D., Jackson, J.B.C., Marquis, R., Oksanen, L., Oksanen, T., Paine, R.T., Pikitch, E.K., Ripple, W.J., Sandin, S.A., Scheffer, M., Schoener, T.W., Shurin, J.B., Sinclair, A.R.E., Soulé, M.E., Virtanen, R., Wardle, D.A., 2011. Trophic downgrading of Planet Earth. Science 333, 301-306, doi: 10.1126/science.1205106.

Estes, J.A., Tinker, M.T., Williams, T.M., Doak, D.F. 1998. Killer whale predation on sea otters linking oceanic and nearshore ecosystems. Science 282, 473–476.

Fischer, V., Bardet, N., Benson, R.B.J., Arkhangelsky, M.S., Friedman, M., 2016. Extinction of fish-shaped marine reptiles associated with reduced evolutionary rates and global environmental volatility. *Nature Communications* 7, 10825, doi: 10.1038/ncomms10825.

Fordyce, R.E. 2003. Cetacean evolution in Eocene-Oligocene oceans revisited. in *From Greenhouse to Icehouse: The Marine Eocene-Oligocene Transition*, D.R. Prothero, L.C. Ivany, E.A. Nesbitt, Eds. (Columbia Univ. Press, New York, 2003), 154–170.

Friedlaender, A.S., Goldbogen, J.E., Hazen, E.L., Calambokidis, J., Southall, B.L., 2014. Feeding performance by sympatric blue and fin whales exploiting a common prey resource. *Mar. Mamm. Sci.* 31, 345–354.

Fröbisch, N.B., Fröbisch, J., Martin Sander, P., Schmitz, L., Rieppel, O., 2013. Macropredatory ichthyosaur from the Middle Triassic and the origin of modern trophic networks. *Proceedings of the National Association of Science* 110, 1393-1397, doi:10.1073/pnas.1216750110.

Garassino, A., Pasini, G., De Angeli, A., Charbonnier, S., Famiani, F., Baldanza, A., Bizzarri, R., 2012. The decapod community from the Early Pliocene (Zanclean) of “La Serra” quarry (San Miniato, Pisa, Toscana, central Italy): sedimentology, systematics, and palaeoenvironmental implications. *Annales de Paléontologie* 98, 1–61.

Gattacceca, J., Deino, A., Rizzo, R., Jones, D.S., Henry, B., Beaudoin, B., Vadeboin F., 2007. Miocene rotation of Sardinia: New paleomagnetic and geochronological constraints and geodynamic implications. *Earth and Planetary Science Letters* 258, 359-377.

Ghinassi, M., 2007. Pliocene alluvial to marine deposits of the Val d'Orcia Basin (Northern Apennines, Italy): sequence stratigraphy and basin analysis. *Rivista Italiana di Paleontologia e Stratigrafia* 113, 459-472.

Ghinassi, M., Lazzarotto, A., 2005. Different source areas feeding the Early pliocene marine Radicofani Basin (Siena): geological and paleogeographical implications. *Bollettino della Società Geologica Italiana*, volume speciale 3, 15-28.

Gingerich, P.D., 2005. Cetacea. In: Rose K.D., Archibald J.D., editors. *Placental mammals: origin, timing, and relationships of the major extant clades*. Baltimore (MD), Johns Hopkins University Press. p. 234-252.

Gingerich, P.D., Ul-Haq, M., von Koenigswald, W., Sanders, W.J., Smith, B.H., Zalmout, I.S., 2009. New protocetid whale from the middle Eocene of Pakistan: birth on land, precocial development, and sexual dimorphism. *PloS One*. 4:e4366.

Goffredo, S., Dubinsky, Z., 2014. The Mediterranean Sea: Its history and present challenges. Springer-Verlag, Berlin, pp. 678.

Goldbogen, J.A., Calambokidis, J., Oleson, E., Potvin, J., Pyenson, N.D., Schorr, G., Shadwick, R.E., 2011. Mechanics, hydrodynamics and energetics of blue whale lunge feeding: efficiency dependence on krill density. *The Journal of Experimental Biology* 214, 131-146.

Greenpeace International, 2009. High seas Mediterranean marine reserves: a case study for the Southern Balearics and the Sicilian Channel. A briefing to the CBD's Expert workshop

[on scientific and technical guidance on the use of biogeographic classification systems and identification of marine areas beyond national jurisdiction in need of protection, Ottawa, 29 September–2 October 2009.](#)

Guinot, G., Adnet, S., Cappetta, H., 2012. An analytical approach for estimating fossil record and diversification events in sharks, skates and rays. PLoS ONE 7(9): e44632. doi:10.1371/journal.pone.0044632.

[Heyning, J.E., Dahlheim, M.E., 1988. *Orcinus orca*. Mammalian Species 304, 1–9, doi:10.2307/3504225.](#)

Higgs, N.D., Glover, A.G., Dahlgren, T.G., Little, C.T.S., 2011. Bone-boring worms: characterizing the morphology, rate, and method of bioerosion by *Osedax mucofloris* (Annelida, Siboglinidae). Biological Bulletin 221:307-316.

Higgs, N.D., Little, C.T.S., Glover, A.G., Dahlgren, T.G., Smith, C.R., Dominici, S. 2012. Evidence of *Osedax* worm borings in Pliocene (~3 Ma) whale bone from the Mediterranean. Historical Biology 24, 269-277.

Higgs, N.D., Gates, A.R., Jones, D.O.B., 2014a. Fish food in the deep sea: revisiting the role of large food-falls. PLOS ONE 9(5): e96016

Higgs, N.D., Glover, A.G., Dahlgren, T.G., Smith, C.R., Fujiwara, Y., Pradillon, F., Johnson, S.B., Vrijenhoek, R.C., Little, C.T.S., 2014b. The morphological diversity of *Osedax* worm borings (Annelida: Siboglinidae). Journal of the Marine Biological Association of the United Kingdom [94](#), 1429–1439, doi: [10.1017/S0025315414000770](#).

Hocking, D.P., Marx, F.G., Park, T., Fitzgerald, E.M.G., Evans, A.R., 2017 A behavioural framework for the evolution of feeding in predatory aquatic mammals. *Proc. R. Soc. B* 284, 20162750, doi: 10.1098/rspb.2016.2750.

Hogler, J.A., 1992. Taphonomy and paleoecology of *Shonisaurus popularis* (Reptilia: Ichthyosauria). *Palaios* 7, 108-117.

Hu, S., Zhang, Q., Chen, Z.-Q., Zhou, C., Lü, T., Xie, T., Wen, W., Huang, J., Benton, M.J., 2011. The Luoping biota: exceptional preservation, and new evidence on the Triassic recovery from end-Permian mass extinction. *Proceedings of the Royal Society B*, 278, 2274–2282, doi:10.1098/rspb.2010.2235.

Huusgaard, R.S., Vismann, B., Kühl, M., Macnaughton, M., Colmander, V., Rouse, G.W., Glover, A.G., Dahlgren, T., Worsaae, K., 2012. The potent respiratory system of *Osedax mucofloris* (Siboglinidae, Annelida) - a prerequisite for the origin of bone-eating *Osedax*? *PLoS ONE* 7(4): e35975. doi:10.1371/journal.pone.0035975.

Jagt, J.W.M, Jagt-Yazykova, E.A., 2016. The Upper Cretaceous and lower Paleogene in the type area of the Maastrichtian stage (72.1–66 Ma). Field guide of the 14th Annual Meeting of the European Association of Vertebrate Paleontologists, 18 pp.

Jenkins, R.G., Kaim, A., Sato, K., Moriya, K., Hikida, Y., and Hirayama, R. 2017. Discovery of chemosynthesis-based association on the Cretaceous basal leatherback sea turtle from Japan. *Acta Palaeontologica Polonica* 62, doi: 10.4202/app.00405.2017.

Johnson, S.B., Warén, A., Lee, R.W., Kano, Y., Kaim, A., Davis, A., Strong, E.E., Vrijenhoek, R.C., 2010. *Rubyspira*, new genus and two new species of bone-eating deep-sea snails with ancient habits. *Biological Bulletin* 219: 166-177.

Jorgensen, S.J., Reeb, C.A., Chapple, T.K., Anderson, S., Perle, C., Van Sommeran, S.R., Fritz-Cope, C., Brown, A.C., Klimley, A.P., Block, B.A., 2010. Philopatry and migration of Pacific white sharks. *Proceedings of the Royal Society B* 277, 679-688.

Kaim, A., Kobayashi, Y., Echizenya, H., Jenkins, R., Tanabe, K., 2008. Chemosynthesis-based associations on Cretaceous plesiosaurid carcasses. *Acta Palaeontologica Polonica* 53: 97–104.

Kiel, S., Goedert, J.L., 2006. Deep-sea food bonanzas: Early Cenozoic whale-fall communities resemble wood-fall rather than seep communities. *Proc. R. Soc. B* 273, 2625–2631, doi: 10.1098/rspb.2006.3620.

Krijgsman, W., Hilgen, F.J., Raffi, I., Sierro, F.J., Wilson, D.S., 1999. Chronology, causes and progression of the Messinian salinity crisis. *Nature*, 400, 652-655.

Kriwet, J., Benton, M.J., 2004. Neoselachian (Chondrichthyes, Elasmobranchii) diversity across the Cretaceous–Tertiary boundary. *Palaeogeography, Palaeoclimatology, Palaeoecology* 214. 181-94.

Lambert, O., Bianucci, G., Post, K., de Muizon, C., Salas-Gismondi, R., Urbini, M., Reumer, J., 2010. The giant bite of a new raptorial sperm whale from the Miocene epoch of Peru. *Nature* 466, 105-108, doi: 10.1038/nature09067

Landini, W., 1977. Revisione degli 'Ittiodontoliti pliocenici' della collezione Lawley. *Palaeontographia Italica* 70, 92-134.

Landini, W., Bianucci, G., Bisconti, M., Carnevale, G., Sorbini, C., 2005. Il Plio-Pleistocene. I vertebrati marini. In Bonfiglio, L., (ed.) *Paleontologia dei vertebrati in Italia. Evoluzione biologica, significato ambientale e paleogeografia*. Museo Civico di Storia Naturale, Verona, 171-175.

Lawley, R., 1876. Nuovi studi sopra i pesci ed altri vertebrati fossili delle colline Toscane. Tipografia dell'Arte della Stampa, Firenze, 122 p.

[Lawton, J.H., 1994. What do Species do in ecosystems? *Oikos* 71, 367-374.](#)

Lazzari, P., Solidoro, C., Ibello, V., Salon, S., Teruzzi, A., Béranger, K., Colella, S., Crise, A. 2012. Seasonal and inter-annual variability of plankton chlorophyll and primary production in the Mediterranean Sea: a modelling approach. *Biogeosciences* 9, 217-233.

[Lewison, R.L., Crowder, L.B., Read, A.J., Freeman, S.A., 2004. Understanding impacts of fisheries bycatch on marine megafauna. *Trends Ecol. Evol.* 19, 598–604.](#)

Lindsay, E.H., Opdyke, N.D., Johnson, N.M., 1980. Pliocene dispersal of the horse *Equus* and late Cenozoic mammalian dispersal events. *Nature* 287, 135-138.

Liotta, D., 1996. Analisi del settore centro-meridionale del bacino pliocenico di Radicofani (Toscana meridionale). Bollettino della Società Geologica Italiana 115, 115-143.

Liu, J. Hu, S., Rieppel, O., Jiang, D., Benton, M.J., Kelley, N.P., Aitchison, J.C., Zhou, C., Wen, W., Huang, J., Xie, T., Lv, T., 2014. A gigantic nothosaur (Reptilia: Sauropterygia) from the Middle Triassic of SW China and its implication for the Triassic biotic recovery. Scientific Reports 4, 7142, [DOI:10.1038/srep07142](https://doi.org/10.1038/srep07142).

[Lourenço, R., Penteriani, V., Rabac, J.E., Korpimäki, R., 2013. Lethal interactions among vertebrate top predators: a review of concepts, assumptions and terminology. Biol. Rev. doi: 10.1111/brv.12054.](#)

[Louwyse, S., Marquet, R., Bosselaers, M., Lambert, O., 2010. Stratigraphy of an early-middle Miocene sequence near Antwerp in Northern Belgium \(Southern North Sea Basin\). Geologica Belgica, 13, 269-284.](#)

Lundsen, L., Schlining, K.L., Frasier, K., Johnson S.B., Kuhnz, L.A., Harvey, J.B.J., Clague, G., Vrijenhoek, R.C., 2010. Time-series analysis of six whale-fall communities in Monterey Canyon, California, USA. Deep-Sea Research I 57, 1573–1584.

[Manganelli, G., Benocci, A., 2014. I cetacei fossili del Museo di Storia Naturale dell'Accademia dei Fisiocritici di Siena. Museologia Scientifica Memorie 13, 103-110.](#)

Mann, K.H., Lazier, J.R.N., 2006. Dynamics of marine ecosystems : biological-physical interactions in the oceans. Third edition. Blackwell Publishing, 496 pp.

Marsili, S., 2006. Analisi Ssistemica, Ppaleoecologica e Ppaleobiogeografica della Sselaciofauna plio-pleistocenica del Mediterraneo. Pisa, PhD Thesis, 246 pp.

Marsili, S., 2007. Revision of the teeth of the genus *Carcharhinus* (Elasmobranchii; Carcharhinidae) from the Pliocene of Tuscany, Italy. *Rivista Italiana di Paleontologia e Stratigrafia* 113 (1), 79–96.

Marsili, S., 2008. Systematic, paleoecologic and paleobiogeographic analysis of the Plio-Pleistocene Mediteranean elasmobranch fauna. *Atti della Società Toscana di Scienze Naturali, Serie A*, 113, 81-88.

Martill, D.M., 1985. The preservation of marine vertebrates in the Lower Oxford Clay (Jurassic) of central England. *Philosophica Transactions of the Royal Society B* 311, 155-165.

Martill, D. M., 1987. A taphonomic and diagenetic case study of a partially articulated ichthyosaur. *Palaeontology* 30, 543-555.

Martill, D. M., Taylor, M. A., Duff, K. L., Riding, J. B., Bown, P. R., 1994. The trophic structure of the biota of the Peterborough Member, Oxford Clay Formation (Jurassic), UK. *Journal of the Geological Society* 151, 173-194.

Martini, I., Aldinucci, M., Foresi, L.M., Mazzei, R., Sandrelli, F., 2011. Geological map of the Pliocene succession of the Northern Siena Basin (Tuscany, Italy). *Journal of Maps* 2011, 193-204.

Martini, I., Foresi, L.M., Bambini, A.M., Riforgiato, F., Ambrosetti, E., Sandrelli, F., 2016. Calcareous plankton bio-chronostratigraphy and sedimentology of the “I Sodi” section (Siena Basin, Italy): a key section for the uppermost Neogene marine deposition in the inner northern Apennines. Ital. J. Geosci. 135, 540-547, doi: 10.3301/IJG.2016.02.

Marx, F., G.M., Collareta, A., Gioncada, A., Post, K., Lambert, O., Bonaccorsi, E., Urbina, M., Bianucci, G., 2017. How whales used to filter: exceptionally preserved baleen in a Miocene cetotheriid. Journal of Anatomy 231, 212-220, doi: 10.1111/joa.12622.

Marx, G.M., Lambert, O., Uhen, M.D., 2016. Cetacean Palaeobiology. Topics in Paleobiology, Wiley-Blackwell, 336 pp.

Marx, F.G., Uhen, M.D., 2010. Climate, Critters, and Cetaceans: Cenozoic dDrivers of the eEvolution of mModern wWhales. Science 327, 993-996.

Melanotte-Rizzoli, P., Artale, V., Borzelli-Eusebi, G.L., Brenner, S., Crise, A., et al., 2014. Physical forcing and physical/biochemical variability of the Mediterranean Sea: a review of unresolved issues and directions for future research. Ocean Science 10, 281-322.

Meyer, C.A., 2011. Amazing graze – Grazing traces of sea urchins on turtles – An example from the Late Jurassic of Switzerland: Annalen des Naturhistorisches Museum in Wien 113, 555-565.

Miyamoto, N., Yoshida, M.A., Koga, H., Fujiwara, Y., 2017. Genetic mechanisms of bone digestion and nutrient absorption in the bone-eating worm *Osedax japonicus* inferred from

transcriptome and gene expression analyses. BMC Evolutionary Biology 17: 17, doi:
[10.1186/s12862-016-0844-4](https://doi.org/10.1186/s12862-016-0844-4).

Motani, R., Jiang, D.-Y., Tintori, A., Sun, Y.-L., S., Hao, W.C., Boyd, A., Hinic-Frlog, S.,
Schmitz, L., Shin, J.-Y. & Sun Z.-Y. 2008. Horizons and assemblages of Middle Triassic
marine reptiles from Panxian, Guizhou, China. Journal of Vertebrate Paleontology 28, 900-
903.

~~Miyamoto, N., Yoshida, M.A., Koga, H., Fujiwara, Y., 2017. Genetic mechanisms of bone
digestion and nutrient absorption in the bone-eating worm *Osedax japonicus* inferred from
transcriptome and gene expression analyses. BMC Evolutionary Biology 17:17. doi:
[10.1186/s12862-016-0844-4](https://doi.org/10.1186/s12862-016-0844-4)~~

Motani, R., Ji, C., Tomita, T., Kelley, N., Maxwell, E., Jiang, D.-Y., Sanders, M., 2013.
Absence of suction feeding ichthyosaurs and its implications for Triassic mesopelagic
paleoecology. PLoS ONE 8(12): e66075. doi:10.1371/journal.pone.0066075.

Motani, R., Jiang, D.-Y., Tintori, A., Rieppel, O., Chen, G.-B., 2014. Terrestrial origin of
viviparity in Mesozoic marine reptiles indicated by Early Triassic embryonic fossils. PLoS
ONE 9(2): e88640. doi:10.1371/journal.pone.0088640.

Motani, R., Chen, X., Jiang, D., Cheng, L., Tintori, A., Rieppel, O., 2015. Lunge feeding in
early marine reptiles and fast evolution of marine tetrapod feeding guilds. Scientific Reports
5, 8900. doi:10.1038/srep08900.

Moulins, A., Rosso, M., Nani, B., Würtz, M., 2007. Aspects of the distribution of Cuvier's beaked whale (*Ziphius cavirostris*) in relation to topographic features in the Pelagos Sanctuary (north-western Mediterranean Sea). *Journal of the Marine Biological Association of the United Kingdom* 87, 177-186, doi: [10.1017/S0025315407055002](https://doi.org/10.1017/S0025315407055002).

Murakami, M., Shimada, C., Hikida, Y., Soeda, Y., Hirano, H. 2014. *Eodelphis kabatensis*, a new name for the oldest true dolphin *Stenella kabatensis* Horikawa, 1977 (Cetacea, Odontoceti, Delphinidae), from the upper Miocene of Japan, and the phylogeny and paleobiogeography of Delphinoidea, *Journal of Vertebrate Paleontology* 34, 491-511, doi: [10.1080/02724634.2013.816720](https://doi.org/10.1080/02724634.2013.816720).

Mutti, E., Davoli, G., Mora, S., Sgavetti, M., 1994. The eastern sector of the south-central folded Pyrenean foreland: criteria for stratigraphic analysis and excursion notes. 2nd International conference on high-resolution sequence stratigraphy, Guidebook. Istituto di Geologia, Università di Parma, Italy.

Muttoni G., Lanci L., Argnani A., Hirt A.M., Cibin U., Abrahamsen N., Lowrie W., 2001. Paleomagnetic evidence for a Neogene two-phase counterclockwise tectonic rotation in the Northern Apennines (Italy). *Tectonophysics* 326, 241-253-

Nalin R., Ghinassi M., Basso D., 2010. Onset of temperate carbonate sedimentation during transgression in a low-energy siliciclastic embayment (Pliocene of the Val d'Orcia Basin, Tuscany, Italy). *Facies* 56, 353-368.

Nalin, R., Ghinassi, M., Foresi, L., Dallanave, E., 2016. Carbonate deposition in restricted basins: a Pliocene case study from the central Mediterranean (Northwestern Apennines). *Journal of Sedimentary Research* 86, 1-32.

[Navarro, J., López, L., Coll, M., Barría, C., Sáez-Liante, R., 2014. Short- and long-term importance of small sharks in the diet of the rare deep-sea shark *Dalatias licha*. *Mar Biol* 161, 1697-1707, doi: 10.1007/s00227-014-2454-2.](#)

Noakes, S.E., Pyenson, N.D., McFall, G., 2013. Late Pleistocene gray whales (*Eschrichtius robustus*) offshore Georgia, U.S.A., and the antiquity of gray whale migration in the North Atlantic Ocean. *Palaeogeography, Palaeoclimatology, Palaeoecology* 392, 502-509.

[Notarbartolo di Sciara, G., 1987. Killer whale, *Orcinus orca*, in the Mediterranean Sea. *Marine Mammal Science* 3, 356-360.](#)

Notarbartolo di Sciara, G., Zanardelli, M., Jahoda, M., Panigada, S., Airoldi, S. 2003. The fin whale *Balaenoptera physalus* (L. 1758), in the Mediterranean Sea. *Mammal Review* 33, 105-150.

Notarbartolo di Sciara, G., Agardy, T., Hyrenbach, D., Scovazzi, T., Van Klaveren, P., 2008. The Pelagos sanctuary for Mediterranean marine mammals. *Aquatic Conservation: Marine and Freshwater Ecosystems* 18, 367-391.

[Notarbartolo di Sciara, G., Podestà, M., Curry, B.E. \(Eds.\), 2016a, Mediterranean marine mammals ecology and conservation. *Advances in Marine Biology* 75, 458 pp.](#)

Notarbartolo di Sciara, G., Castellote, M., Druon, J.-N., Panigada, S., 2016b. Fin whales, *Balaenoptera physalus*: at home in a changing Mediterranean Sea?. In: G. Notarbartolo Di Sciara, M. Podestà, B.E. Curry, eds., *Advances in Marine Biology* 75, 75-101, doi: [10.1016/bs.amb.2016.08.002](https://doi.org/10.1016/bs.amb.2016.08.002).

Panigada, S., Notarbartolo di Sciara, G., 2012. *Balaenoptera physalus* (Mediterranean subpopulation). The IUCN Red List of Threatened Species 2012, e.T16208224A17549588, doi: [10.2305/IUCN.UK.2012.RLTS.T16208224A17549588.en](https://doi.org/10.2305/IUCN.UK.2012.RLTS.T16208224A17549588.en).

Parrish, F.A., 2009. Do monk seals exert top-down pressure in subphotic ecosystems? *Mar. Mamm. Sci.* 25, 91-106, doi: [10.1111/j.1748-7692.2008.00245.x](https://doi.org/10.1111/j.1748-7692.2008.00245.x).

Pascucci, V., Costantini, A., Martini, I.P., Dringoli, R., 2006. Tectono-sedimentary analysis of a complex, extensional, Neogene basin formed on thrust-faulted, Northern Apennines hinterland: Radicofani Basin, Italy. *Sedimentary Geology*, 183, 71-97.

Patzkowsky, M.E., Holland S.M., 2012. Stratigraphic paleobiology. Understanding the distribution of fossil taxa in time and space. Chicago: University of Chicago Press 256 pp.

Pauly, D., Trites, A.W., Capuli, E., Christensen, V., 1998. Diet composition and trophic levels of marine mammals. *ICES Journal of Marine Science*, 55, 467-481.

Pilleri, G., 1987. The cetacea of the Italian Pliocene with a descriptive catalogue of the species in the Florence Museum of Paleontology. Ostermundigen: Brain Anatomy Institute, University of Berne, Berne, 160 p.

Pimiento, C., Griffin, J.N., Clements, C.F., Silvestro, D., Varela, S., Uhen, M.D., Jaramillo, C., 2017. The Pliocene marine megafauna extinction and its impact on functional diversity. *Nature Ecology & Evolution*, doi: [10.1038/s41559-017-0223-6](https://doi.org/10.1038/s41559-017-0223-6).

Pirotta, E., Matthiopoulos, J., MacKenzie, M., Scott-Hayward L., Rendell, L., 2011. Modelling sperm whale habitat preference: a novel approach combining transect and follow data. *Marine Ecology Progress Series* 436, 257-272, do: [10.3354/meps09236](https://doi.org/10.3354/meps09236).

Podestà, M., Azzellino, A., Cañadas, A., Frantzis, A., Moulins, A., Rosso, M., Tepsich, P., Lanfredi, C., 2016. Cuvier's beaked whale *Ziphius cavirostris*, distribution and occurrence in the Mediterranean Sea: high-use areas and conservation threats. In: Notarbartolo di Sciara, G., Podestà, M., Curry, B.E. (Eds.), *Mediterranean Marine Mammals Ecology and Conservation*. *Advances in Marine Biology* 75, 105–142, doi: [10.1016/bs.amb.2016.07.007](https://doi.org/10.1016/bs.amb.2016.07.007).

Pokines, J.T., Symes, S.A., 2013. *Manual of forensic taphonomy*. CRC Press, Boca Raton, FL, 496 p.

Polis, G.A., Anderson, W.B., Holt, R.D., 1997. Toward an integration of landscape and food web ecology: The dynamics of spatially subsidized food webs. *Annual Review of Ecology and Systematics* 28, 289-316.

Polis, G.A., Myers, C.A., Holt, R.D., 1989. The ecology and evolution of intraguild predation: potential competitors that eat each other. *Annual Review of Ecology and Systematics* 20, 297-330, doi: [10.1146/annurev.es.20.110189.001501](https://doi.org/10.1146/annurev.es.20.110189.001501).

Praca, E., Gannier, A., 2008. Ecological niches of three teuthophageous odontocetes in the northwestern Mediterranean Sea. *Ocean Science* 4, 49–59. <http://dx.doi.org/10.5194/os-4-49-2008>.

Prista, G.A., Agostinho, R.J., Cachão, M.A., 2015. Observing the past to better understand the future: a synthesis of the Neogene climate in Europe and its perspectives on present climate change. *Open Geoscience* 7, 65-83.

Pyenson, N.D., Haasl, D.M., 2007. Miocene whale-fall from California demonstrates that cetacean size did not determine the evolution of modern whale-fall communities. *Biology Letters (Palaeontology)* 3, 709-711, doi: 10.1098/rsbl.2007.0342.

Pyenson, N.D., Lindberg, D.R., 2011. What happened to gray whales during the Pleistocene? The ecological impact of sea-level change on benthic feeding areas in the North Pacific Ocean. *PLoS ONE* 6(7): e21295. doi:10.1371/journal.pone.0021295

Raffi, S., Stanley, S., Marasti, R., 1985. Biogeographic patterns and Plio-Pleistocene extinction of *Bivalvia* in the Mediterranean and southern North Sea. *Paleobiology* 11, 368-388.

Raymo, M.E., Hearty, P., De Conto, R., O'Leary, M., Dowsett, H.J., Robinson, M.M., Mitrovica, J.X., 2009. PLIOMAX: Pliocene maximum sea level project. *PAGES News* 17(2), 58-59.

Rendell, L., Frantzis, A. 2016. Mediterranean sperm whales, *Physeter macrocephalus*: the precarious state of a lost tribe. In: Notarbartolo di Sciara, G., Podestà, M., Curry, B.E. (Eds.), *Mediterranean Marine Mammals Ecology and Conservation. Advances in Marine Biology*. 37-74, doi: [10.1016/bs.amb.2016.08.001](https://doi.org/10.1016/bs.amb.2016.08.001).

Reolid, M., Santos, A., Mayoral, E., 2015. Grazing activity as taphonomic record of necrobiotic interaction: A case study of a sea turtle carapace from the Upper Jurassic of the Prebetic (south Spain). *Revista Mexicana de Ciencias Geológicas* 32(1), 21-28.

Roemer, G.W., Gompfer, M.E., Van Valkenburgh, B., 2009. The ecological role of the mammalian mesocarnivore. *BioScience* 59, 165-173, doi: [10.1525/bio.2009.59.2.9](https://doi.org/10.1525/bio.2009.59.2.9).

Roman, J., Estes, J.A., Morissette, L., Smith, C., Costa, D., McCarthy, J.B., Nation, S.N., Pershing, A., Smetacek, V., 2014. Whales as marine ecosystem engineers. *Frontiers in Ecology and the Environment* 12 (7), 15 pp.

Roman, J., McCarthy, J.J., 2010. The whale pump: marine mammals enhance primary productivity in a coastal basin. *PLoS ONE* 5: e13255.

Roveri, M., Flecker, R., Krijgsman, W., Lofi, J., Lugli, S., Manzi, V., Sierro, F.J., Bertini, A., Camarlenghi, A., De Lange, G., Govers, R., Hilgen, F.J., Hübscher, C., Meijer, P.T.H., Stoica, M., 2014. The Messinian Salinity Crisis: Past and future of a great challenge for marine sciences. *Marine Geology* 352, 25-58.

Sarti, C., Lanzetti, A., 2014. I cetacei fossili del museo geologico Giovanni Capellini dell'università di Bologna. *Museologia Scientifica Memorie*, 70-78.

Scarponi, D., Kowalewski, M., 2004. Stratigraphic paleoecology: bathymetric signatures and sequence overprint of mollusk associations from upper Quaternary sequences of the Po Plain, Italy. *Geology* 32, 989-992.

Scheyer, T.M., Romano, C., Jenks, J., Bucher, H., 2014. Early Triassic Marine Biotic Recovery: The Predators' Perspective. *PLoS ONE* 9(3): e88987. doi:10.1371/journal.pone.0088987.

Schmeisser McKean, R.L., Gillette, D.D., 2015. Taphonomy of large marine vertebrates in the Upper Cretaceous (Cenomanian-Turonian) Tropic Shale of southern Utah. *Cretaceous Research* 56, 278-292.

Sergio, F., Schmitz, O.J., Krebs, C.J., Holt, R.D., Heithaus, M.R., Wirsing, A.J., Ripple, W.J., Ritchie, E., Ainley, D., Oro, D., Jhala, Y., Hiraldo, F., Korpimäki, E., 2014. Towards a cohesive, holistic view of top predation: a definition, synthesis and perspective. *Oikos* 123, 1234-1243.

Slater, G.J., Goldbogen, J.A., Pyenson, N.D., 2017. Independent evolution of baleen whale gigantism linked to Plio-Pleistocene ocean dynamics. *Proc. R. Soc. B* 284, 20170546. <http://dx.doi.org/10.1098/rspb.2017.0546>.

Smith, C., Glover, A.G., Treude, T., Higgs, N.D., & Amon, D.J. 2015. Whale-Fall Ecosystems: Recent Insights into Ecology, Paleoecology, and Evolution. Annual Review of Marine Science 7: 10.1-10.26.

Smith, F.A., Payne, J.L., Heim, N.A., Balk, M.A., Finnegan, S., Kowaleski, M., Lyons, K., McClain, C.R., McShea, D.W., Novack-Gottshall, P.M., Spaeth Anich, P., Wang, S.C., 2016, Body Size Evolution Across the Geozoic. Annu. Rev. Earth Planet. Sci. 44, 523-53, doi: 10.1146/annurev-earth-060115-012147.

Sorbi, S., Domning, D.P., Vaiani, S.C., Bianucci, G., 2012. *Metaxytherium subapenninum* (Bruno, 1839) (Mammalia, Dugongidae), the latest sirenian of the Mediterranean Basin. Journal of Vertebrate Paleontology 32, 686-707.

Sorbini, C., Bianucci, G., Baldassarri, M., 2014. La balena fossile di Montopoli in Val d'Arno (Pisa). Museologia Scientifica, Memorie 13, 90-92.

~~Sorbi, C., Bianucci, G., Baldassari, M., 2014. La balena fossile di Montopoli in Val d'Arno (Pisa). Museologia Scientifica 13, 90-92.~~

Stambler, N., 2014. The Mediterranean Sea – Primary productivity. In: Goffredo, S., Dubinsky, Z., eds., The Mediterranean Sea: Its history and present challenges. Springer-Verlag, Berlin, 113-121.

Stanley, S., 1975. Adaptive themes in the evolution of the Bivalvia (Mollusca). Annual Review of Earth and Planetary Sciences 3, 361-385.

Steeman, M., 2010. The extinct baleen whale fauna from the Miocene-Pliocene of Belgium and the diagnostic cetacean ear bones. *Journal of Systematic Palaeontology* 8, 63-80, doi: [10.1080/14772011003594961](https://doi.org/10.1080/14772011003594961).

Steeman, M., Hebsgaard, M.B., Fordyce, R.E., Ho, S.Y.W., Rabosky, D.L., Nielsen, R., Rahbek, C., Glenner, H., Sørensen, M.V., Willerslev, E., 2009. Radiation of extant cetaceans driven by restructuring of the oceans. *Systematic Biology* 58, 573-585.

Stinnesbeck, W., Frey, E., Rivas, L., Pardo Pérez, J., Leppe Cartes, M., Salazar Soto, C., Zambrano Lobos, P., 2014. A Lower Cretaceous ichthyosaur graveyard in deep marine slope channel deposits at Torres del Paine National Park, southern Chile. *Geological Society of America Bulletin* doi: [10.1130/B30964.1](https://doi.org/10.1130/B30964.1)

Taboada, S., Riesgo, A., Bas, M., Arnedo, M.A., Cristobo, J., Rouse, G.W., Avila, C., 2015. Bone-eating worms spread: insights into shallow-water *Osedax* (Annelida, Siboglinidae) from Antarctic, Subantarctic, and Mediterranean Waters. *PLoS One* 10(11): e0140341.

Tinelli, C., Ribolini, A., Bianucci, G., Bini, M., Landini, W., 2012. Ground penetrating radar and palaeontology: the detection of sirenian fossil bones under a sunflower field in Tuscany (Italy). *Comptes Rendus Palevol* 11, 445-454.

Tinelli, C. (2013). *Marine vertebrates from Pliocene shell beds from Tuscany (Italy): prospecting, taphonomy, palaeoecology and systematic palaeontology*. Pisa: PhD Thesis, 164 pp.

Tomašových, A., Dominici, S., Zuschin, M., Merle, D., 2014. Onshore-offshore gradient in metacommunity turnover emerges only over macroevolutionary time-scales. *Proceedings of the Royal Society B* 281, 1-10.

Tresguerres, M., Katz, S., & Rouse, G.W., 2013. How to get into bones: proton pump and carbonic anhydrase in *Osedax* boneworms. *Proceedings of the Royal Society B* 280: 20130625.

Ugolini, R., 1899. Lo *Steno bellardii* Portis del Pliocene di Orciano Pisano. *Atti della Società Toscana di Scienze Naturali, Memorie* 17, 132-143.

Ugolini, R., 1902. Il *Monachus albiventer* Bodd. del Pliocene di Orciano. *Palaeontographia Italica* 8, 1–20.

Ugolini, R., 1907. Resti di vertebrati marini del Pliocene di Orciano. In: *Atti del Congresso dei Naturalisti Italiani promosso dalla Società Italiana di Scienze Naturali 1906. Tipografia degli Operai, Milano, 195-207.*

Uhen, M.D., 2008. New protocetid whales from Alabama and Mississippi, and a new cetacean clade, Pelagiceti. *Journal of Vertebrate Paleontology*, 28, 589-593.

Volkenandt, M., O'Connor, I., Guarini, J.-M., Berrow, S., O'Donnell, C., 2015. Fine-scale spatial association between baleen whales and forage fish in the Celtic Sea. *Canadian Journal of Fisheries and Aquatic Sciences* 73, 197-204, doi: 10.1139/cjfas-2015-0073.

Young, M.T., Brusatte, S.L., Beatty, B.L., Brandalise De Andrade, M., Desojo, J.B., 2012. Tooth-on-tooth interlocking occlusion suggests macrophagy in the Mesozoic marine *Crocodylomorph Dakosaurus*. The Anatomical Record 295, 1147-1158.

Williams, R., Okey, T.A., Wallace, S., Gallucci, V.F., 2010. Shark aggregation in coastal waters of British Columbia. Marine Ecology Progress Series 414, 249-256.

Woodley, T.H., Gaskin, D.E., (1996). Environmental characteristics of North Atlantic right and fin whale habitat in the lower Bay of Fundy. Canada. Canadian Journal of Zoology 74, 75-84.

Figure and table captions:

Fig. 1 - Location of sedimentary logs within the largest Pliocene basins of Tuscany. Fine Basin (FB): Pagliana (1), Pieve Vecchia (2) and Orciano Pisano (3). Volterra-Era Basin (VEB): Parlascio (4), Lajatico (5), Fabbrica (6) and Volterra (7). Era Basin (EB): San Lorenzo (8), La Serra (9), Poggio al lupo (10), San Maiano (11), Canneto (12), Casenuove (13), Castelfiorentino (14), Fiano (15) and San Gimignano (16). Orcia-Ombrore Basin: Arcille (17) and Poggio alle Mura (18). Siena-Radicofani Basin (SRB): Siena (19), Monteaperto (20), Castelnuovo Berardenga (21), and Radicofani (22) and Fastelli (23). Chiana Basin (CB): Sinalunga (24), Cetona (25) and Allerona (26).

Fig. 2 - Pliocene stratigraphy of Tuscany, see Fig. 1 for the location of numbered sections. All logs measured and described by the authors, except Volterra (log 7: Bianucci et al. 1998), Arcille (log 17: Tinelli et al., 2012; Tinelli, 2013), Siena (log 19: Bianucci et al., 2001), Monteaperto (log 20: Martini et al., 2011), Castelnuovo Berardenga (log 21: Martini et al.,

2016) and Radicofani (log 22: Ghinassi and Lazzarotto, 2005)(chronostratigraphy based on biostratigraphic data from several sources, see main text). See Fig. 1 for the location of numbered logs. (Chronostratigraphy of localities 25 (Poltriciano, Cetona: Capellini, 1883) and 26 (Allerona: Danise, 2010) are unknown logs 21-23 not shown). Correlation between sections is also shown as boundaries of planktonic foraminifera biozones (dashed lines), following the scheme of Sprovieri (1992), based on available biostratigraphic studies for each basin (see main text for relevant references).

Fig. 3 - NumberAbundance of fossil ~~large marine vertebrate~~MM records in major museums of Tuscany-catalogues, distributed by locality of provenance. Each record ranges from a single fragment or ~~single~~ tooth, to ~~a quasi-~~articulated, nearly complete skeleton. A: ~~whales and dolphins~~masticates and odontocetes; B: large sharks; C: ~~dugongs~~sirenians — ~~manscuba~~ diver for scale in each figure. Symbols for basins as in Fig. 1.

Fig. 4 - Detailed sedimentary logs measured at three localities, representing three different stratigraphic contexts for the large marine vertebrate fossil record of the Tuscan Pliocene. The succession at Orciano Pisano is included in synthems S3-S4 of FB, at Arcille-Poggio alle Mura-Camigliano in synthem S2 of OOB, at Sinalunga in synthem S6 of CB. See Figs.1-2 for the location of the numbered localities and references in the main text for facies analysis and sequence stratigraphy of synthem S5. Arcille log from Tinelli et al., 2012, and Tinelli, 2013.

Fig. 5 - Taphonomy of large marine vertebrates at Poggio alle Mura (Figs. 5A-5D) and Arcille (Fig. 5E, see Fig. 4 for the sequence stratigraphic and sedimentary context; plan view of the sirenian skeleton is modified from Tinelli et al., 2012), synthem S2. A: Plan view of the Poggio alle Mura undetermined balaenopterid. Parts of the skeleton are quasi-articulated,

5401 others are scattered, but not far from the original position. B: Quasi-articulated vertebrae of
5402 the same specimen, lying on top of a *Haustator* shell bed. C: Side view of a vertebra on top
5403 of the densely-packed shell bed. D. Detail of the shell bed, in top view. The turritelline
5404 gastropod *Haustator vermicularis* is visible in the upper left, a large fragment of wood in the
5405 lower right, with the ichnofossil *Teredolites* produced by wood-dwelling teredinid bivalves,
5406 in the centre of the photograph. E: Plan view of one of the Arcille dugongsirenian specimens,
5407 ~~belonging to the species of~~ *Metaxitherium appenninicum*. Same scale as in 5A, the arrows
5408 points to the North.

5419
5420
5421 Fig. 6 - Taphonomy of a 10m-long, undetermined balaenopterid at Orciano Pisano (see Fig.
5422 4 for the sequence stratigraphic and sedimentary context), synthem 4. A: Planimetry of the
5423 quasi-articulated and nearly complete skeleton. B: Detail of the central part of the skeleton
5424 in the field. The cortex layer of vertebrae and flipper bones is badly consumed, whereas
5425 some of the costae are still pristine. C: Lateral view of a turritellid shell bed, below, and the
5426 surface where the whale skeleton lied (dashed line), about 15 cm above the shell bed. The
5427 sediment is a very fine-grained silty sand, completely bioturbated (large vertical burrows are
5428 visible). D: Top view detail of the turritellid shell bed. At the center a valve of *Yoldia nitida*,
5429 surrounded by a few specimens of the turritellid *Archimediella spirata*.

5430
5431
5432 Fig. 7 - Taphonomy of an incomplete, undetermined mystecete at Castel San Gimignano,
5433 synthem 4, comprising articulated torso elements. A: Planimetry of the articulated elements.
5434 B: Detail of one of the limbs in the field (trowel for scale = 22 cm): humerus, radius and ulna
5435 are in anatomical relationship; the cortex layer is well preserved, suggesting quick burial of
5436 the carcass. C: Bones of the chest region; on the background the massive sandstone
5437 associated with the fossil whale. Articulated shoreface pectinid bivalves (*Pecten*
5438 *flabelliformis*) were interspersed in the sandstone (photographs by Fabio Cozzini, 1985).

Fig. **87** - Taphonomy of a 5m-long, undetermined balaenopterid at Sinalunga (see Fig. 4 for a tentative sequence stratigraphic interpretation), synthem 6. A: Oblique view of the fully articulated skeleton lying in a gravelly sandstone, stratified in the lower part, massive in the upper. Vertebrae are tightly connected as if in life. B: Plan view of the flipper, with carpals, metacarpals and phalanges in perfect anatomical connection. C: The gravelly sandstone lies above a bioturbated muddy sandstone, with vertical burrows (*Ophiomorpha*). D: The lower part of the unit with the whale skeleton is formed by three fining-upward beds. Each bed grades from gravel to medium-grained sand. Articulated and empty shoreface bivalves (e.g., *Callista chione*), not in life position, are interspersed with the coarse gravel.

Fig. **98** - Quantitative analysis of the facies type and sequence stratigraphic distribution of large marine vertebrates, Pliocene of Tuscany (N = 39, see Tab. 2). A: The vast majority of cases (77%) are associated with fine-grained muddy sediments of the shelf, a few are found in deltaic coarse-grained strata. B: Most **LMVsMM** (69%) **are** found in highstand deposits, a few in proximity of the maximum flooding interval, or in transgressive deposits.

Fig. **109** - NMDS ordination of bulk samples (N = 72, see Fig. 2 for their stratigraphic position), based on the distribution of standardised abundances of 329 mollusc species (further explanation on multivariate techniques in the main text). A: Samples are subdivided based on the associated sedimentary facies types (F2-F5 in Tab. 1). The main axis ordines samples along a paleodepth gradient, from shallowest to deepest, moving from the left to the right side of the bivariate plot. B: Same ordination, with an estimate of absolute paleodepth of each sample based on score along the main axis and calibrated through the average modern depth distribution of 23 extant species characterising the Pliocene dataset (abundance > 0.15%, see text; regression logarithmic curve in the inset). Vertebrates

recovered in proximity of some of the samples (N = 13) are plotted on the diagram, confirming that, on average, the LMVMM fossil record is concentrated on the open shelf at an estimated depth of 30-300 m (M = dugongs_sirenians; S = sharks; P = pinnipeds; W = whales).

Fig. 11 - Occupancy of trophic levels by Pliocene marine mammals and sharks in the north-western Mediterranean, expressed by number of species per trophic level (see Tabs 3, 5 for explanation and references). This figure, summing up data for the whole epoch, spanning circa 2,8 My, closely matches the association found in one single synthem S4, of much shorter duration (mid-Piacenzian, 3-4 hundred thousand years).

Tab. 1 - Sedimentary facies types.

Tab. 2 - Stratigraphic, taphonomic and paleoenvironmental framework for Pliocene large marine vertebrates marine mammals recovered of in Tuscany, with abundance data (N = 3964). A: Highly-aArticulated and quasi-articulated skeleton; C: Highly-eComplete and quasi-complete skeleton; C+PC: Cranial and post-cranial bonesremains.

Tab. 3 - Paleoecology of Pliocene large mammals compared toand ecology of modern NWMS analogues.

Tab. 4 - Geographic distribution of Pliocene large mammals in Tuscany.

Tab. 5 . Paleoecology of Pliocene sharks compared toand ecology of modern NWMS analogues.

Tab. 6 - Geographic distribution of Pliocene sharks in Tuscany.

Supplement Material

Fig. S1 - Dendrogram resulting from cluster analysis of a dataset of 336 species distributed in 72 samples (standardised abundance, square-root transformed, Bray-Curtis similarity).

Tab. S1 - Ranked total average of standardised abundance of Pliocene molluscs.

Tab. S2 - Score on main axis of NMDS ordination (NMDS1) of Pliocene mollusc species (N=329).

Tab. S3 - Average depth of extant mollusc species with >1,5% abundance in the Pliocene dataset (N=23).

Tab. S4 - Estimated depth of Pliocene samples (N=72) based on NMDS1.

Pliocene stratigraphic paleobiology in Tuscany and the fossil record of marine megafauna

Abstract

Tuscany has a rich Pliocene record of marine megafauna (MM), including mysticetes, odontocetes, sirenians and seals among the mammals, and six orders of sharks among the elasmobranchs. This is reviewed with respect to paleogeography and sequence-stratigraphic occurrence in six different basins. Conditions at the ancient seafloor are explored by means of sedimentary facies analysis, taphonomy and multivariate techniques applied to a large quantitative dataset of benthic molluscs. MM is rare or absent in most basins during the Zanclean, except in one basin, and most abundant in Piacenzian deposits in all six basins. MM occurs preferentially in fine-grained shelfal highstand deposits of small-scale depositional sequences, or at condensed horizons of the maximum flooding interval. It is rare in shallow marine paleoenvironments and nearly absent in bathyal paleosettings. Paleogeographic and paleoecological evidence and a comparison with modern patterns of marine upwelling suggest that a wedge of nutrient-rich waters sustained in the offshore during the Pliocene a high biomass of primary producers and a community of apex consumers and mesopredators, similarly to the modern Northwestern Mediterranean Sea, with a higher species-richness than the modern. The highest MM diversity coincides with the mid-Piacenzian warm period, suggesting that facies control does not obscure a link between climate and diversity. We underline however that not all marine environments were suitable for marine mammal preservation. Buoyant carcasses were preferentially dismembered and destroyed in high-energy shallow waters, with the possible exception of delta front deposits, where sudden sediment input occasionally buried pristine carcasses. We hypothesise that carcasses sunken on the seafloor below the shelf break underwent destruction through the activity of a whale-fall biota of modern type, specialised in the

consumption of decomposing tissues, both soft and mineralised. A taphonomic window was left between storm wave base and the shelf break. Here water pressure is high enough to prevent the formation of decomposing gases and the resurfacing of carcasses, while the lack of a specialised whale-fall biota slows down bone degradation with respect to deeper settings. Sedimentation rate was high enough to cover skeletal material before its complete destruction. An estimate of paleobathymetries based on multivariate techniques suggests that the preferential depth for the inclusion of MM in the fossil record was 30-300 m. The results are compared with major Mesozoic and Cenozoic MM records worldwide. Available evidence suggests that the late Neogene radiation of large whales, true ecosystem engineers, and their size increase, triggered the radiation of a bone-eating fauna that hampered, and hampers, MM preservation in the deep sea. Stratigraphic paleobiology and an ecosystem-level approach deliver useful insights in the nature of the fossil record.

Pliocene stratigraphic paleobiology in Tuscany and the fossil record of marine megafauna

Stefano Dominici (Università di Firenze, Museo di Storia Naturale, Firenze, Italy)

Silvia Danise (Plymouth University, School of Geography, Earth and Environmental Sciences, Plymouth, United Kingdom)

Marco Benvenuti (Università di Firenze, Dipartimento di Scienze della Terra, Firenze, Italy)

Abstract

Tuscany has a rich Pliocene record of marine megafauna (MM), including mysticetes, odontocetes, sirenians and seals among the mammals, and six orders of sharks among the elasmobranchs. This is reviewed with respect to paleogeography and sequence-stratigraphic occurrence in six different basins. Conditions at the ancient seafloor are explored by means of sedimentary facies analysis, taphonomy and multivariate techniques applied to a large quantitative dataset of benthic molluscs. MM is rare or absent in most basins during the Zanclean, except in one basin, and most abundant in Piacenzian deposits in all six basins. MM occurs preferentially in fine-grained shelfal highstand deposits of small-scale depositional sequences, or at condensed horizons of the maximum flooding interval. It is rare in shallow marine paleoenvironments and nearly absent in bathyal paleosettings. Paleogeographic and paleoecological evidence and a comparison with modern patterns of marine upwelling suggest that a wedge of nutrient-rich waters sustained in the offshore during the Pliocene a high biomass of primary producers and a community of apex consumers and mesopredators, similarly to the modern Northwestern Mediterranean Sea, with a higher species-richness than the modern. The highest MM diversity coincides with the mid-Piacenzian warm period, suggesting that facies control does not obscure a link

between climate and diversity. We underline however that not all marine environments were suitable for marine mammal preservation. Buoyant carcasses were preferentially dismembered and destroyed in high-energy shallow waters, with the possible exception of delta front deposits, where sudden sediment input occasionally buried pristine carcasses. We hypothesise that carcasses sunken on the seafloor below the shelf break underwent destruction through the activity of a whale-fall biota of modern type, specialised in the consumption of decomposing tissues, both soft and mineralised. A taphonomic window was left between storm wave base and the shelf break. Here water pressure is high enough to prevent the formation of decomposing gases and the resurfacing of carcasses, while the lack of a specialised whale-fall biota slows down bone degradation with respect to deeper settings. Sedimentation rate was high enough to cover skeletal material before its complete destruction. An estimate of paleobathymetries based on multivariate techniques suggests that the preferential depth for the inclusion of MM in the fossil record was 30-300 m. The results are compared with major Mesozoic and Cenozoic MM records worldwide. Available evidence suggests that the late Neogene radiation of large whales, true ecosystem engineers, and their size increase, triggered the radiation of a bone-eating fauna that hampered, and hampers, MM preservation in the deep sea. Stratigraphic paleobiology and an ecosystem-level approach deliver useful insights in the nature of the fossil record.

1. Introduction

The modern marine megafauna (MM) includes all marine mammals, seabirds, sea turtles and sharks, apex consumers that influence their associated ecosystems (Lewison et al., 2004), both pelagic and nearshore, through top-down forcing and trophic cascades, and now severely affected by human impact (Estes et al., 1998, 2011, 2016). On a macroevolutionary scale, predation pressure has shaped the evolution of marine preys, with

121
122 feedbacks on predators, setting the stage for the Mesozoic marine revolution (Vermeji, 1977;
123
124 Chen and Benton, 2012; Benton et al., 2013). The new ecosystem structure started in the
125
126 the Early and Middle Triassic with several lineages of Actinopterygian fishes (Chen and
127
128 Benton, 2012), continuing with marine reptiles possessing feeding styles (Fröbisch et al.,
129
130 2013; Motani et al., 2015; but see also Motani et al., 2013) and reproductive adaptations
131
132 (Motani et al., 2014) of modern type. Triassic and Jurassic novelties underwent a prolonged
133
134 crisis during the Cretaceous, with the gradual extinction of plesiosaurians, mosasaurs
135
136 (Benson et al., 2010) and ichthyosaurs (Fischer et al., 2016), and a diversity drop of sharks
137
138 (Guinot et al., 2012). A marine megafauna of comparable size returned in the Paleogene,
139
140 with the new diversification of neoselachian elasmobranchs (Kriwet and Benton, 2004) and
141
142 the evolution of large marine mammals: Eocene archaeocetes (Uhen, 2008, Gingerich et
143
144 al., 2009) and Oligocene odontocetes and mysticetes (Gingerich, 2005; Marx and Uhen
145
146 2010, Berta, 2012; Marx et al., 2016) empowered by high metabolic rates and new anatomic
147
148 features (Armfield et al., 2013). Among the largest vertebrates of all times, after a dramatic
149
150 size increase at the outset of glacial age (Marx et al., 2016; Bisconti et al., 2017; Slater et
151
152 al., 2017), baleen and sperm whales are among today's ocean's ecosystem engineers
153
154 (Roman et al., 2014) with which to compare their Mesozoic analogues (Smith et al., 2016).
155
156
157 Notwithstanding a crucial role in ecology and evolution, the nature and distribution of the
158
159 MM fossil record has been less explored, compared to that of marine invertebrates and
160
161 terrestrial vertebrates. Available data suggest a strong correlation between taxic diversity
162
163 and the number of marine fossiliferous formations, resulting in megabiases in the fossil
164
165 record (e.g., Cretaceous: Benson et al., 2010). Within its vast history, studies on the
166
167 geologically recent marine megafauna offer important insights, considering our better
168
169 knowledge of: 1) geological setting, in terms of outcrop extent and high-resolution
170
171 stratigraphy; 2) ecologic role played by individual species, whether extant or extinct, in terms
172
173 of habitat, trophic role, life histories and population structure, thanks to a comparison with
174
175
176
177
178
179
180

181
182 extant descendants, or close relatives; 3) MM taphonomy, based on actuopaleontology. A
183
184 recent global study revealed that MM extinction peaked in the late Pliocene, between 3.8
185
186 and 2.4Ma, linked to the sudden drop in the extension of nearshore environments after a
187
188 large sea level regression (Pimienta et al., 2017), confirming that the fossil record offers
189
190 important clues on the vulnerability of keystone marine species to climate change. We
191
192 contribute here to a better understanding of the Pliocene fossil record by reviewing the rich
193
194 MM of Tuscany, in Italy. In particular, we consider all reports of Pliocene marine mammals
195
196 and sharks and revise taphonomy and sedimentary facies associated with all known
197
198 findings, setting them within a sequence stratigraphic framework. We also reconstruct the
199
200 paleoenvironmental context and review data on a part of the marine ecosystem through the
201
202 paleoecology of fossil invertebrates on a regional basis, following a stratigraphic
203
204 paleobiological approach that can be applied to both the recent and the distant geological
205
206 past (Patzkowsky and Holland, 2012). Published studies that have taken this direction are
207
208 still a few, examples concerning Jurassic ichthyosaurs, plesiosaurs and pliosaurs (McMullen
209
210 et al., 2014), Cretaceous turtles, plesiosaurs, bony fish and sharks (Schemisser McKean
211
212 and Gillette, 2015), Eocene archaeocetes, sea cows and sharks (Peters et al. 2009), and
213
214 Neogene marine mammals and sharks (Boessenecker et al., 2014). All of these papers
215
216 record the co-occurrence of shelly faunas, only one undertaking quantitative studies of the
217
218 distribution of fossil invertebrates (Jurassic of the Sundance Formation: McMullen et al.,
219
220 2014, see also Danise and Holland, 2017). The benefits of an outcrop-scale sequence
221
222 stratigraphic approach include: (1) an independent record of relative sea-level change to
223
224 test paleobiological hypotheses (see also Pyenson and Lindberg, 2011; Noakes et al.,
225
226 2013); (2) a chronostratigraphic scheme for high-resolution correlations; (3) a means to
227
228 recognise minor and major breaks of the record; (4) an ecological and sedimentary
229
230 framework for taphofacies distribution (Patzkowsky and Holland, 2012); and (5) an
231
232
233
234
235
236
237
238
239
240

independent control of onshore-offshore patterns of fossil assemblages (e.g., Tomašových et al., 2014).

Researchers that study the geologic history of marine ecosystems have focused on patterns of ecological restructuring based on the taxonomy of selected groups (e.g., Thorne et al., 2011; Benton et al., 2013; Scheyer et al., 2014; Fischer et al., 2016), at the expenses of a more holistic approach that includes functional diversity and embraces as many ecosystem components as possible (Dineen et al., 2014). By analogy with ecologists who shift focus from models based on single groups (e.g., Steeman et al., 2009) to an all-embracing vision of marine life (Lawton, 1994; Sergio et al., 2014), connecting food web ecology with landscape ecology (Polis et al., 1997; Estes et al., 2011), stratigraphic paleobiology can draw from the fossil record and offer multidimensional insights on the complex geological history of modern marine ecosystems.

After revising fossil MM hosted in major museums of Tuscany, both isolated and articulated remains, we focus on all fossil bones that can be stratigraphically (e.g., Bianucci et al., 1995, 1998, 2001; Tinelli, 2013) and taphonomically framed (e.g., Dominici et al., 2009; Bianucci, 2010; Danise and Dominici, 2014). MM lists for the Mediterranean Pliocene have been recently updated (marine mammals: Landini et al., 2005; Bianucci et al., 2009a; Sorbi et al., 2012; Bianucci and Vomero, 2014; sharks: Marsili, 2006). Species-level ecological data are available on modern apex consumers and mesocarnivores (Pauly et al., 1998; Cortés, 1999), with detailed information made available for Mediterranean species following conservation concerns (marine mammals: Notarbartolo di Sciara et al., 2016; sharks: Cavanagh and Gibson, 2007), allowing for a detailed paleoecological evaluation of the Tuscan fossil record. The actualistic approach is also viable for species of benthic molluscs, about half of which are still extant in modern Mediterranean bottoms (55% of extant species of Mediterranean and North Sea bivalves, excluding strictly brackish and bathyal forms, i.e., 202 out of 367 species, survives from the Zanclean: Raffi et al., 1985). The regional

quantitative study of molluscan assemblages was the basis for an independent assessment of paleoenvironments, paleoecology and paleobathymetry. A revision of abundance distributions of marine molluscs, the largest contributors to Mediterranean Pliocene shell beds and a key component of Mesozoic and Cenozoic marine ecosystems (Stanley, 1975; Vermeij, 1977), allowed to further explore the structure and composition of Pliocene marine ecosystems, and reconstruct a paleobathymetric gradient (e.g., Scarponi and Kowaleski, 2004) along which to frame MM occurrence. The present work must necessarily start with a review of the chronostratigraphy and physical stratigraphy of the Tuscan marine Pliocene.

1. Geological setting

The Pliocene succession of Tuscany was deposited in a complex setting characterised by continental collision related to the later evolution of the Northern Apennines chain. According to a well-established hypothesis, the region, affected by shortening before the Middle-Late Miocene, accommodated by NE-verging thrust and fold systems, underwent crustal extension during the late Neogene and the Quaternary (DeCelles, 2012; Fig. 1). Crustal extension generated differential subsidence in a series of normal-fault controlled hinterland sedimentary basins, filled throughout by continental and shallow marine, mostly clastic successions (Martini and Sagri, 1993; Pascucci et al., 2006; Brogi, 2011). An alternative hypothesis places the late orogenic hinterland basins in a more complex tectonic setting characterised by the alternation of compressive, extensional and transcurrent stress fields (Benvenuti et al., 2014; Bonini et al., 2014).

The Neogene Tuscan basins considered in this work include, from West to East, and from North to South, the Fine Basin (FB; Bossio et al., 1997), the Volterra-Era Basin (VEB; Bossio et al., 1994), The Elsa Basin (EB; Benvenuti et al., 2014), the Ombrone-Orcia Basin (OOB; Bossio et al., 1991; Nalin et al., 2010), the Siena-Radicofani Basin (SRB; Ghinassi and

Lazzarotto, 2005; Martini et al., 2011, 2016), and the Chiana Basin (CB; Fig. 1: Pesa Basin not considered here). With one exception (OOB, see below), these basins show a shape conditioned by the structural and physiographic features of the inner portion of the Northern Apennines. Their NW-SE general elongation reflects the trend of the thrust-related anticline ridges developed during earlier collisional stages. These compressive structures have bounded most basins through their infilling, only to be obliterated by younger parallel normal fault systems, leaving an invariant stratigraphic onlap of the Pliocene successions onto the basin margins. Despite a NW-SE distribution of the hinterland basins, the structural setting is responsible for a NE-trending physiographic and paleogeographic gradient, where the FB is closest, and the CB furthest, from offshore settings throughout the late orogenic phase, with important implications for the facies architecture and the distribution of marine vertebrates and shell beds. Differential active uplift of the basin shoulders during the Pliocene, coupled with important erosional phases, resulted in a different preservation of the original stratigraphic architecture. The infill during the Zanclean is generally characterised by relatively continuous open marine successions, the correlative fluvial-coastal systems missing due to uplift and erosion of basin margins. On the other hand, the Piacenzian infill is characterised by genetically-connected, fluvial, coastal and shallow marine facies tracts, particularly well-preserved in the EB, hinting to reduced uplift of the marginal areas. The modern physiography of OOB, escaping the structurally-controlled geometry of the other basins, mimics instead an original fluvial network developed during the latest Messinian, flooded after the Salinity Crisis (Bossio et al., 1991; Benvenuti et al., 2015), filled during the Pliocene, and finally disrupted by post-Pliocene uplift and erosion. This difference in the structural history also justifies the preservation in OOB of Zanclean fluvial and shallow marine facies (Fig. 2).

1.1 Pliocene Stratigraphy

The Neogene succession of Tuscany is up to 2000 m-thick, about half of which belongs to the uppermost Miocene-Pleistocene interval (Bossio et al., 2004; 1997; Benvenuti et al., 2014). The Pliocene has been traditionally subdivided into three main informal lithostratigraphic units: continental conglomerates and sandstones at the base, overlain by the “Blue Clay Formation” (*Argille Azzurre*: Zanclean-lower Piacenzian), marking the post-Messinian Mediterranean marine transgression and forming the thickest part of the basin infill (e.g., Bossio et al., 1994; 1997; Ghinassi and Lazzarotto, 2005), and the “Upper Sands” (*Sabbie superiori*: Piacenzian-Gelasian) and conglomerates, deposited during the ensuing regression. Several finer lithostratigraphic units have been introduced to define the local stratigraphy, resulting in a complex and largely informal lithostratigraphic terminology which includes Zanclean lower “Blue Clays” and Zanclean-Piacenzian upper “Blue Clays” (Capezzuoli et al., 2015), the latter eventually further separated by the widespread occurrence of Piacenzian carbonates (Nalin et al., 2016). The lower Zanclean (OOB: Ghinassi, 2007; Nalin et al., 2010; Dominici et al., 2012) and the Piacenzian, are characterised by the high-frequency alternation of coarse-grained and fine-grained facies, ranging from fluvial to marine shelf settings (Benvenuti et al., 1995; 2007; 2014; Martini et al. 2011, Fig. 2). The dynamics of the Pliocene infilling are better-understood in the EB, where six synthems have been defined, each up to more than 200 m-thick, further subdivided in a number of elementary and composite depositional sequences and chronologically calibrated through marine biostratigraphy and continental vertebrate biochronology (Benvenuti and Del Conte, 2013; Benvenuti et al., 2014, with references).

2. Materials and methods

Stratigraphic sections were measured and described at several localities (Fig. 1). Siliciclastic and carbonate facies were described, subdivided into groups of facies based on lithology, sedimentary structures and ichnology, and interpreted in terms of process and depositional environment (Tab. 1). Each group represents a set of individual facies forming monogenic associations (in the sense of Mutti et al., 1994), i.e., the meter-scale stacking of facies which express the autocyclic behaviour of specific depositional systems within a given accommodation space (Benvenuti and Del Conte, 2013). Sequence stratigraphic concepts have been applied to reconstruct the dynamics of basin infills at a hierarchy of scales, advancing hypotheses on controlling factors. The chronostratigraphic subdivision of Benvenuti et al. (2014), which divides the Pliocene into six synthem, S1-S6 from older to younger, was extended to all six Tuscan basins by referring to available biostratigraphic schemes (Fig. 2). The sequence stratigraphic interpretation of S2 in OOB is based on Tinelli (2013). Other parts of the S1-S3 succession were drawn based on available lithostratigraphic literature (see below). Studies integrating sedimentary facies analysis, biostratigraphy and magnetostratigraphy (Nalin et al., 2016) have helped correlating carbonate bodies typical of S4. The reader is referred to Benvenuti et al. (2007; 2014) for details on facies analysis and sequence stratigraphic interpretation of synthem S3-S6.

First, data on the geographic distribution of fossil marine mammals, large sharks and sirenians were largely based on collections housed at the Natural History museums of the University of Florence (UFMSN), University of Pisa (UPMSN), and Accademia de' Fisiocritici of Siena (AFMSN), the three largest collections of Tuscany, and at the Geological Museum Giovanni Capellini, Bologna University (MGGC). In particular, counts of cetaceans were based on UFMSN collections (Mysticeti and Odontoceti, N = 142), large shark on the sum of UFMSN, UPMSN and AFMSN collections (Carcharhiniformes and Lamniformes, N = 337, data synthesised from Marsili, 2006), sirenians from all reports in Tab. 2 (N = 10). Each record is formed either by a single element (e.g., whale bone, shark tooth), by a few

elements of the same individual, or by a whole, quasi-articulated skeletons. A large proportion of this dataset lacks precise location, allowing only for some crude stratigraphic attribution (Fig. 3).

On a second step, all fossil Tuscan Pliocene cetaceans, sirenians and pinnipeds that could be framed within the available high-resolution stratigraphic framework and associated with taphonomic data, were selected. At this step, after excluding unidentified MM remains, a dataset of 64 specimens (cetaceans N = 50; sirenians N = 10; pinnipeds N = 4) was assembled. Association with shark teeth is frequent (55% of 25 cases according to Danise and Dominici 2014 for the Italian Pliocene; see also Bianucci et al. 2002, 2010). The majority of the 64 specimens are included in the catalogue of UFMSN, UPMSN, AFMSN and MGGC, whereas a few are stored in smaller collections of the municipalities of Montaione, Scandicci (Florence province), and Certaldo (Pisa province), one in a private property (Castello di Villa Banfi, near Montalcino, Siena province), and one in the Museum National d'Histoire Naturelle in Paris (France). Whenever possible, large marine vertebrates were coded by synthem (N = 60) and depositional environment (N = 54). We analysed abundance distributions among marine mammals, and species richness of marine mammals and sharks. To infer Pliocene paleoecology, fossil taxa recognised in Tuscany were compared with their closest descendants, focusing on the species today living in the North-Western Mediterranean Sea (NWMS).

All known MM-bearing sedimentary facies are associated with a mollusc-dominated benthic fauna. The third step of the analysis concerned a quantitative study of shell beds, allowing: 1) to interpret the regional evolution from a perspective independent from the sedimentary facies, 2) to characterise MM paleoenvironmental and bathymetric distribution, and 3) to explore the structure of the benthic component of marine ecosystems and to identify underlying environmental controls. 72 Bulk samples were collected at major shell beds at bed resolution throughout the succession and sieved with 1 mm mesh size. Fossils of

bivalves, gastropods and scaphopods were identified to species level. The minimum number of individuals was calculated following standard approaches (see Patzkowsky and Holland, 2012), resulting in a richness of 525 species (S) and a total abundance of 64206 individuals (N). We coded each fossil assemblage by synthem, tract of small-scale depositional sequence, and depositional environment. Most samples belonged to facies types F2-F5 (Tab. 1; see also Tomašových et al., 2014). Facies F6 usually lacks macrofossils and allowed for the collection of only one sample. No samples were collected in facies F1, lacking marine shells, facies F7, mostly devoid shells, and facies F8, which is richly fossiliferous, but lacks aragonite shells and is associated with specimens hardly extractable from the rock. The resulting quantitative dataset served for statistical analyses on the distribution of species-level abundances on a siliciclastic shelf depositional system, performed with the software Primer 6.0 (Clarke and Gorley, 2006). Analyses included clustering and nMDS ordination techniques on a Bray-Curtis similarity matrix, of standardised, square-root transformed data (72 samples; S = 333 and N = 63518 after the exclusion of singletons). To test statistically whether there is a significant difference between two or more groups of sampling units based on sedimentary facies, we performed an analysis of similarity (ANOSIM). To interpret the outcomes of the quantitative study and the significance of clusters we used species-level autoecologic information available for the most abundant species, based on the distribution of extant forms. This information, retrieved from the Marine Biodiversity and Ecosystem Functioning EU website (MARBEF: www.marbef.org), included the average life depth of 23 modern species that in our dataset had an overall abundance > 0.15%.

3. Results

3.1. Unconformity-bounded units

3.1.1 Synthem S1: the early Zanclean transgression

The Miocene-Pliocene transition, marking the return to marine conditions after the Messinian salinity crisis (Krijgsman et al., 1999), is recorded in limited exposures of earliest Zanclean, open marine mudstones resting both unconformably or conformably onto latest Messinian non-marine deposits (*Lago-Mare*), an isochronous boundary being dated in the Mediterranean at 5.33 Ma (Roveri et al., 2014). Differences depend on the specific structurally-controlled distribution of hinterland basins, where an uplifting chain determined the presence of thresholds delaying the early Zanclean marine flooding from inner (EB, OOB: Benvenuti et al., 2015a) to outer hinterland basins (FB, VEB). In inner basins such as EB, where S1 has been defined, continental deposition continued into the earliest Zanclean, marine flooding occurring within the MPL1 biozone (references in Benvenuti et al., 2015a). A chronostratigraphical equivalent of S1 is represented in FB and VEB (Bossio et al., 1978) by an open-shelf mudstone, conformably resting on latest Messinian *Lago-Mare* deposits (Roveri et al., 2014). An apparently analogous situation is documented in OOB, where MPL1 shelfal mudstones of the early Zanclean (normal chron C3n) rest on a Messinian to basal Pliocene paleovalley fill (Benvenuti et al., 2015a).

3.1.2 Synthem S2: Zanclean differential preservation

Synthem S2, as recognised in the EB (Benvenuti et al. 2014), is represented by relatively thin fluvial conglomerates unconformably resting on S1 (biozones MPL1-MPL2), capped by S3 (biozone MPL4a: Bossio et al., 1993; 2001), comprising important stratigraphic gaps at its base and top. On the other hand, in FB (Bossio et al., 1997), VEB (Bossio et al., 1994)

and SRB (Ghinassi and Lazzarotto, 2005), the same chronostratigraphic interval is recorded by monotonous epibathyal mudstones several hundred meters thick, locally intercalated with delta-front hyperpycnal sandstones and conglomerates. In a very broad sense, the concept of S2 is extended to these basins, by assuming that erosional unconformities in EB pass to into correlative conformities in rapidly subsiding adjacent basins, where thick successions could be accommodated. Apart from exceptions, no shells were found in bathyal mudstone or in deltaic sandstone. In OOB, the same time span is marked by a N-S facies gradient characterised by a single deepening-upward succession, from fluvio-deltaic sandstone to shelfal mudstone, replaced by a succession made of four distinct regressive-transgressive units in the Orcia valley to the north (Ghinassi, 2005; Benvenuti et al., 2015b). A laterally-continuous shell bed, with sharks remains and skeletons and articulated bones of whales, sirenians and large teleost fishes (Danise, 2010; Sorbi et al., 2012; Tinelli, 2013), marks a major transgressive surface overlain by open shelf mudstones (Sorbi et al., 2012; Tinelli, 2013: biozone MPL2) that is hypothetically traced along a NE-SW profile (Figs. 4, 5).

3.1.3 Synthem S3: Zanclean-Piacenzian transition

Synthem S3 is subdivided in EB into a lower and an upper interval (Benvenuti et al., 2014). The lower division is represented by deepening-upward, coarse-grained delta front system, overlain by an upper mudstone division from an open shelf setting. The upper part is rich with shell beds, and occasional articulated whale skeletons, associated with shark teeth, have been recovered (Danise and Dominici, 2014). Similar shelf mudstones of the MPL4 biozone crop out in VEB and SRB (Bossio et al., 1993; Riforgiato et al., 2005) whereas biostratigraphically equivalent mudstones in FB testify to an upper epibathyal paleoenvironment.

3.1.4 Synthem S4: early Piacenzian warm climate and high sea-level

Synthem S4 (Benvenuti et al., 2014) has been recognised in FB and VEB by facies similarities and chronostratigraphic correlation. In EB, S4 comprises a lower interval dominated by richly fossiliferous, massive mudstone or very-fine-grained sandstone (prodelta-inner shelf), overlain by bioclast-rich sandstones recording prograding mixed carbonate-clastic ramp, outcropping in the southeastern part of the Elsa valley. Equivalent deposits, also comprised in biozone MPL4b, are patchily distributed in FB, VEB, SRB, OOB and other basins of southern Tuscany (Ghinassi and Nalin, 2010; Ghinassi, personal communication, 2015). The upper interval of S4 is formed by a succession of delta front sandstones, passing in EB eastern margin to a few tens of m-thick fluvial succession, hinting at an original depositional gradient. S4 is apparently missing due to erosion north of San Gimignano (EB), and around Lajatico (VEB). Biostratigraphic data allow to refer S4 to the upper part of biozone MPL4b and the lower part of MPL5a, thus comprising the mid-lower part of the Piacenzian, globally characterised between 3.264-3.025 Ma by warm climate and relatively high sea level (Raymo et al., 2009; Dowsett et al., 2013; Prista et al., 2015).

3.1.5 Synthem S5: mid-Piacenzian high-frequency sea level variation

Synthem S5, recognised in EB and VEB, is bounded below by an erosional unconformity that cuts deeply into underlying units, bringing S5 directly on top of S3 (EB: log 15; VEB: logs 5-6 in Fig. 1). S5 is up to about 200 m in EB, where it has been subdivided into a hierarchy of small-scale depositional sequences (Benvenuti et al., 2007; Dominici et al., 2008: see following paragraphs). Each composite depositional sequence forms a tens-of-m-thick asymmetric sedimentary cycle, composed by a deepening-upward part, from fluvial or coastal coarse-grained sediments, to open shelf mudstones, sometimes topped by a

regressive shoreface or delta sandstone, other times directly overlain by the next sequence through a sharp contact. Fluvial, brackish-water, and other intertidal deposits mark the lower part of each composite sequence, usually topped by a laterally-continuous shell bed, from a few cm to a few dm-thick, representing a surface of transgression. Shell beds are particularly well-developed around the middle part of sequences, where they separate shoreface and delta sandstones from overlying open shelf mudstones, marking the time of maximum flooding (MFS). Large marine vertebrates, including articulated whale skeletons and large sharks (Danise and Dominici, 2014), are often recovered both at MFS and overlying mudstone (Fig. 1). Towards the north-eastern margin of EB, cyclothemic fluvial conglomerates, sandstones and mudstones replace coastal and fully marine deposits, testifying to an original facies gradient. In the central part of EB, composite sequences are stacked to form a deepening-upward succession, with a topmost thick and laterally-continuous open shelf mudstone interval, directly onlapping the S4-S5 basal unconformity on the eastern EB (log 15, Fiano: Fig. 1).

3.1.6 Synthem S6: Piacenzian-Gelasian climate change and regression

As S5, synthem S6 is also built through a hierarchy of small-scale depositional sequences, better expressed in EB, but also documented in SRB and CB. In EB, fluvial coarse-grained sandstones fill a deep valley incised in S5 deposits (logs 8-10 in Fig. 1), resting on the basal unconformity of S6. Intertidal or coastal lagoon deposits form the transgressive systems tract of the composite depositional sequence. A laterally-continuous shell bed testifies to the MFS of S6, topped by highstand shoreface and delta front sandstones. The Piacenzian-Gelasian boundary, corresponding to a major climatic transition from warmer-moister to colder-drier conditions (Benvenuti et al., 1995b; 2007), is marked by the Gauss-Matuyama reversal detected at Montopoli (Lindsay et al., 1980). Highstand marine sandstones are

characterised by the recovery of two mysticete skeletons, at Montopoli (EB, see Capellini, 1905) and Sinalunga (CB, Fig. 7). Fully continental environments were established throughout the Gelasian in all basins here under study (e.g., Benvenuti and Del Conte, 2013; Benvenuti et al., 2014; Bianchi et al., 2015).

3.2 Sedimentary facies and facies associations

Seven groups of siliciclastic facies and one group of carbonate facies, with very different fossil content, have been recognised (Tab. 1). Siliciclastic facies form a paleoenvironmental gradient from terrestrial to marine and, in the case of marine facies, from shallowest to deepest (Fig. 8a). Facies types are fluvial conglomerate and sandstone, and alluvial mudstone (F1); intertidal to very shallow subtidal mudstone and muddy sandstone (F2); shallow subtidal coarse- and medium-grained sandstone with sparse conglomerate (F3); deep subtidal muddy fine-grained sandstone (F4); open shelf sandy mudstone (F5); outer shelf to upper bathyal mudstone (F6); outer shelf and bathyal turbidite sandstone and conglomerate (F7). An eighth group is formed by facies deposited subtidally in limited mixed carbonatic-siliciclastic ramps (F8), irrespective of depth (Nalin et al., 2016). Facies F6-F7 are restricted to Zanclean deposits (synthems S1-S3); facies F5 and F8 characterise the upper Zanclean-middle Piacenzian interval (synthems S3-S4); facies F1-F4 characterise the upper Piacenzian (synthems S5-S6). The OOB succession, characterised by facies F1-F4, is an exception within the Zanclean.

3.3 Elementary depositional sequences (EDS)

Both fluvial (Benvenuti and Del Conte, 2013) and marine facies groups (siliciclastics: Benvenuti et al., 2007; 2014; carbonates: Nalin et al., 2016) are stacked to form facies

associations which record cyclic variations of depositional and environmental conditions in response to a change in accommodation space. Physical surfaces and the intervening deposits allowed to subdivide depositional sequences in systems tracts (Benvenuti et al., 2007; Dominici et al., 2008). At the simplest scale, these hybrid facies associations form elementary depositional sequences, up to 10-20 m thick, in their turn stacked to form composite sequences (original concepts from Mutti et al., 1994). This hierarchy is particularly evident in synthems S5-S6, formed at a time of pronounced glacio-eustatic oscillations and expressed around coastal settings, where maximum facies contrast allows for the expression of subtle cycle of sea level variation (e.g., Benvenuti and Dominici, 1992; Benvenuti et al., 2007; Dominici et al., 2008). Analogue sharp facies contrast within Zanclean EDS in OBB (Tinelli, 2013), but is otherwise absent in deeper sediments (facies F5-F7). EDSs have different expressions depending on the time interval and the sedimentary basin.

3.3.1 Zanclean EDS (synthems S1-S3)

In most basins, deposition of synthems S1-S2-S3 takes place at outer-shelf or bathyal depths, well below the point on a depositional profile where the rate of relative sea level change is zero (equilibrium point). Here the sediment supply is not sufficient to fill the available accommodation space and an aggradational style of deposition prevails, with the result that in most Zanclean settings smaller cycles of sea level variations are not marked by a facies change. The sharp facies change recorded where the monotonous muddy deposition is interrupted by turbidite sandstone and conglomerate, is connected with synthem boundaries and major tectonic phases of restructuring of the region. MM and shell beds are practically absent. This situation reverses in the Orcia-Ombrone basin, where depths of deposition are shallower and EDS are expressed. At Arcille deltaic sandy conglomerates and sandstones (facies F3) are overlain by a fluvial cross-bedded sandstone

(facies F1a), separated by a transgressive surface (TS) from an overlying bioturbated shallow marine sandstone. A *Haustator vermicularis* shell bed (Danise, 2010; Tinelli, 2013) forms the MFS separating the shoreface sandstone from an open marine mudstone with scattered shells (facies F5), marking a sudden and prolonged deepening of the basin (biozone MPL2: Sorbi et al., 2012; Tinelli, 2013). The succession is topped by deposits from shallower depths, expression of the falling-stage (FSST), below the upper SB (Fig. 4). Similar small-scale depositional sequences, expressed through fining-upward cycles no less than 40 m-thick, are also present in synthem S3 at Case al Poggio, near Siena (biozones MPL3-MPL4a: Bianucci et al., 2001) and at Castelfiorentino (biozone MPL4b: Benvenuti et al., 2014).

3.3.2 Piacenzian EDS (synthems S4-S6)

As depth of deposition shallows during the Piacenzian, and cycles of sea level variation widen, the cyclic stacking of EDS becomes the typical depositional theme (Benvenuti et al., 2007, 2014; Dominici et al., 2008). In FB, the northwesternmost basin, depths remain considerable and facies change is more subtle. Pliocene at Orciano Pisano is traditionally assigned to the “Blue Clays” formation (Bossio et al., 1997), but two distinct bodies were evidenced since the late nineteenth century (D’Ancona, 1867). The lower one is formed by grey claystone with very rare shells, the upper one by muddy, very-fine grained gray sandstone richly fossiliferous. The lower part of the latter interval outcrops at the foothill of the small town of Orciano Pisano, around the locality Case Nuove (Bianucci and Landini, 2005; Berta et al., 2015). Here a laterally persistent shell bed is dominated by the turritellid *Archimediella spirata*, overlain by a 25 m-thick monotonous sandstone interval with intercalated shell beds or sparse shells, becoming muddier upward. The *Archimediella* shell bed is interpreted as the TS of an EDS, coinciding with the lower SB of a lower Piacenzian EDS. The overlying muddy fine-grained sandstone is the TST and HST (MPL5a: Dominici

et al., 2009; Fig. 6). At shallower settings, in all basins to the East and South of FB, EDS of synthem S4 take the form of an alternation of mudstone and carbonate (facies F5 and F8: Nalin et al., 2016), or mudstone and sandstone (facies F5 and F3-F4: Benvenuti et al., 2014; biozone MPL5a). Middle and upper Piacenzian EDS form an alternation of coastal mudstone and sandstone (facies F1-F2-F3: Benvenuti and Dominici, 1992; Dominici, 1994), with MFS and HST marked by a shell bed topped by a lower shoreface sandstone, or a shelf mudstone (facies F4-F5: Benvenuti et al., 2007; Dominici et al., 2008).

4. Distribution of large marine vertebrates

4.1 Geographic distribution

The MM geographic distribution is listed in Table 2. All MM that was geographically located, irrespective of stratigraphy, was plotted in Fig. 3. The largest number were recovered in FB in Northwestern Tuscany, with a peak at a few sites around the small town of Orciano Pisano, in the Pisa province. This coincides with the highest number of known species, including mammals, elasmobranchs, turtles, and large bony fishes. MM is also abundant in the province of Siena, around Volterra (VEB; Bianucci and Landini 2005), and around San Quirico, particularly rich with elasmobranchs according to the available data (SRB; Marsili, 2006). A fourth basin with a consistent number of findings is OOB, where cetaceans, sharks, sirenians and large teleost fishes have been unearthed.

4.2 MM stratigraphic distribution

4.2.1 Synthem S1-S2

In the vicinity of Saline di Volterra, one of the sites with remains of *Pliophoca etrusca* (Berta et al. 2015) and sperm whale (Tab. 2), S1 is represented by a bathyal mudstone (facies F6, biozone MPL1). Lower-middle Zanclean MM is otherwise absent, with the exception of S2 in OOB, where large marine vertebrates are concentrated at the maximum flooding interval, outcropping at Poggio alle Mura (Danise, 2010), Camigliano (Sorbi et al., 2012) and Arcille (Tinelli et al., 2012; Tinelli, 2013), in biozone MPL2. At Poggio alle Mura a slightly disarticulated balaenid whale skeleton (WOM1-1 in Tab. 2) has been excavated in contact with the laterally-persistent *Haustator* shell bed at the maximum flooding interval (Figs. 4, 5; Danise, 2010; Tinelli, 2013). At Camigliano and Arcille, tens of km from Poggio alle Mura, the *Haustator* shell bed is associated with other articulated skeletons and isolated MM remains, including several specimens of the sirenian *Metaxytherium subapenninum* (MOM2-1-5), large bony fishes, rays, and sharks *Carcharias taurus*, *Carcharhinus* sp., *Galeocerdo cuvieri*, and *Squatina* sp. (Sorbi et al., 2012; Tinelli et al., 2012; Tinelli, 2013). A partial skull of the delphinid *Etruridelphis giulii*, with right and left dentaries fractured, but nearly complete and with most teeth still in their alveoli (DSi2-1 in Tab. 2), was recovered southeast of Siena, near Chianciano Terme (SRB) at the top of a mudstone (facies F6) intercalated with turbiditic sandstone beds (Facies F7), topped by a monotonous mudstone interval (upper Zanclean, uppermost part of biozone MPL3: Bianucci et al., 2009b). In the same basin, two undetermined beaked whales and bones of four different specimens attributed to *Metaxytherium subapenninum* were found in the middle of a fining-upward succession, at the boundary between biozones MPL3 and MPL4a (Bianucci et al., 2001; WSi2-1-2 and MSi2-1-4 in Tab. 2), suggesting that findings at the two SRB localities belong to the same MFS, in the upper part of S2. Three specimens of marine mammals were identified in Synthem S1, seven in Synthem S2.

4.2.2 Synthem S3

The upper part of S3 yielded an articulated and well-preserved balenopterid skeleton found in a 30 m-thick mudstone succession at Castelfiorentino, in the Elsa basin (EB, WEI3-1). These strata are richly fossiliferous, with several shell beds with epifaunal cemented taxa, such as vermetid gastropods, oysters and corals (Facies F5), in a normal-polarity magnetostratigraphic interval (Gauss chron: Andrea Albanelli, personal communication, 1999). Based on physical stratigraphic correlation, this can be assigned to a lower Piacenzian HST. A skull of *Hemisyntrachelus* sp. (Aldinucci et al., 2011) and partly articulated vertebrae and costae of a dolphin skeleton (Arbeid et al., 2015) were recently excavated a few hundred meters apart one from the other, along a monotonous mudstone S3 succession intercalated with several *Ostrea* and *Serpulorbis* shell beds, near Certaldo (facies F5, DEI3-1-2 in Tab. 2). Overall, specimens of marine mammals attributed to Synthem S3 were eleven (some are uncertain and may come from the upper part of S2: Tab. 2).

4.2.3 Synthem S4

Synthem S4 yielded the highest abundance and species-richness of the Tuscan MM. In the Fine basin, the locality of Orciano Pisano is represented in Table 2 by 18 records of whales, dolphins (a partial skeleton: Bianucci, 1996; Bianucci et al., 2009), seals (Berta et al., 2015), tens of other unidentified cetacean elements, hundreds of shark teeth and vertebrae (Fig. 3), and sea turtles, large bony fishes, and sea birds hosted in museum collections (Bianucci and Landini, 2005; Marsili, 2007b; Cioppi and Dominici, 2011). In locality Case Nuove, a single middle Piacenzian transgressive surface has yielded a whole, articulated whale skeleton (Figs. 4, 6), teeth of blue and white sharks and bones of sea birds (Dominici et al., 2009). From the same site comes a skeleton of the monk seal *Pliophoca etrusca* (Berta et

al., 2015), and possibly many other museum specimens labelled “Orciano Pisano”, suggesting that this interval forms the most prolific bonebed of the region (Tab. 2). A few cm above the *Archimediella* shell bed, glauconitic and deeply bioeroded whale bones (Danise, 2010), associated with ichnological evidence of the activity of *Osedax* bone-eating worms (see Higgs et al., 2012), lie in a bioturbated muddy, fine-grained sandstone with a complex boxwork of *Ophiomorpha* and *Thalassinoides* trace fossils (Fig. 6), associated with a diverse paleocommunity of molluscs and other benthic invertebrates with complex trophic connections (Dominici et al., 2009; Danise et al., 2010). At the boundary between FB and VEB, an incomplete skull and skeleton of the dolphin *Etruridelphis giulii* was recovered near Lorenzana (Lawley, 1876; Bianucci, 1996; Bianucci et al., 2009b), at a locality associated with a muddy sandstone interval (facies F4) in synthem S4. An incomplete and articulated mysticete was recovered in a sandstone at San Gimignano, associated with pectinid bivalves (Fig. 7; facies F3; Elsa basin, EB: Danise and Dominici, 2014), here tentatively assigned to the uppermost part of the synthem (WEI4-1 in Tab. 2). Overall, specimens of marine mammals attributed to Synthem S4 were 28.

4.2.4 Synthem S5

The next MM richer stratigraphic interval are Piacenzian deposits of synthem S5. An incomplete and disarticulated balaenid skeleton was found in intertidal deposits of the lower part of the synthem, at Casenuove (facies F2, EB; Bianucci et al., 1995; Collareta et al., 2016; WEI5-1 in Tab. 2). A large balaenid was recovered higher up section, a few meters above a laterally-continuous very thick and complex *Haustator vermicularis* shell bed, up to 2 m-thick (Benvenuti et al., 1995a), traced laterally for 2 km to the east of San Miniato (“*Turritella strata*”, De Stefani, 1874), and forming a surface of transgression within the TST of S5 (Benvenuti et al., 2007; 2014; Dominici et al., 2008). The MFS is formed by a

Glycymeris insubricus shell bed, separating around San Miniato shoreface sandstone (facies F3 or F4) from offshore mudstone (F5; Benvenuti et al., 2007). The balaenid skeleton was almost articulated and bioeroded, closely associated with teeth of the great white shark and other scavengers (Borselli and Cozzini, 1992; Bianucci et al., 2002; Danise and Dominici, 2014), in the early HST of synthem S5 (Benvenuti et al., 2007; Dominici et al., 2008; WEI5-2 in Tab. 2). The *Glycymeris* shell bed can be traced laterally for several km. In the vicinity of Fiano, it includes bioeroded and encrusted gravels inherited from underlying successions, interpreted as revinement deposits. The interval of maximum flooding is marked by the stacking of at least three distinct shell beds, all including a high-diversity association with bioeroded and encrusted shells. This situation suggests that balaenid WEI5-2 lies in correspondence of an interval of low rates of sedimentation. In SRB, near Castelnovo Berardenga, shelfal mudstones (facies type F5) have yielded MM remains at a few localities. Delphinid remains were found at the “I Sodì” quarry and at Troiola (DSi5-1-2, in Tab. 2). Bones of a beaked whale and undetermined mysticetes are reported from Guistrigona (Manganelli and Benocci, 2014) and a fragmentary specimen of the monk seal *Pliophoca etrusca* from Castelnovo Berardenga (Berta et al., 2015). A very rich shark fauna, including sawsharks, thresher, frilled, bluntnose sixgill, bramble, gulper, kitten, sand tiger, shorten mako, basking and requiem sharks, an association suggesting an upper slope paleoenvironment for the surroundings of Castelnovo Berardenga Scalo (Cigala-Fulgosi et al., 2009; the same mudstone interval at the same locality has been interpreted as a shelfal deposit: Martini et al., 2016). Judging from historical accounts (Capellini, 1883), the killer whale *Orcinus citonensis* (DCh4-1) was recovered in a sandy unit lying on top of a thick mudstone interval (S3 or S4) and is tentatively assigned to S5. Specimens of marine mammals attributed to Synthem S5 were 10.

4.2.5 Synthem S6

A fairly complete skeleton of a large balaenid whale was recovered in the second half of the 19th Century in the EB near Montopoli Valdarno (Capellini 1905), in open shelf strata attributable to the interval of maximum flooding of S6 (WEI6-1 in Tab. 2). The MFS of S6, of uppermost Piacenzian age, is marked in EB by a laterally continuous *Pteria phalenacea* shell bed, with a high-diversity association of macroinvertebrates, including a rich decapod paleocommunity (Garassino et al., 2012). In the vicinity, near Palaia, a right whale (*Eubalena* sp.) was recovered in 1974 in sandy mudstones, in association with mollusc shells and teeth of the great white shark (*Carcharodon carcharias*: Bisconti, 2002; Sorbini et al., 2014). A tightly articulated balaenopterid skeleton was found at Sinalunga (WCh4-1 in Tab. 2), in deltaic sandstones and conglomerates (Fig. 8). Marine mammal specimens attributed to Synthem S6 were three.

4.3 MM facies type distribution

In an ideal deepening-up gradient, multi-element findings of marine mammals are very rare in intertidal and very shallow subtidal paleoenvironments (facies type F2, 1,5%: Fig. 9A), moderately represented in delta or shoreface sandstones (facies type F3, 4,6%), most abundant in sandy mudstone of open shelf settings (facies type F5, 70%), rare in outer shelf and bathyal sediments (facies type F6-F7, 1,5%). The most pristine and complete skeletons are associated with gravelly well-sorted sands from event sedimentation, suggesting a negative relationship between taphonomic loss and sedimentary processes at delta fronts.

In the tightly articulated and pristine Sinalunga balaenopterid (WCh4-1 in S4-CB), the cortical surface of the tightly-connected vertebrae is practically intact, and carpal, metacarpal and phalanges of the flipper are in perfect anatomical connection, as if a sudden depositional event buried a fresh carcass (Fig. 8; similar pristine skeleton are found in deep

water turbiditic succession: Stinnesbeck et al., 2014). Another pristine and tightly articulated skeleton, belonging to a killer whale (WCh3-1 in CB), was collected at Cetona in the second half of the 19th century, in a locality associated with sandstones, also possibly of deltaic origin. Large vertebrates embedded in fine-grained, muddy matrix (shelf deposit formed below storm wave base) and those associated with laterally-persistent shell beds (condensed deposits) are slightly disarticulated and fairly complete, showing signs of long permanence in a low energy, well-oxygenated seafloor before the final burial. Bioerosion of bones caused by phototrophic cyanobacteria and algae, heterotrophic fungi and bacteria (Orciano balaenopterid WFi4-1 in S4-FB), and eventually by whalebone-eating siboglinid worm of genus *Osedax* (on a ziphiid humerus, WFi4-14: Higgs et al., 2012), occurred at condensed intervals, in association with glauconite (Danise, 2010). In one instance, a condensed shelly interval is traced for a few kilometers, connecting bioturbated shoreface sandstones yielding slightly disarticulated sirenian skeletons (MOm2-1, MOm2-2) and other MM (Tinelli, 2013), with open shelf deposits yielding slightly disarticulated whale remains (WOM2-1 in S2-OOB).

Regarding a sequence stratigraphic interpretation, pristine skeletons from delta front sediments can be part of the FSST (Fig. 4; in an alternative interpretation it may belong to the early TST, when incised valleys are filled with coastal deposits). TST deposits account for 6,2% of cases (Fig. 9B). Much more frequently, articulated or slightly disarticulated skeletons are associated with the HST of the corresponding depositional sequence, lying above the MFS (56,9% of cases), or within the maximum flooding interval, above a laterally-persistent shell bed (10,8%). Bone bioerosion is less pronounced in the late HST, when sedimentation rate increases (Castelfiorentino whale WEI3-2 in S3-EB).

4.4 MM abundance and species-richness

Some taxa need revision, but a conservative estimate of the different morphologies suggest that at least 17 marine mammal species lived in the NWMS during the Pliocene (possibly more than 20, an estimate for the whole epoch, i.e., 5,332-2,588 Ma), against nine presently living in the same area (plus two occasional visitors). Among the cetaceans, six families were present, against only five presently living in the Ligurian Sea (Tab. 3). The most abundant Pliocene species of Tuscany are the sirenian *Metaxytherium subapenninum* (N = 10), the dolphin *Etruridelphis giulii* (N = 7) and the delphinid *Hemisyntrachelus cortesii* (N = 5; Tab. 4). Overall abundance and species richness are not randomly distributed, but are maximum in Piacenzian strata of synthem S4, dated at 3.2-3.0 Ma (Tab. 2, Figs 2, 9C-D), particularly in the FB and VEB (Tab. 4). A species list of marine mammals summing up fossils found around Orciano Pisano and at La Rocca, near Volterra, yields a species richness (S) of 13. This Piacenzian peak in marine mammal diversity is matched by the fossil record of sharks, also maximum near Orciano Pisano (S = 27), with a second peak in S5, around Castelnuovo Berardenga (SRB, S = 16). Differently from the marine mammals, showing a complete turnover from the Pliocene to the recent (Tab. 3), 34% of Pliocene shark species are still extant in the NWMS (Tabs. 5-6). Marine mammal frequency and diversity gradually decreases in S5-S6.

5. Paleoenvironment-fauna relations

The paleoecology of Pliocene MM of Tuscany can be reconstructed by comparison with the ecology of their modern relatives. This approach can be applied at the family or genus level for marine mammals (Tab. 3), at the genus or species level for sharks (Tab. 5). With the only exception of the sirenian *Metaxytherium subapenninum*, feeding on seagrasses (Domining, 2001), and small demersal sharks (e.g., catsharks, frilled sharks), all MM studied here are pelagic forms that had no direct connection to conditions at the seafloor (Tabs. 4

and 6). The paleoecology of benthic habitats informs however on the situation of the overlying water column in terms of factors that matter to the distribution of pelagic organisms, such as water depth, salinity and nutrient levels. In the second place, since all MM remains after death ultimately sink to the seafloor, benthic paleoecology is also a means to understand taphonomic controls on MM distribution. Cluster analysis based on the distribution of mollusc species in 72 samples resulted in the identification of four main groups of samples, roughly corresponding to the four main facies types recognised based on lithology and sedimentary structures (F2-F5 in Tab. 1). Clusters are formed by samples from outer shelf and upper slope (three samples), open shelf (34 samples), shoreface (23 samples), and transitional settings, such as brackish-water coastal lagoons and tidal flats (12 samples: see Supplement Material, Fig. S1). ANOSIM confirms that sedimentary facies type can broadly predict what benthic assemblage it will yield (Tab. 7; general $R = 0,632$). The difficulty to discriminate between upper and lower shoreface facies, and between shoreface and open shelf facies is confirmed by overlaps in sample distribution in the NMDS ordination diagram (Fig. 10A). Samples AG1, MON1 and MON2 allow to re-interpret the associated sandstones, originally included in upper shoreface facies type, as offshore deposits. The presence of gravels and cobbles intensely bioeroded by bivalves and polychaete (ichnofossils include *Gastrochaenolithes*, *Meandripolydora* and *Caulostrepsis*) and encrusted by balanids, oysters, serpulids, and bryozoans, suggests they are part of condensed beds resulting from transgressive pulses (hiatal concentrations). The relationship between facies type and mollusc association is broadly summarised in the following paragraphs (see online Supplement Information for a list of characterising species).

5.1 Coastal lagoon, tidal flat and embayment

Intertidal faunas are always associated with facies type F2 and are characterised by low-diversity associations, sometimes with less than 10 taxa and dominated by one or two species, including species today living in brackish waters of the Mediterranean, at intertidal or very shallow subtidal depths (e.g. *Cerastoderma edule*, *Nassarius reticulatus*, *Scrobicularia plana*: Pérès and Picard, 1964). Facies type F2 is also associated with samples having a species richness higher than the preceding and including species typical of seagrass bottoms and known to withstand moderate changes of salinity. In only one instance a large vertebrate was associated with intertidal deposits (a balaenid, lying on top of large wood fragments: Bianucci, 1995; Collareta et al., 2016).

5.2 Upper shoreface

Facies type F3 is associated with a high-diversity assemblage representing a paleocommunity dominated by suspension feeders adapted to shifting sandy bottoms, with bivalves typical of modern shoreface sandy bottoms (e.g., families Glycymeridae, Tellinidae, Donacidae and Veneridae). Among extinct species of this recurring assemblage, some are large-sized or have very thick shells. Some species of this group indicate the presence of vegetated bottoms. Small pyramidellid gastropods are parasitic on echinoderms, also typical of the upper shoreface.

5.3 Lower shoreface

Species richness further increases in collections associated with facies type F4 (lower shoreface). Species typical of this recurring assemblage include both suspension-feeding and detritus-feeding bivalves and gastropods. The following gastropod families are usually represented by several species: Trochidae, Rissoidae, Cerithiidae (from vegetated

bottoms), Naticidae, Muricidae, Turridae, Conidae, Terebridae, Bullidae, Cylichnidae (carnivores), Pyramidellidae (echinoderm parasites). Many bivalve species occur in both facies types F2 and F3. At three different sites and at different stratigraphic units large marine vertebrates, including mysticetes, sirenians and sharks, were recovered in association with shell beds dominated by the gregarious turritellid gastropod *Haustator vermicularis* (Fig. 6C-D).

5.4 Offshore and upper slope

Another important set of species recurred in facies type F5 (mudstone deposited in offshore bottoms at shelf depths). Among characterising gastropods are the suspension feeders (*Turritella tricarinata*, *Archimediella spirata* and *Petalochoncus intortus*), deposit feeders (*Aporrhais uttingeriana*) and carnivores or scavengers (*Epitonium frondiculoides*, *Nassarius semistriatus*, *Mitrella nassoides*). Also the bivalves occupy many different ecological niches (e.g., infaunal detritus feeders, epifaunal suspension feeders, either free-living, byssate, or cemented). Outer shelf and uppermost bathyal sediments from F6 mudstones, studied at only one location, are characterised by a separate set of carnivorous gastropods and by a few small bivalve species. Ubiquitous molluscs include species found from intertidal to outer shelf depths (e.g., *Corbula gibba*), and those preferential of open marine waters, from lower shoreface to outer shelf. Most multi-element findings of marine mammals are associated with sandy mudstones from open shelf settings, below storm wave base, also in association with an *Archimediella spirata* shell bed (Fig. 6C-D). No molluscs were recovered in facies type F7, with the exception of bathyal mudstone in the lowermost Pliocene of FB, associated with sparse specimens of the gryphaeid epifaunal bivalve *Neopycnodonte navicularis* (not sampled).

5.5 Carbonate platform

A recurring benthic assemblage associated with the highly fossiliferous facies type F8, not included in the quantitative analysis, consists of the large pectinid bivalves *Gigantopecten latissima* and *Hinnites crispus* and by a mixture of photozoans (large benthic foraminifera), hard-substrata dwellers (brachiopods, echinoderms), encrusters (red algae, bryozoa) and bioeroders (clionid sponges). All fossil-rich carbonates are associated with synthem S4, in the mid-Piacenzian (Fig. 2, see also Nalin et al., 2016).

6. Paleodepths

Multivariate techniques are usefully applied to stratigraphic and paleobiologic analysis (Scarponi and Kowalewski, 2004). We used the results of the ordination analysis to estimate absolute depths of the final resting place of some large vertebrates listed in Tab. 2. Samples in the NMDS ordination plot following a water depth gradient, with shallower samples to the left (low values of NMDS axis 1) and deeper samples to the right (high values of axis 1). Therefore, NMDS values of axis 1 can be used as a proxy for relative water depth. We calculated absolute palaeodepths by fitting a logarithmic regression curve between absolute paleodepth of 23 modern species (data from MARBEF database), common in our dataset, and the values on NMDS axis 1 (Supplement Material: Tabs. S1-S4). The regression analysis, with $R^2 = 0.813$, indicates that scores along the nMDS main axis are a good predictor of the preferred depth for the 23 modern species (Fig. 10B, inset), thus supporting the bathymetric interpretation. This allowed to estimate the absolute depth of the 72 samples, which ranged from 0.4 m to 365 m, i.e., from intertidal to upper slope depths (in accordance with a previous estimate of absolute paleodepths in the upper part of the Pliocene of EB, based on counts of foraminifera: Dominici et al., 2007). Facies type F2 is

deposited at 0-5 m depth, F3 at 3-30 m, F4 at 10-100 m, and F5 at 40-300 m, F5 at around 350 m depth (only one sample: Fig. 10B). The position of fossil cetaceans, sirenians, pinnipeds and large sharks (respectively W, M, P and S, N = 13) was plotted near the corresponding shell bed in the NMDS ordination. The resulting pattern shows that all MM considered, associated with open shelf settings and with facies types F4-F5, cluster around -100 m, spanning -30-300 m.

7. Factors of the Pliocene NWMS marine megafauna fossil record

The detailed sequence-stratigraphic framework and the abundant shell beds, offering an independent check on sedimentary facies distribution by benthic paleoecology, allows also to explore factors behind MM geographic and stratigraphic distribution, and to sort out evolutionary, ecological, and taphonomic drivers of this particular fossil record. NWMS Pliocene distribution can be compared with similar studies in different settings and at different times, to draw conclusions on the general quality of MM fossil record.

7.1 Evolutionary control

Available data allow a meaningful comparison of NWMS MM diversity across the Pliocene, particularly detailed for marine mammals (Fig. 9D). Although the study suggests a facies control, with MM remains being generally associated with late TST-HST muddy sandstones and mudstones from lower shoreface and offshore shelf paleosettings, the temporal pattern of biodiversity recorded on a regional basis likely reflects also a paleobiologic phenomenon, since lower shoreface and offshore shelf sediments are represented in all synthems. Marine

mammals are unrecorded in the thickest part of synthem S1-2 (Zanclean, mostly bathyal mudstone and turbiditic sandstone), but they are present in S2, in OOB (Zanclean shoreface and open shelf deposits) and SRB (Zanclean upper slope deposits). MM gradually rises in S3, at the upper Zanclean-lower Piacenzian, whereas it suddenly peaks in synthem S4, where species richness of both marine mammals and sharks is highest. S5 reflects a lower diversity of marine mammals, but still a high diversity of sharks, while values of both groups drop to the lowest abundance and species richness in S6, during the upper Piacenzian-lowermost Gelasian (Fig. 9D).

The S4 diversity peak coincides with the middle part (3.264–3.025 Ma) of the Piacenzian, a time interval in which the earth experienced global average temperature 1.84 °C–3.60 °C warmer than the pre-industrial period (Dowsett et al., 2013). Climatic impact is testified by the widespread occurrence of carbonate deposits in S4 (Fig. 2), with sedimentary facies indicative of warm-temperate to subtropical conditions, with summer sea-surface temperature considerably warmer than 20°C and winter temperatures colder than 20°C (Nalin et al., 2016). This suggests a causative link between global climate and biodiversity, S4 diversity peak recording a global phenomenon, possibly an increase of speciation rate connected with global warming. Similarly, we propose that the lower diversities recorded at S5-S6 are the regional expression of an increase in extinction rate related to climatic cooling and global sea-level drop, ultimately leading to the global MM extinction event recorded on a coarser scale at the Pliocene-Pleistocene boundary (Pimiento et al., 2017): the finer stratigraphic resolution adopted here suggests a stepwise extinction event. NWMS data also point to a selective effect, extinction being recorded by marine mammals, with a 100% regional turnover between Pliocene and Recent (Tab. 3), but not as much by the shark fauna, with 34% of the species still living in the Mediterranean, while still others have shrank their distribution to subtropical latitudes. Also the Piacenzian teleost fish fauna (Cigala Fulgosi et al., 2009) and benthic molluscs (Raffi et al., 1985) show a high percentage of

holdovers, suggesting that marine mammals have been particularly prone to climatic change (see also Steeman et al., 2009).

Estimates of body size in Pliocene mysticetes of Tuscany, with several specimens reaching 10 m (Danise and Dominici, 2014: seven specimens in Tab. 1), are comparable to global values derived from the literature for this time interval (Lambert et al., 2010), confirming that NWMS baleen whales were larger than their Miocene analogues, and smaller than modern forms. As with regard to odontocetes, the most common Pliocene delphinid, *Etruridelphis*, was larger than the modern analogue *Stenella* (Bianucci et al., 2009b). The same is true for *Hemisyntrachelus cortesii*, larger than modern *Tursiops* (Bianucci, 1997a). On the other hand, the largest extant delphinid *Orcinus orca*, reaching 9 m, is about twice as long as *Orcinus citonensis* (Heyning and Dahlheim, 1988). The high percent of holdover points to a more conservative figure for larger sharks (Tab. 5), but the presence in the Pliocene of the gigantic *Carcharocles megalodon* and some large thermophilic species today restricted to lower latitudes suggests that impoverishment of the fauna is coupled with an average decrease in size (Marsili, 2008). Comparing sizes suggests an overall restructuring of NWMS MM during the last three million years (see also Bisconti, 2009).

7.2 Ecological control

The composition of the Pliocene NWMS MM is affected in the first place by the availability of food. At the lowest trophic level, inferring from the ecology of the modern MM (Tabs. 3, 5), we found herbivore sirenians feeding on seagrasses (trophic level, TL = 2,0), all others being carnivores, thus having TL > 3,0 (Fig. 11). The lowermost levels among the carnivores were occupied by baleen whales and whale sharks (TL = 3,2-3,4), filter-feeding on pelagic crustaceans and fishes (krill). Roughsharks, catsharks, and houndsharks, with many species living in the modern Mediterranean, have a relatively species-poor Pliocene record,

probably due to a preservation bias related to their small size. At TL = 3,7-4,0 were one species of catshark and a monk seal, both feeding near the seafloor on crustaceans, teleost fishes and cephalopods. At TL = 4,1 were sandbar, tiger and blue sharks, feeding on teleost fishes, cephalopods and on marine mammals. At the same level, in slope environments, kitefin sharks mainly fed on other sharks. The majority of MM species were found at TL = 4,2, with smaller odontocetes (three species) and 11 species of sharks, including several requiem sharks, a houndshark and a hammerhead. Larger dephinids, sperm whales, beaked whales, together with mackerel, sand and sand tiger sharks, occupied high trophic levels (TL = 4,3), followed at the top of the global NWMS food web by white shark, megalodon shark, one species of sevengill shark (genus *Notorynchus*), and killer whales, all feeding on marine mammals and smaller sharks (TL = 4,4-4,7). With no exception, all highest levels encountered in modern NWMS offshore pelagic and nearshore communities were occupied during the Pliocene by an analogous MM, often by the same species (large sharks), or by congeneric or con-familial species (marine mammals: Tab. 3, 5). The Pliocene pelagic ecosystem, typified by the mid-Piacenzian S4 association, must have been however trophically more diversified (Fig. 11), including aquatic megaherbivores, several balaenid filter feeders, larger and more diverse dolphins, and sharks species today extinct (e.g., *Carcharocles megalodon*) or restricted to tropical seas (e.g., *Galeocerdo cuvier*). With the exception of TL = 2, all Pliocene NWMS MM were either apex predators of their community, or mesopredators, occupying trophic positions below apex predators. The definitions of apex predators (or top predators) and mesopredators are relative and to an extent context-dependent (species that in a contact are apex predators, in another are mesopredators, e.g., Estes et al., 1998). Since predation is a trophic interaction in which one animal (predator) consumes another (prey) as a source of energy (food), irrespective of the means by which this is accomplished (Lourenço et al., 2013), filter-feeding baleen whales can be considered apex-predators of their community (e.g., Lewiston et al., 2004; Notarbartolo di Sciara et al.,

2016). Among sharks, 68% of living Mediterranean elasmobranches are ranked as top predators, with a trophic level of 4 or more (Goffredo and Dubinsky, 2014, including superorder Batoidea, against 22% of teleost fishes), an estimate that can be extended to Pliocene NWMS MM. Relationships between apex predators and mesopredators are complex and hard to define in ecology, involving predation on other predators (intraguild predation, combining competition and predation: Polis et al., 1989), where consumption and competition need to be proved (Lourenço et al., 2013). In the Mediterranean Pliocene, paleontological evidences of carnivores serving as food to MM include killing of prey (Bianucci et al., 2010) and scavenging (Cigala Fulgosi, 1990; Bianucci et al., 2002; Dominici et al., 2009). The occurrence of intraguild predation must have been far more extended than what taphonomy can prove, however, given a Pliocene diversity of NWMS very large raptorial feeders higher than the modern, including the killer whale *Orcinus citonensis*, the large delphinid *Hemisyntrachelus cortesii* (phylogenetically related to the modern killer whale: Murakami et al., 2014), the white shark *Carcharodon carcharias*, the largest shark of all times *Carcharocles magalodon* (Marsili, 2008), and a diverse association of smaller carnivores, including monk seal, delphinids and sharks with $4,0 < TL < 4,2$ (Tab. 3, 5), candidate prey for larger raptorial feeders. This interaction likely exerted in its turn a control on community structure at lower trophic levels through processes like “mesopredator release” and trophic cascades (Roemer et al., 2009), eventually linking pelagic and nearshore communities, including benthic animals and plants, like in many modern ecosystems (Estes et al., 2011, 2016), down to slope depths (e.g., Parrish, 2009). Large raptorial feeders could exert a control on the diversity of the filter-feeding MM, like it has been suggested on a global scale since the Miocene (Lambert et al., 2010), and for the Pliocene by Bisconti (2003), when baleen whales were more diversified, both in terms of species richness, size range, and feeding strategies, including both skim- and lunge-filter feeding (respectively balaenid and balaenopterid whales: Berta et al., 2016; Hocking et al.,

2101
2102 2017), minimising competition for food and diversifying spatial niches (see also Marx et al.,
2103
2104 2017, for the upper Miocene). A further important top-down control on community structure
2105
2106 is suggested by modern studies on the role of baleen and sperm whales as nutrient vectors,
2107
2108 both in horizontal (during seasonal migration) and vertical direction (during daily feeding
2109
2110 migration: Roman and McCarthy, 2010), a process particularly important in oligotrophic seas
2111
2112 (Alleger et al., 2017), like large sectors of the Mediterranean. Finally (literally, after death),
2113
2114 MM becomes a detrital sources of energy and habitat for deep sea whale-fall communities
2115
2116 (Roman et al., 2014; Smith et al., 2015), with paleontological evidence available for the
2117
2118 NWMS (Dominici et al., 2009; Higgs et al., 2012; Baldanza et al., 2013).
2119

2120
2121 Much evidence suggests that a bottom-up control on the structure of NWMS MM community
2122
2123 was exerted by wind-driven upwelling currents, through enhanced oceanic productivity and
2124
2125 concentration of preys. Over geologic time, these factors may have exerted a selection
2126
2127 favoring large size, triggering the Plio-Pleistocene emergence of whale gigantism in several
2128
2129 lineages (Slater et al., 2017). The largest among marine top predators can travel long
2130
2131 distances and cross oceans, but tend to congregate in shallow waters with abundant prey.
2132
2133 Baleen whales, abundant in boreal eutrophic waters (Woodley and Gaskin, 1996), with
2134
2135 population size under the control of food availability (Croll et al., 2005), always require high
2136
2137 prey density for efficient bulk filter feeding (Goldbogen et al., 2011). Six different species of
2138
2139 mysticetes currently foraging in the Southern ocean, among which the largest animals that
2140
2141 have ever lived in world oceans, exploit the high biomass of Antarctic krill, their main food
2142
2143 resource. Great white sharks are abundant in the offshore of California (Jorgensen et al.,
2144
2145 2010), Australia-New Zealand, South Africa, and in the Mediterranean (Bonfil et al., 2005),
2146
2147 clustering in proximity of seal colonies, including monk seals. The distribution in the modern
2148
2149 Mediterranean is no exception to general rule, only in a smaller scale, with large sharks and
2150
2151 marine mammals congregating in productive areas, such as the Strait of Sicily and the
2152
2153 Balearic Archipelago (Tabs. 3, 5), with the second largest animal on earth, the fin whale,
2154
2155
2156
2157
2158
2159
2160

showing a movement pattern that parallels seasonal variability in available feeding habitat (Notarbartolo di Sciara, 2016), contributing to the horizontal transfer of nutrients. Tuscany faces the Ligurian Sea, where a deep-water upwelling current coming from Southeast convects nutrients to the water surface, leading to high levels of primary productivity in its western sectors, extending westward to the Provençal and Balearic Seas, with spring algal blooms. These waters host different trophic regimes in an otherwise oligotrophic Mediterranean Sea (Lazzari et al., 2012; Melanotte-Rizzoli et al., 2014; Stambler, 2014). Supporting a conspicuous biomass of zooplankton (Cuzin-Roudy 2011), the Ligurian Sea sustains large populations of fin whales (*Balaenoptera physalis*) and striped dolphins (*Stenella coeruleoalba*: Notarbartolo di Sciara et al., 2008). Mediterranean-resident fin whales have adapted to exploit localised mesoscale hotspots of productivity that are highly variable in space and time (Notarbartolo di Sciara et al., 2016), feeding behaviors possibly mediated by the depth of prey and species-specific behaviours, allowing to minimise competition with other large filter feeders (see Friedlaender et al., 2014). The sperm whale *Physeter macrocephalus* uses habitat across a range of depths and a specialised diet (Rendell and Frantzis, 2016), gathering along NWMS steeper slopes, where water currents allows for higher trophic level biomass (Pirodda et al., 2011). Beaked whales are particularly abundant in the Ligurian Sea and central Tyrrhenian Sea, preferring submarine canyons at slope depths (Podestà et al., 2016). The largest among common Mediterranean delphinids, *Grampus griseus*, is frequent in the Ligurian and Tyrrhenian seas where it forages on cephalopods at depths 300-1500, where upwelling currents are most effective (Azzellino et al., 2016).

The NWMS existed as a Liguro-Provençal back-arc oceanic basin since the upper Miocene, when Sardinia rotated to its present position (Gattacceca et al., 2007), and in coincidence with the formation of the hinterland basins here under study (Muttoni et al., 2001), justifying the assumption that the modern NWMS trophic regime is a feature that dates back at least

to the upper Miocene. Consistently, diatomites deposited in Tuscany during the Messinian, before the salinity crisis (Bossio et al., 1997; Roveri et al., 2014), indicate that high productivity was a primitive feature of the Ligurian Sea. Paleontological evidences include the high diversity and abundance of Pliocene MM in FB, the closest to modern upwelling areas (Fig. 3), and the association of several MM with “*Turritella* beds”, turritelline gastropods showing gregarious habit and high abundance in areas of high primary productivity (Allmon, 1988). Both a comparison with the modern and paleoecologic data strongly suggest that a wedge of NWMS nutrient-rich waters intersecting the Tuscan shelf exerted a strong bottom-up control on community structure through mixing and upwelling of nutrients, stimulation of phytoplankton blooms, followed by zooplankton increase, while nekton and vertebrates tracked plankton concentrations, as is typical of modern upwelling systems throughout the world (Polis et al., 1997). High surface primary productivity in the study area during the Pliocene would have caused a concentration of detritus-falls, supporting a diverse community of deep-sea scavengers.

7.3 Taphonomic control

Taphonomic data on articulation and completeness of MM specimens are available for large mammals, the shark record being formed mainly by isolated teeth collected through superficial picking during the years (Cigala Fulgosi et al., 2009), with only a few contextualised studies (e.g., Bianucci et al., 2002; Dominici et al., 2009). Our record includes several marine mammals with a high degree of articulation (14%) and completeness (27%), or with at least cranial and post-cranial elements of the same individual (38%). All articulated specimens and the most complete skeletons are associated with shelf mudstones, usually forming the HST of the relative depositional sequence (Tab. 2), with two exceptions, one associated with delta-front, coarse-grained beds (WCh6-1, Fig. 6), another from outer shelf

or upper slope sediments (DSi2-1). These data point to a strong environmental control on the quality of the MM fossil record, at least concerning articulation and completeness of skeletons, very shallow and very deep (bathyal) depths being generally unfavourable to the preservation of MM bones (Fig. 9A). To explain these results, hypotheses are based on the available data on carcasses of MM in modern marine environments.

The biostratinomy of lung-breathing marine mammals depends on water depth (Allison et al. 1991; Smith et al., 2005). The vast majority of mammals are negatively buoyant and sink after death, but in shallow waters the low hydrostatic pressure allows putrefaction gases to develop and carcasses to resurface. Skeletons are scavenged (Dicken, 2008) and disintegrate while floating, leading to the preferential deposition of isolated bones. At deeper settings, where high hydrostatic pressure allows the carcass to lay relatively undisturbed on the seafloor, the skeleton becomes only slightly disarticulated after soft tissue removal (Reisdorf et al., 2012). The subsequent fate of deep-water, disarticulated skeletons depends on the nature of the scavenging fauna and other elements of the whale-fall community, and on the time of exposure on the seafloor (Boessenecker et al. 2014). At depths deeper than the slope breaks, where sedimentation rate is very low, carcasses are exposed for a long time and the skeleton is rapidly disintegrated. Time-series analyses carried out at modern whale-fall communities at slope depths (range 382-2893 m: Lundsén et al., 2010) suggests that carcasses up to 17 m are rapidly degraded, with the deepest whale carcasses disappearing after only seven years of exposure on the sea floor. Larger skeletons may persist on deeper settings for decades, but if not buried, they also ultimately undergo complete destruction. Although environmental forcing triggered by higher temperatures, active currents and sediment transport, clearly plays a role, a specialised whale-fall fauna rapidly consumes both soft and mineralised tissues. This fauna is characterised by low diversity and high abundance of microorganisms, most likely as a result of both specialisation to nutrient enrichment and high growth rates. Whale-fall habitats likely

undergo a temporal microbial succession from primarily heterotrophic to more heterotrophic/chemosynthetic metabolisms until the whale biomass is completely exploited (Smith et al., 2015). Of all the specialised taxa, bone-eating polychaetes of genus *Osedax*, with their soft root-like tissues that erode the bones to access nutrients (Tresguerres et al., 2013; Minamoto et al., 2017), are the primary cause of bone disintegration, particularly of denser bones (Higgs et al., 2011). Found also at shelf depths, but invariably in low abundance (Huusgaard et al., 2012; Higgs et al., 2014b), bone-eating worms occur in high numbers in the deep sea (Smith et al., 2015) where they act as biodiversity regulators (Alfaro-Lucas et al., 2017). The general paucity of novel taxa on shallow-water whale falls suggests that species-rich, specialised whale-fall communities develop only in the food-poor deep sea. Accordingly, among new animal species described in the recent literature from whale falls, only about 10% have been found on whale remains at depths of less than 260 m (12 out of 129 new species: Smith et al., 2015). Of all MM investigated so far, only whales are known to host a whale-fall community, but also carcasses of large elasmobranchs undergo rapid destruction at bathyal depths, teeth being all that eventually remains (Higgs et al., 2014a; teeth are also lost by sharks during feeding: Pokines and Symes, 2013). As a consequence, over geological time no large marine vertebrate is expected to be recovered at depths greater than the shelf break. On the opposite side, the lack of a biota specialised in exploiting large organic falls, coupled with higher rates of deposition in proximity of sediment sources, make it more probable that the most articulated and complete whale skeletons become part of the fossil record of shelf settings below storm wave base.

8. Comparison with other studies

The study of sedimentary facies uncovers some environmental factors that directly control the taphonomy of large marine vertebrates. Such abiotic drivers include water pressure,

2401 wave energy and sedimentation rate — three factors summarised by water depth, and
2402
2403 upwelling, bringing to the surface deep sea nutrients and concentrating preys. The
2404
2405 taphonomic pathway of large marine carcasses is also driven by biotic factors that change
2406
2407 in geological time in response to coevolution between bacteria, scavengers and their
2408
2409 substrates. We now explore the multifaceted nature of MM taphonomy by reviewing
2410
2411 Mesozoic and Cenozoic studies where sufficient data for stratigraphic paleobiology are
2412
2413 available.
2414
2415
2416
2417
2418

2419 8.1 Abiotic factors 2420 2421 2422 2423

2424 According to one of the few previous studies on the sequence stratigraphic distribution of
2425
2426 MM, Jurassic ichthyosaurs, plesiosaurs, and pliosaurs of the Sundance Seaway, in North
2427
2428 America, display facies control and are found primarily in offshore mudstone and at
2429
2430 condensed intervals at the maximum flooding surface, rather than shoreface and estuarine
2431
2432 sandstone (McMullen et al., 2014). Taphonomic data on Upper Cretaceous marine reptiles
2433
2434 and large fishes suggest that partially articulated and disarticulated skeletons are associated
2435
2436 with little biological activity and relatively rapid burial by muddy sediments, deposited in the
2437
2438 North American seaway during an interval of maximum flooding (Schemisser McKean and
2439
2440 Gillette, 2015). Upper Cretaceous mosasaur remains are particularly concentrated in fine-
2441
2442 grained shelf deposits in Europe (Jagt and Jagt-Yazykova, 2016). Complete, partially
2443
2444 articulated whale skeletons of archaeocetes, together with bones and teeth of sirenians and
2445
2446 sharks, are abundant in offshore marine flooding surfaces (MFS) in the Priabonian late TST
2447
2448 of Egypt. In a further parallelism with the Pliocene of Tuscany, Late Eocene well-articulated
2449
2450 whales are associated with rapidly accumulating shoreface sediments of the FSST,
2451
2452 comprising “*Turritella* shell beds” (Peters et al., 2009). During the Oligocene, eomysticete
2453
2454 whale bones were deposited at shelf depth below storm wave base. The occurrence of
2455
2456
2457
2458
2459
2460

sparse traces attributed to *Osedax* and the association with a glauconitic limestone testify to the exposure of bones on the seafloor without undergoing complete destruction (Boessenecker and Fordyce, 2014), in a manner similar to MM at Orciano (see Danise et al., 2010; Higgs et al., 2012). Burdigalian/Langhian MM recovered in the Antwerpen Sands, in Belgium (including isolated remains of a baleen whale, several odontocetes and a pinniped) are fragmented, worn and associated with clayey sandstone rich in glauconite, suggesting long exposure on the sea-floor (Louwye et al., 2010). Bones are concentrated at the base of a coarsening-upward succession, on top of shallow marine, coarse-grained sandstone, suggesting this is a surface of maximum flooding. An association of MM taxonomically comparable to that here under study is encountered in the Mio-Pliocene of the Purisima Formation, in Central California. Taphofacies differ in some aspects. The Pliocene of California yields laterally persistent bonebeds with polished and phosphatised bones, and abundant phosphate nodules that are absent in Tuscany, indicating times of higher sediment starvation during transgressive pulses, in an area of much stronger nutrient content (the California Current system is a northern-hemisphere analogue of the Peruvian upwelling system, associated with the economically most important fish stocks in the world: Mann and Lazier, 2006). Shoreface deposits indicate stronger wave energy, and the preferential absence of molluscs in bonebeds indicates chemical destruction of carbonate shells (Boessenecker et al., 2014, in a quantitative MM taphonomic study). Episodic sedimentation, however, causes the preferential preservation of articulated remains in the Californian offshore as in the Pliocene of NWMS. The late Miocene Pisco Formation in Perú offers another, more extreme example of MM taphonomy in a eutrophic setting. Here almost two hundred cetaceans, pinnipeds, and sharks were described in an exceptional state of preservation (Bianucci et al., 2015), within a monotonous succession of finely laminated white diatomites (Di Celma et al., 2015), suggesting very high primary productivity in an area of intense upwelling and volcanic activity. Algal blooms sustained high biomass of apex

predators (see Marx and Uhen, 2010), triggering at the same time anoxic conditions at the seafloor where MM carcasses remained intact (Brand et al., 2004; see also Gioncada et al., 2016; Marx et al., 2017, also in offshore paleosettings). Finally, the stratigraphic distribution of Pliocene cetaceans in western Emilia (Italy) shows an uneven distribution of findings (N = 24, dolphins and baleen whales) and a strong positive correlation with offshore mudstones (no findings in shoreface sandstones, rare occurrences in epibathyal mudstones: Freschi and Cau, 2016), paralleling the distribution of Tuscan Pliocene MM.

8.2 Biotic factors

Many reviews of Triassic (Camp, 1980; Hogler, 1992; Motani et al., 2008; Hu et al., 2011; Liu et al., 2014) and Early Jurassic marine reptiles (Benton and Taylor, 1984) report mostly good preservation and a high degree of completeness and articulation of skeletal material. This record might be partially controlled by prevailing anoxic or dysoxic conditions in the bottom waters of many Mesozoic fossiliferous deposits (e.g., Middle Triassic Besano and Guangling Formations, Lower Jurassic Blue Lias and Posidonia Shale Formations), which precluded organism activity within the sediment, and prevented predation or scavenging of the carcasses on the sea bottom (Beardmore and Furrer, 2016). Evidence of advanced levels of disarticulation or bone degradation (Martill, 1985; sauropterygians, crocodilians, ichthyosaurs and fishes from the Middle Jurassic Oxford Clay Formation), is mostly attributed to physical factors (e.g., weathering on the sea floor). Up to the early-Late Cretaceous, biological activity is testified by circumstantial evidence of scavenging (*Hybodus* teeth associated with marine reptile skeletons, Martill et al., 1994), and by the more common occurrence of microbial mats, grazers and encrusters (Martill, 1987; Meyer, 2011; Danise et al., 2014; Reolid et al., 2015), but lack traces of bone-eating worms and sulphophilic fauna typical of modern whale falls. The siboglinid *Osedax* is an evolutionary

novelty in possessing a root system that hosts heterotrophic mutualists and secretes bone-dissolving acids (Tresguerres et al., 2013; Miyamoto et al., 2017), and an ecosystem engineer (Alfaro-Lucas et al., 2017). Genus *Osedax* is today associated with whale falls worldwide (Taboada et al., 2015), but its impact on MM has changed in time. The oldest trace fossils attributable to a bone-eating fauna is found on early-Late Cretaceous plesiosaur and sea turtle bones (about 100 Ma: Danise and Higgs, 2015). Time estimates suggest that *Osedax* diverged from other siboglinids in the Middle Cretaceous (around 108 Ma: Taboada et al., 2015). However, if the bone-eating worm lives also on the bones of birds and terrestrial mammals (Rouse et al., 2011), its global nature and high species diversity in modern oceans suggest that whale falls, as complex and species-rich habitats, have been the most important biodiversity generators (Higgs et al., 2014b; Smith et al., 2015). And although the first appearance of *Osedax* is concomitant with the occurrence of large marine reptiles and teleost fishes in the oceans, long before whale evolution, and although their distribution is not limited to large carcasses (Pyenson and Haas, 2007), the radiation of ocean-going mysticetes at the Oligocene onset of the Antarctic Circumpolar Current (Fordyce, 2003) clearly increased available substrata worldwide. The steady increase of cetacean size during the Neogene, with a dramatic pulse in the last five million years, when Neoceti surpassed 10 m length and reached 30 m in the Pleistocene (Lambert et al., 2010; Slater et al., 2017), would have thus triggered a second and more massive radiation of bone-eating worms (Kiel and Goedert, 2006). Consistently with this hypothesis, the preferential distribution of modern *Osedax* in high latitude settings worldwide (Taboada et al., 2015) suggests that biodiversity hotspots coincide with the feeding grounds of larger cetaceans. Other bone-eaters of modern deep water whale-fall ecosystem belong to the group of abyssochrysid snails, with fossils found on Late Cretaceous plesiosaur (Kaim et al. 2008) and sea turtle bones (within a chemosynthesis-based association: Jenkins et al., 2017). Modern abyssochrysid whalebone-eaters of genus *Rubyspira*, hosting a specific and

exclusive microbiome (Aronson et al., 2017), split during the upper Eocene/lower Oligocene (Johnson et al., 2010). Species of *Rubyspira* benefited too from the radiation of ocean-going whales. Although scanty, available evidence on the geological history of bone-eaters thus makes the ephemeral nature of large carcasses in modern deep seas — and their absence in bathyal deposits of the Pliocene of NWMS — a larger-than-life model for the Mesozoic and the early Paleogene.

9. Conclusions

- 1) Sedimentary facies in the Pliocene of Tuscany are vertically stacked to form small-scale depositional sequences particularly in the upper half, Piacenzian part of the succession, with laterally-continuous shell beds marking transgressive surfaces and intervals of maximum flooding. Small-scale sedimentary sequences are stacked to form six major, unconformity-bounded stratigraphic units (synthems) of regional extension, forming a high-resolution framework to study the chronostratigraphic distribution of marine megafauna (MM).
- 2) Benthic biotopes, identified through a quantitative study of a large mollusc dataset, can be arranged to form an ideal onshore-offshore, bathymetric gradient, connecting terrestrial environments with deep sea epibathyal bottoms, consistently with the distribution of sedimentary facies. MM remains and shell beds are present in all marine paleoenvironments, but their distribution is uneven. The fossil record of large vertebrates, including marine mammals (abundance data) and sharks (presence-absence data), is particularly rich in sediments deposited between storm wave base and shelf break, moderately rich in shoreface sediments, very poor in epibathyal sediments. Paleobathymetric estimates suggest that a peak of abundance of better preserved skeletons occurs around 100 m depth, within the 30-300 m range.

- 2701
2702
2703
2704
2705
2706
2707
2708
2709
2710
2711
2712
2713
2714
2715
2716
2717
2718
2719
2720
2721
2722
2723
2724
2725
2726
2727
2728
2729
2730
2731
2732
2733
2734
2735
2736
2737
2738
2739
2740
2741
2742
2743
2744
2745
2746
2747
2748
2749
2750
- 3) Species-richness of MM and abundance of marine mammal remains steadily increase during the Zanclean, reaching a peak in synthem 4, possibly as a response to the mid-Piacenzian warm period, and gradually decrease until reaching a minimum in proximity of the Plio-Pleistocene boundary.
 - 4) Comparison with ecological data on a global scale, paleogeographic and topographic considerations on the North-Western Mediterranean Sea (NWMS), and the occasional association of Pliocene MM with *Turritella* beds, suggest that upwelling and high-nutrient conditions in the Northwestern Mediterranean offshore sustained during the Pliocene a rich community of apex predators and mesopredators. Higher taxonomic MM diversity in the NWMS during the Pliocene, suggests higher niche-partitioning with respect to the modern NWMS and a top-down control on community structure.
 - 5) A comparison with studies on the biota exploiting tissues of large food particles sunken on the seafloor, both modern and ancient, suggests that the poor epibathyal record of Pliocene larger vertebrates of Tuscany may be caused by the destructive action of bone-eating invertebrates. This biotic driver of the marine vertebrate fossil record was less efficient before the radiation of ocean-going whales in the late Eocene-lower Oligocene. On the other hand, bone-eaters played a major taphonomic role on a global scale after the Pliocene-Pleistocene increase in whale size.
 - 6) Due to resurfacing of carcasses in shallow waters, the MM fossil record through the Mesozoic-Cenozoic reviewed through stratigraphic paleobiology shows a strong facies control, being more continuous in rocks deposited below storm wave base and above the shelf-slope break during sea-level transgressions and highstands.

Acknowledgements

2751
2752
2753

We thank Michelangelo Bisconti (SCNHM) for helpful discussion on mysticete taxonomy and paleoecology, Giovanni Bianucci (DSTUP) for his insightful review on the first draft of

2754
2755
2756
2757
2758
2759
2760

2761
2762 this paper, and an anonymous reviewer. Maurizio Forli helped by sharing his thorough
2763
2764 knowledge of fossiliferous Pliocene outcrops of Tuscany.
2765
2766
2767

2768 **References**

2769
2770
2771
2772
2773 Alfaro-Lucas, J.M., Shimabukuro, M., Ferreira, G.D., Kitazato, H., Fujiwara, Y. and Sumida,
2774
2775 P.Y.G. 2017. Bone-eating Osedax worms (Annelida: Siboglinidae) regulate biodiversity of
2776
2777 deep-sea whale-fall communities. Deep Sea Research II
2778
2779 <http://dx.doi.org/10.1016/j.dsr2.2017.04.011>.
2780
2781

2782
2783
2784 Barría, C., Coll, M., Navarro, J., 2015. Unravelling the ecological role and trophic
2785
2786 relationships of uncommon and threatened elasmobranchs in the western Mediterranean
2787
2788 Sea. Mar. Ecol. Prog. Ser. 539, 225-240, doi: 0.3354/meps11494.
2789
2790

2791
2792
2793 Barría, C., Navarro, J., Coll, M., 2017. Trophic habits of an abundant shark in the
2794
2795 northwestern Mediterranean Sea using an isotopic non-lethal approach, Estuarine, Coastal
2796
2797 and Shelf Science, doi: 10.1016/j.ecss.2017.08.021.
2798
2799

2800
2801 Alderighi, L., Bianucci, G., Tinelli, C., Petri, A., 2011. Certaldo (FI). Località Canonica:
2802
2803 scoperta e recupero di delfino fossile. Notiziario della Soprintendenza per i Beni Archeologici
2804
2805 della Toscana 7, 250-252.
2806
2807

2808
2809 Almond, W. 1988. Armfield, B.A., Zheng, Z., Bajpai, S., Vinyard, C.J., Thewissen, J.G.M.,
2810
2811 2013. Development and evolution of the unique cetacean dentition. PeerJ 1:e24; DOI
2812
2813 10.7717/peerj.24.
2814
2815
2816
2817
2818
2819
2820

Arbeits, B., Cioppi, E., Dominici, S., Wierer, U., 2015. Certaldo (FI). Località San Giorsole: recupero di resti paleontologici. Notiziario della Soprintendenza per i Beni Archeologici della Toscana 11, 154.

Aronson, H.S., Zellmer, A.J. and Goffredi, S.K., 2017. The specific and exclusive microbiome of the deep-sea bone-eating snail, *Rubyspira osteovora*. FEMS Microbiology Ecology, 93, fiw250, doi: 10.1093/femsec/fiw250.

Azzellino, A., Aioldi, S., Gaspari, S., Lanfredi, C., Moulins, A., Podestà, M., Rosso, M., Tepsich, P., 2016. Risso's dolphin, *Grampus griseus*, in the Western Ligurian Sea: trends in population size and habitat use. In: Notarbartolo di Sciara, G., Podestà, M., Curry, B.E. (Eds.), Mediterranean Marine Mammals Ecology and Conservation. Advances in Marine Biology 75, 207-234, doi: 10.1016/bs.amb.2016.08.003.

Baldanza, A., Bizzarri, R., Famiani, F., Monaco, P., Pellegrino, R., Sassi, P., 2013. Enigmatic, biogenically induced structures in Pleistocene marine deposits: a first record of fossil ambergris. Geology 41, 1075-1078, doi: 10.1130/G34731.1.

Beardmore, S. R., Furrer, H., 2016. Evidence of a preservational gradient in the skeletal taphonomy of Ichthyopterygia (Reptilia) from Europe. Palaeogeography, Palaeoclimatology, Palaeoecology 443, 131-144.

Benson, R.B.J., Butler, R.J., Lindgren, J., Smith, A.S., 2010. Mesozoic marine tetrapod diversity: mass extinctions and temporal heterogeneity in geological megabiases affecting vertebrates. Proceedings of the Royal Society B: Biological Sciences, 277 (1683), 829-834.

Benvenuti, M., Bertini, A., Conti, C., Dominici, S., 2007. Integrated analyses of litho- and biofacies in a Pliocene cyclothemic, alluvial to shallow marine succession (Tuscany, Italy). *Geobios* 40, 143–158.

Benvenuti, M., Bertini, A., Conti, C., Dominici, S., Falcone, D., 1995a. Analisi stratigrafica paleoambientale integrata del Pliocene dei dintorni di San Miniato. Quaderni del Museo di Storia Naturale di Livorno, Supplemento, 29–49.

Benvenuti, M., Degli Innocenti, D., 2001. The Pliocene deposits in the Central-Eastern Valdelsa Basin (Florence, Italy), revised through facies analysis and unconformity-bounded stratigraphic units. *Rivista Italiana di Paleontologia e Stratigrafia* 107, 265-286.

Benvenuti, M., Del Conte, S., 2013. Facies and sequence stratigraphic modeling of a Upper Pliocene–Lower Pleistocene fluvial succession (Valdelsa Basin, central Italy). *Sedimentary Geology* 294, 303-314.

Benvenuti, M., Del Conte, S., Scarselli, N., Dominici, S., 2014. Hinterland basin development and infilling through tectonic and eustatic processes: latest Messinian-Gelasian Valdelsa Basin, Northern Apennines, Italy. *Basin Research* 26, 387-402.

Benvenuti, M., Dominici, S., 1992. Facies analysis, paleoecology and sequence stratigraphy in a Pliocene siliciclastic succession, San Miniato (Pisa, Italy). *Bollettino della Società Paleontologica Italiana* 31, 241-259.

Benvenuti, M., Dominici, S., Rook, L., 1995b. Inquadramento stratigrafico- deposizionale delle faune a mammiferi villafranchiane (unita` faunistiche Triversa e Montopoli) del Valdarno Inferiore nella zona a Sud dell'Arno. Il Quaternario 8, 457-464.

Benvenuti, M., Moratti, G., Sani, F., Bonini, M., Oriol, L.O., Papini, M., Rook, L., Cavallina, C., Cavini, L., 2015. Messinian-earliest Zanclean tectonic-depositional dynamics of the Cinigiano-Baccinello and Velona basins (Tuscany, Italy). Italian Journal of Geosciences, 134, 237-254.

Berta, A., 2012. *Return to the sea: the life and evolutionary times of marine mammals*. University of California Press, Berkeley, California, 205 pp.

Berta, A., Kienle, S., Bianucci, G., Sorbi, S., 2015. A Reevaluation of *Pliophoca etrusca* (Pinnipedia, Phocidae) from the Pliocene of Italy: phylogenetic and biogeographic implications. Journal of Vertebrate Paleontology, DOI: 10.1080/02724634.2014.889144 e889144, doi: 10.1080/02724634.2014.889144.

Berta, A., Lanzetti, A., Ekdale, E.G., and Deméré. T.A., 2016. From teeth to baleen and raptorial to bulk filter feeding in Mysticete cetaceans: The role of paleontological, genetic, and geochemical data in feeding evolution and ecology. Integrative and Comparative Biology 56, 1271-1284, doi: 10.1093/icb/icw128.

Bianchi, V., Ghinassi, M., Aldinucci, M., Boaga, J., Brogi, A., Deiana, R., 2015. Tectonically driven deposition and landscape evolution within upland incised valleys: Ambra Valley fill, Pliocene–Pleistocene, Tuscany, Italy. Sedimentology 62, 897-927, doi: 10.1080/17445647.2013.829412.

Bianucci, G., 1995. A new record of baleen whale from the Pliocene of Tuscany (Italy). Atti della Società Toscana di Scienze Naturali 102, 101-104.

Bianucci, G., 1996. The Odontoceti (Mammalia, Cetacea) from Italian Pliocene. Systematics and phylogenesis of Delphinidae. Palaeontographia Italica, 83: 73-167.

Bianucci, G., 1997a. *Hemisyntrochelus cortesii* (Cetacea, Delphinidae) from Pliocene sediments of Campore Quarry (Salsomaggiore Terme, Italy). Bollettino della Società Paleontologica Italiana 36, 75-83.

Bianucci, G., 1997b. The Odontoceti (Mammalia Cetacea) from Italian Pliocene. The Ziphiidae. Palaeontographia Italica 84, 163–192.

Bianucci, G., 1999. I cetacei fossili del Museo di Storia Naturale dell'Università di Pisa. Atti della Società Toscana di Scienze naturali, Memorie, Serie A 103, 63-68.

Bianucci, G., Bisconti, M., Landini, W., Storai, T., Zuffa, M., Giuliani, S., Mojette, A., 2002. Trophic interactions between white sharks (*Carcharodon carcharias*) and cetaceans: a comparison between Pliocene and recent data. 33-48. In Vacchi, M., La Mesa, G., Serena, F., Sèret, B., (eds). Proceedings 4ht Meeting of the European Elasmobranch Association, Livorno (Italy) 27–30 September 2000. Imprimerie F. Paillart, Abbeville, France, 209 pp.

Bianucci, G., Di Celma, C., Landini, W., Post, K., Tinelli, C., de Muizon, C., Gariboldi, K., Malinverno, E., Cantalamessa, G., Gioncada, A., Collareta, A., Gismondi, R.-S., Varas-Malca, R., Urbina, M., Lambert, O., 2016. Distribution of fossil marine vertebrates in Cerro

Colorado, the type locality of the giant raptorial sperm whale *Livyatan melvillei* (Miocene, Pisco Formation, Peru). Journal of Maps, 12, 543-557, doi: 10.1080/17445647.2015.1048315.

Bianucci, G., Landini, W., 2005. I paleositi a vertebrati fossili della Provincia di Pisa. Atti della Società Toscana di Scienze Naturali, Memorie serie A, 110, 1-22.

Bianucci, G., Mazza, P., Merola, D., Sarti, G., Cascella, A., 2001. The early Pliocene mammal assemblage of Val di Pugna (Tuscany, Italy) in the light of calcareous nannoplankton biostratigraphical data and paleoecological observations. Rivista Italiana di Paleontologia e Stratigrafia 107, 425-438.

Bianucci, G., Sarti, G., Catanzariti, R., Santini, U., 1998. Middle Pliocene cetaceans from Monte Voltraio (Tuscany, Italy). Biostratigraphical, paleoecological and paleoclimatic observations. Rivista Italiana di Paleontologia e Stratigrafia 104, 123-130.

Bianucci, G., Sorbini, C., 2014. Le collezioni a cetacei fossili del Museo di Storia Naturale dell'Università di Pisa. Museologia Scientifica Memorie 13, 93-102.

Bianucci, G., Sorbi, S., Vaiani, S.C., Landini, W., 2009a. Pliocene marine mammals from Italy: A systematic and stratigraphic overview. In: International conference on vertebrate palaeobiogeography and continental bridges across Tethys, Mesogea, and Mediterranean Sea, Bologna, Abstract book, 9-12.

Bianucci, G., Sorce, B., Storai, T., Landini, W., 2010. Killing in the Pliocene: shark attack on a dolphin from Italy. Palaeontology 53, 457-470, doi: 10.1111/j.1475-4983.2010.00945.x.

Bianucci, G., Vaiani, S.C., Casati, S., 2009b. A new delphinid record (Odontoceti, Cetacea) from the Early Pliocene of Tuscany (Central Italy): systematics and biostratigraphic considerations. *Neues Jahrbuch für Geologie und Paläontologie Abhandlungen* 254, 275-292.

Bianucci, G., Vomero, V., 2014. The cetacean collections of Italian museums. *Museologia Scientifica, Memorie* 13, Volume 2, 135 pp.

Bisconti, M., 2002. An early Late Pliocene right whale (genus *Eubalaena*) from Tuscany (Central Italy). *Bollettino della Società Paleontologica Italiana* 41, 83-91.

Bisconti, M., 2003. Evolutionary history of Balaenidae. *Cranium* 20, 9-50.

Bisconti, M., Lambert, O., Bosselaers, M., 2017. Revision of "*Balaena*" *belgica* reveals a new right whale species, the possible ancestry of the northern right whale, *Eubalaena glacialis*, and the ages of divergence for the living right whale species. *PeerJ* 5, e3464, doi: 10.7717/peerj.3464.

Boessenecker, R.W., Perry, F.A., Schmitt, J.G., 2014. Comparative taphonomy, taphofacies, and bonebeds of the Mio-Pliocene Purisima Formation, Central California: strong physical control on marine vertebrate preservation in shallow marine settings. *PLoS ONE* 9(3): e91419. doi:10.1371/journal.pone.0091419.

Boessenecker, R.W., Fordyce, R.E., 2014. Trace fossil evidence of predation upon bone-eating worms on a baleen whale skeleton from the Oligocene of New Zealand. *Lethaia* 48, 326-331.

Boldrocchi, G., Kiszka, J., Purkis, S., Storai, T., Zinzula, L., Burkholder, D. 2017. Distribution, ecology, and status of the white shark, *Carcharodon carcharias*, in the Mediterranean Sea. *Reviews in Fish Biology and Fisheries*, doi: 10.1007/s11160-017-9470-5.

Commented [1]:

Bonfil, R., Meyer, M., Scholl, M.C., Johnson, R., O'Brien S., et al., 2005. Transoceanic migration, spatial dynamics, and population linkages of white sharks. *Science* 310, 100-103.

Borselli, V., Cozzini, F., 1992. Il recupero di un cetaceo fossile in località Ponte a Elsa (Pisa). *Museologia Scientifica* 8, 9-22.

Bossio, A., Mazzei, R., Salvatorini, G., Sandrelli, F., 1993. Geologia dell'area compresa tra Siena e Poggibonsi ("Bacino del Casino"). *Atti della Società Toscana di Scienze Naturali* 107, 69-85.

Bossio, A., Mazzanti, R., Mazzei, R., Pascucci, V., Salvatorini, G.F., Sandrelli, F., 1994. Il Bacino di Volterra: notizie preliminari sull'evoluzione della sua area centro meridionale durante il Pliocene. *Studi Geologici Camerti*, volume speciale 1, 19-31.

Bossio, A., Foresi, L.M., Mazzanti, R., Mazzei, R., Salvatorini, G.F., 1997. Note micropaleontologiche sulla successione miocenica del Torrente Morra e su quella Pliocenica del bacino dei fiumi Tora e Fine (Provincia di Livorno e Pisa). *Atti della Società Toscana di Scienze Naturali* 104, 85-134.

3241
3242
3243
3244 Brogi, A., 2011. Bowl-shaped basin related to low-angle detachment during continental
3245 extension: the case of the controversial Neogene Siena Basin (central Italy, Northern
3246 Apennines). *Tectonophysics*, 499, 54-76.
3247
3248
3249

3250
3251
3252 Camp, C.L., 1980. Large ichthyosaurs from the Upper Triassic of Nevada.
3253 *Palaeontographica Abteilung A*, A170 (4-6), 139- 200.
3254
3255
3256
3257

3258
3259 Canese, S., Cardinali, A., Fortuna, C.M., Giusti, M., Lauriano, G., Salvati, E., Greco, S.,
3260 20067. The First identified winter feeding ground of finn whales (*Balaenoptera physalus*) in
3261 the Mediterranean Sea. *Journal of the Marine Biological Association of the U.K.* 86, 903-
3262 907.
3263
3264
3265
3266
3267

3268
3269 Capellini, G., 1876. Sulle balene fossili toscane. *Atti della Reale Accademia dei Lincei* 3, 9-
3270 14.
3271
3272
3273
3274
3275

3276 Capellini, G., 1883. Di un'Orca fossile scoperta a Cetona in Toscana. *Memorie*
3277 *dell'Accademia delle Scienze dell'Istituto di Bologna* 4, serie 4, 1-25.
3278
3279
3280
3281

3282 Capellini, G. 1905. Balene fossili toscane - III. *Idiocetus guicciardinii*. *Memorie*
3283 *dell'Accademia delle Scienze dell'Istituto di Bologna* 2, serie 6, 71-80.
3284
3285
3286
3287
3288

3289 Cavanagh, R.D., Gibson, C., 2007. Overview of the conservation status of cartilaginous
3290 fishes (Chondrichthyans) in the Mediterranean Sea. Gland, Switzerland and Malaga, Spain:
3291 The World Conservation Union (IUCN), vi + 42 pp.
3292
3293
3294
3295
3296
3297
3298
3299
3300

- 3301
3302 Chen, Z.-Q., Benton, M.J., 2012. The timing and pattern of biotic recovery following the end-
3303 Permian mass extinction. *Nature Geoscience* 5, 375-383.
3304
3305
3306
3307
3308 Cigala Fulgosi, F., 1990. Predation (or possible scavenging) by a great white shark on an
3309 extinct species of bottle-nosed dolphin in the Italian Pliocene. *Tertiary Research*, 12, 17-6.
3310
3311
3312
3313
3314 Cigala Fulgosi, F., Casati, S., Orlandini, A., Persico, D., 2009. A small fossil fish fauna, rich
3315 in *Chlamydoselachus* teeth, from the Late Pliocene of Tuscany (Siena, central Italy).
3316 *Cainozoic Research* 6, 3-23.
3317
3318
3319
3320
3321
3322
3323 Cioppi, E., 2014. I cetacei fossili a Firenze, una storia lunga più di 250 anni. *Museologia*
3324 *Scientifica Memorie* 13, 81-89.
3325
3326
3327
3328
3329 Cioppi, E., Dominici, S., 2011. Pliocene marine vertebrates, in: Monechi, S., Rook, L. (Eds.),
3330 The Museum of Natural History of the University of Florence, Volume 3, The geological and
3331 paleontological collections: Firenze, Firenze University Press, 224-235.
3332
3333
3334
3335
3336
3337
3338 Clarke, K.R., Gorley, R.N., 2006. PRIMER v6: User Manual/Tutorial. Plymouth: PRIMER-
3339 E., 190 p.
3340
3341
3342
3343
3344
3345 Cockcroft, V.G., Cliff, C., Ross, G.J.B., 1989. Shark predation on Indian Ocean bottlenose
3346 dolphins *Tursiops truncatus* off Natal, South Africa. *South African Journal of Zoology* 24,
3347 305-310.
3348
3349
3350
3351
3352
3353
3354
3355
3356
3357
3358
3359
3360

Collareta, A., Bosselaers, M. & Bianucci, G., 2016. Jumping from turtles to whales: a Pliocene fossil record depicts an ancient dispersal of *Chelonibia* on mysticetes. *Rivista Italiana di Paleontologia e Stratigrafia* 122, 35-44.

Compagno, L.J.V., 2002. Sharks of the world. An annotated and illustrated catalogue of shark species known to date. Volume 2. Bullhead, mackerel and carpet sharks (Heterodontiformes, Lamniformes and Orectolobiformes). *FAO Species Catalogue for Fishery Purposes* 1 (2) Rome, 269 p.

Cortés, E., 1999. Standardized diet compositions and trophic levels of sharks. *ICES Journal of Marine Science* 56, 707–717, doi: 10.1006/jmsc.1999.0489.

Croll, D.A., Marinovic, B., Benson, S., Chavez, F.P., Black, N., Ternullo, R., Tershy, B.R., 2005. From wind to whales: trophic links in a coastal upwelling system. *Marine Ecology Progress Series* 289, 117-130.

Cuzin-Roudy, J., 2011. Reproductive strategies of the Mediterranean krill, *Meganyctiphanes norvegica* and the Antarctic krill, *Euphausia superba* (Crustacea: Euphausiacea). *Invertebrate Reproduction & Development* 23, 105-114, doi: 10.1080/07924259.1993.9672301.

Danise, S., 2010. Modern and fossil shallow water whale fall communities. University of Florence, PhD Thesis, 178 pp.

Danise, S., Dominici, S., 2014. A record of fossil shallow-water whale falls from Italy. *Lethaia*, 47, 229–243.

Danise, S., Higgs, N.D., 2015. Bone-eating *Osedax* worms lived on Mesozoic marine reptile deadfalls. *Biology Letters* 11: 20150072.

Danise, S., Holland, S.M., 2017. Faunal response to sea-level and climate change in a short-lived seaway: Jurassic of the Western Interior, USA. *Palaeontology*, doi: 10.1111/pala.12278.

Danise, S., Twitchett, R.J., Matts, K., 2014. Ecological succession of a Jurassic shallow-water ichthyosaur fall. *Nature Communications* 5: 4789, doi: 10.1038/ncomms5789 | www.nature.com/naturecommunications.

DeCelles, P.G., 2012. Foreland basin systems revisited: variations in response to tectonic settings. In: C., Busby, and A., Azor (eds.), *Tectonics of Sedimentary Basins: Recent Advances*. Hoboken: Wiley-Blackwell, 405-426.

De Stefani, C., 1874. I terreni subapennini dei dintorni di S. Miniato al Tedesco. *Bollettino Malacologico Italiano* 7, 5-88.

Di Celma, C., Malinverno, E., Gariboldi, K., Gioncada, A., Rustichelli, A., Pierantoni, P.P., Bianucci, G., 2015. Stratigraphic framework of the late Miocene to Pliocene Pisco formation at Cerro Colorado (Ica Desert, Peru). *Journal of Maps*. doi:10.1080/17445647.2015.1047906.

Dicken, M.L., 2008. First observations of young of the year and juvenile great white sharks (*Carcharodon carcharias*) scavenging from a whale carcass. Marine and Freshwater Research 59, 596-602.

Dineen, A.A., Fraiser, M.L., Sheehan, P.M., 2014. Quantifying functional diversity in pre- and post-extinction paleocommunities: A test of ecological restructuring after the end-Permian mass extinction. Earth-Science Reviews 136, 339-349.

Dominici, S., 1994. Regressive-transgressive cycles from the Pliocene of the San Miniato area (Tuscany, Italy): paleoecology and sequence stratigraphy. Bollettino della Società Paleontologica Italiana Supplement 2, 117-126.

Dominici, S., Rook, L., Benvenuti, M., Abbazzi, L., 1995. Tapir remains in paralic deposits of Pliocene age in lower Valdarno (Tuscany, Italy): Facies analysis and taphonomy. Geobios 28, 131-135.

Dominici, S., Conti, C., Benvenuti, M., 2008. Foraminifer communities and environmental change in marginal marine sequences (Pliocene, Tuscany, Italy). Lethaia 41, 447-460.

Dominici, S., Cioppi, E., Danise, S., Betocchi, U., Gallai, G., Tangocci, F., Valleri, G., Monechi, S., 2009. Mediterranean fossil whale falls and the adaptation of mollusks to extreme habitats. Geology 37, 815–818.

Domning, D.P., 2001. Sirenians, seagrasses, and Cenozoic ecological change in the Caribbean. Palaeogeography, Palaeoclimatology, Palaeoecology 166, 27-50.

Dowsett, H.J., Robinson, M.M., Stoll, D.K., Foley, K.M., Johnson, A.L.A., Williams, M.,
Riesselman, C.R., 2013. The PRISM (Pliocene Palaeoclimate) reconstruction: Time for a
paradigm shift, *Philosophical Transactions of the Royal Society*, 371(2001):
doi:10.1098/rsta.2012.0524.

Ellis, J.A., Musick, J.A., 2007. Ontogenetic changes in the diet of the sandbar shark,
Carcharhinus plumbeus, in lower Chesapeake Bay and Virginia (USA) coastal waters.
Environmental Biology and Fishing 80, 51–67, doi: 10.1007/s10641-006-9116-2.

Estes, J.A., Burdin, A., Doak, D.F., 2016. Sea otters, kelp forests, and the extinction of
Steller's sea cow. *PNAS* 113, 880–885, doi: 10.1073/pnas.1502552112.

Estes, J.A., Terborgh, J., Brashares, J.S., Power, M.E., Berger, J., Bond, W.J., Carpenter,
S.R., Essington, T.E., Holt, R.D., Jackson, J.B.C., Marquis, R., Oksanen, L., Oksanen, T.,
Paine, R.T., Pikitch, E.K., Ripple, W.J., Sandin, S.A., Scheffer, M., Schoener, T.W., Shurin,
J.B., Sinclair, A.R.E., Soulé, M.E., Virtanen, R., Wardle, D.A., 2011. Trophic downgrading
of Planet Earth. *Science* 333, 301-306, doi: 10.1126/science.1205106.

Estes, J.A., Tinker, M.T., Williams, T.M., Doak, D.F. 1998. Killer whale predation on sea
otters linking oceanic and nearshore ecosystems. *Science* 282, 473–476.

Fischer, V., Bardet, N., Benson, R.B.J., Arkhangelsky, M.S., Friedman, M., 2016. Extinction
of fish-shaped marine reptiles associated with reduced evolutionary rates and global
environmental volatility. *Nature Communications* 7, 10825, doi: 10.1038/ncomms10825.

Fordyce, R.E. 2003. Cetacean evolution in Eocene-Oligocene oceans revisited. in *From Greenhouse to Icehouse: The Marine Eocene-Oligocene Transition*, D.R. Prothero, L.C. Ivany, E.A. Nesbitt, Eds. (Columbia Univ. Press, New York, 2003), 154–170.

Friedlaender, A.S., Goldbogen, J.E., Hazen, E.L., Calambokidis, J., Southall, B.L., 2014. Feeding performance by sympatric blue and fin whales exploiting a common prey resource. *Mar. Mamm. Sci.* 31, 345–354.

Fröbisch, N.B., Fröbisch, J., Martin Sander, P., Schmitz, L., Rieppel, O., 2013. Macropredatory ichthyosaur from the Middle Triassic and the origin of modern trophic networks. *Proceedings of the National Association of Science* 110, 1393-1397, doi:10.1073/pnas.1216750110.

Garassino, A., Pasini, G., De Angeli, A., Charbonnier, S., Famiani, F., Baldanza, A., Bizzarri, R., 2012. The decapod community from the Early Pliocene (Zanclean) of “La Serra” quarry (San Miniato, Pisa, Toscana, central Italy): sedimentology, systematics, and palaeoenvironmental implications. *Annales de Paléontologie* 98, 1–61.

Gattacceca, J., Deino, A., Rizzo, R., Jones, D.S., Henry, B., Beaudoin, B., Vadeboin F., 2007. Miocene rotation of Sardinia: New paleomagnetic and geochronological constraints and geodynamic implications. *Earth and Planetary Science Letters* 258, 359-377.

Ghinassi, M., 2007. Pliocene alluvial to marine deposits of the Val d'Orcia Basin (Northern Apennines, Italy): sequence stratigraphy and basin analysis. *Rivista Italiana di Paleontologia e Stratigrafia* 113, 459-472.

Ghinassi, M., Lazzarotto, A., 2005. Different source areas feeding the Early pliocene marine Radicofani Basin (Siena): geological and paleogeographical implications. *Bollettino della Società Geologica Italiana*, volume speciale 3, 15-28.

Gingerich, P.D., 2005. Cetacea. In: Rose K.D., Archibald J.D., editors. *Placental mammals: origin, timing, and relationships of the major extant clades*. Baltimore (MD), Johns Hopkins University Press. p. 234-252.

Gingerich, P.D., Ul-Haq, M., von Koenigswald, W., Sanders, W.J., Smith, B.H., Zalmout, I.S., 2009. New protocetid whale from the middle Eocene of Pakistan: birth on land, precocial development, and sexual dimorphism. *PloS One*. 4:e4366.

Goffredo, S., Dubinsky, Z., 2014. *The Mediterranean Sea: Its history and present challenges*. Springer-Verlag, Berlin, pp. 678.

Goldbogen, J.A., Calambokidis, J., Oleson, E., Potvin, J., Pyenson, N.D., Schorr, G., Shadwick, R.E., 2011. Mechanics, hydrodynamics and energetics of blue whale lunge feeding: efficiency dependence on krill density. *The Journal of Experimental Biology* 214, 131-146.

Greenpeace International, 2009. High seas Mediterranean marine reserves: a case study for the Southern Balearics and the Sicilian Channel. A briefing to the CBD's Expert workshop on scientific and technical guidance on the use of biogeographic classification systems and identification of marine areas beyond national jurisdiction in need of protection, Ottawa, 29 September–2 October 2009.

Guinot, G., Adnet, S., Cappetta, H., 2012. An analytical approach for estimating fossil record and diversification events in sharks, skates and rays. PLoS ONE 7(9): e44632. doi:10.1371/journal.pone.0044632.

Heyning, J.E., Dahlheim, M.E., 1988. *Orcinus orca*. Mammalian Species 304, 1–9, doi:10.2307/3504225.

Higgs, N.D., Glover, A.G., Dahlgren, T.G., Little, C.T.S., 2011. Bone-boring worms: characterizing the morphology, rate, and method of bioerosion by *Osedax mucofloris* (Annelida, Siboglinidae). Biological Bulletin 221:307-316.

Higgs, N.D., Little, C.T.S., Glover, A.G., Dahlgren, T.G., Smith, C.R., Dominici, S. 2012. Evidence of *Osedax* worm borings in Pliocene (~3 Ma) whale bone from the Mediterranean. Historical Biology 24, 269-277.

Higgs, N.D., Gates, A.R., Jones, D.O.B., 2014a. Fish food in the deep sea: revisiting the role of large food-falls. PLOS ONE 9(5): e96016

Higgs, N.D., Glover, A.G., Dahlgren, T.G., Smith, C.R., Fujiwara, Y., Pradillon, F., Johnson, S.B., Vrijenhoek, R.C., Little, C.T.S., 2014b. The morphological diversity of *Osedax* worm borings (Annelida: Siboglinidae). Journal of the Marine Biological Association of the United Kingdom 94, 1429–1439, doi: 10.1017/S0025315414000770.

Hocking, D.P., Marx, F.G., Park, T., Fitzgerald, E.M.G., Evans, A.R., 2017 A behavioural framework for the evolution of feeding in predatory aquatic mammals. Proc. R. Soc. B 284, 20162750, doi: 10.1098/rspb.2016.2750.

- Hogler, J.A., 1992. Taphonomy and paleoecology of *Shonisaurus popularis* (Reptilia: Ichthyosauria). *Palaios* 7, 108-117.
- Hu, S., Zhang, Q., Chen, Z.-Q., Zhou, C., Lü, T., Xie, T., Wen, W., Huang, J., Benton, M.J., 2011. The Luoping biota: exceptional preservation, and new evidence on the Triassic recovery from end-Permian mass extinction. *Proceedings of the Royal Society B*, 278, 2274–2282, doi:10.1098/rspb.2010.2235.
- Huusgaard, R.S., Vismann, B., Kühl, M., Macnaughton, M., Colmander, V., Rouse, G.W., Glover, A.G., Dahlgren, T., Worsaae, K., 2012. The potent respiratory system of *Osedax mucofloris* (Siboglinidae, Annelida) - a prerequisite for the origin of bone-eating *Osedax*? *PLoS ONE* 7(4): e35975. doi:10.1371/journal.pone.0035975.
- Jagt, J.W.M, Jagt-Yazykova, E.A., 2016. The Upper Cretaceous and lower Paleogene in the type area of the Maastrichtian stage (72.1–66 Ma). Field guide of the 14th Annual Meeting of the European Association of Vertebrate Paleontologists, 18 pp.
- Jenkins, R.G., Kaim, A., Sato, K., Moriya, K., Hikida, Y., and Hirayama, R. 2017. Discovery of chemosynthesis-based association on the Cretaceous basal leatherback sea turtle from Japan. *Acta Palaeontologica Polonica* 62, doi: 10.4202/app.00405.2017.
- Johnson, S.B., Warén, A., Lee, R.W., Kano, Y., Kaim, A., Davis, A., Strong, E.E., Vrijenhoek, R.C., 2010. *Rubyspira*, new genus and two new species of bone-eating deep-sea snails with ancient habits. *Biological Bulletin* 219: 166-177.

- Jorgensen, S.J., Reeb, C.A., Chapple, T.K., Anderson, S., Perle, C., Van Sommeran, S.R., Fritz-Cope, C., Brown, A.C., Klimley, A.P., Block, B.A., 2010. Philopatry and migration of Pacific white sharks. *Proceedings of the Royal Society B* 277, 679-688.
- Kaim, A., Kobayashi, Y., Echizenya, H., Jenkins, R., Tanabe, K., 2008. Chemosynthesis-based associations on Cretaceous plesiosaurid carcasses. *Acta Palaeontologica Polonica* 53: 97–104.
- Kiel, S., Goedert, J.L., 2006. Deep-sea food bonanzas: Early Cenozoic whale-fall communities resemble wood-fall rather than seep communities. *Proc. R. Soc. B* 273, 2625–2631, doi: 10.1098/rspb.2006.3620.
- Krijgsman, W., Hilgen, F.J., Raffi, I., Sierro, F.J., Wilson, D.S., 1999. Chronology, causes and progression of the Messinian salinity crisis. *Nature*, 400, 652-655.
- Kriwet, J., Benton, M.J., 2004. Neoselachian (Chondrichthyes, Elasmobranchii) diversity across the Cretaceous–Tertiary boundary. *Palaeogeography, Palaeoclimatology, Palaeoecology* 214. 181-94.
- Lambert, O., Bianucci, G., Post, K., de Muizon, C., Salas-Gismondi, R., Urbini, M., Reumer, J., 2010. The giant bite of a new raptorial sperm whale from the Miocene epoch of Peru. *Nature* 466, 105-108, do: 10.1038/nature09067
- Landini, W., 1977. Revisione degli 'Ittiodontoliti pliocenici' della collezione Lawley. *Palaeontographia Italica* 70, 92-134.

Landini, W., Bianucci, G., Bisconti, M., Carnevale, G., Sorbini, C., 2005. Il Plio-Pleistocene. I vertebrati marini. In Bonfiglio, L., (ed.) Paleontologia dei vertebrati in Italia. Evoluzione biologica, significato ambientale e paleogeografia. Museo Civico di Storia Naturale, Verona, 171-175.

Lawley, R., 1876. Nuovi studi sopra i pesci ed altri vertebrati fossili delle colline Toscane. Tipografia dell'Arte della Stampa, Firenze, 122 p.

Lawton, J.H., 1994. What do Species do in ecosystems? *Oikos* 71, 367-374.

Lazzari, P., Solidoro, C., Ibello, V., Salon, S., Teruzzi, A., Béranger, K., Colella, S., Crise, A. 2012. Seasonal and inter-annual variability of plankton chlorophyll and primary production in the Mediterranean Sea: a modelling approach. *Biogeosciences* 9, 217-233.

Lewison, R.L., Crowder, L.B., Read, A.J., Freeman, S.A., 2004. Understanding impacts of fisheries bycatch on marine megafauna. *Trends Ecol. Evol.* 19, 598–604.

Lindsay, E.H., Opdyke, N.D., Johnson, N.M., 1980. Pliocene dispersal of the horse *Equus* and late Cenozoic mammalian dispersal events. *Nature* 287, 135-138.

Liotta, D., 1996. Analisi del settore centro-meridionale del bacino pliocenico di Radicofani (Toscana meridionale). *Bollettino della Società Geologica Italiana* 115, 115-143.

Liu, J. Hu, S., Rieppel, O., Jiang, D., Benton, M.J., Kelley, N.P., Aitchison, J.C., Zhou, C., Wen, W., Huang, J., Xie, T., Lv, T., 2014. A gigantic nothosaur (Reptilia: Sauropterygia)

from the Middle Triassic of SW China and its implication for the Triassic biotic recovery. Scientific Reports 4, 7142, doi:10.1038/srep07142.

Lourenço, R., Penteriani, V., Rabac, J.E., Korpimäki, R., 2013. Lethal interactions among vertebrate top predators: a review of concepts, assumptions and terminology. Biol. Rev. doi: 10.1111/brv.12054.

Louwye, S., Marquet, R., Bosselaers, M., Lambert, O., 2010. Stratigraphy of an early-middle Miocene sequence near Antwerp in Northern Belgium (Southern North Sea Basin). Geologica Belgica, 13, 269-284.

Lundsen, L., Schlining, K.L., Frasier, K., Johnson S.B., Kuhnz, L.A., Harvey, J.B.J., Clague, G., Vrijenhoek, R.C., 2010. Time-series analysis of six whale-fall communities in Monterey Canyon, California, USA. Deep-Sea Research I 57, 1573–1584.

Manganelli, G., Benocci, A., 2014. I cetacei fossili del Museo di Storia Naturale dell'Accademia dei Fisiocritici di Siena. Museologia Scientifica Memorie 13, 103-110.

Mann, K.H., Lazier, J.R.N., 2006. Dynamics of marine ecosystems : biological-physical interactions in the oceans. Third edition. Blackwell Publishing, 496 pp.

Marsili, S., 2006. Analisi sistematica, paleoecologica e paleobiogeografica della selaciofauna plio-pleistocenica del Mediterraneo. Pisa, PhD Thesis, 246 pp.

- Marsili, S., 2007. Revision of the teeth of the genus *Carcharhinus* (Elasmobranchii; Carcharhinidae) from the Pliocene of Tuscany, Italy. *Rivista Italiana di Paleontologia e Stratigrafia* 113 (1), 79–96.
- Marsili, S., 2008. Systematic, paleoecologic and paleobiogeographic analysis of the Plio-Pleistocene Mediterranean elasmobranch fauna. *Atti della Società Toscana di Scienze Naturali, Serie A*, 113, 81-88.
- Martill, D.M., 1985. The preservation of marine vertebrates in the Lower Oxford Clay (Jurassic) of central England. *Philosophical Transactions of the Royal Society B* 311, 155-165.
- Martill, D. M., 1987. A taphonomic and diagenetic case study of a partially articulated ichthyosaur. *Palaeontology* 30, 543-555.
- Martill, D. M., Taylor, M. A., Duff, K. L., Riding, J. B., Bown, P. R., 1994. The trophic structure of the biota of the Peterborough Member, Oxford Clay Formation (Jurassic), UK. *Journal of the Geological Society* 151, 173-194.
- Martini, I., Aldinucci, M., Foresi, L.M., Mazzei, R., Sandrelli, F., 2011. Geological map of the Pliocene succession of the Northern Siena Basin (Tuscany, Italy). *Journal of Maps* 2011, 193-204.
- Martini, I., Foresi, L.M., Bambini, A.M., Riforgiato, F., Ambrosetti, E., Sandrelli, F., 2016. Calcareous plankton bio-chronostratigraphy and sedimentology of the “I Sodi” section

(Siena Basin, Italy): a key section for the uppermost Neogene marine deposition in the inner northern Apennines. Ital. J. Geosci. 135, 540-547, doi: 10.3301/IJG.2016.02.

Marx, F., G.M., Collareta, A., Gioncada, A., Post, K., Lambert, O., Bonaccorsi, E., Urbina, M., Bianucci, G., 2017. How whales used to filter: exceptionally preserved baleen in a Miocene cetotheriid. Journal of Anatomy 231, 212-220, doi: 10.1111/joa.12622.

Marx, G.M., Lambert, O., Uhen, M.D., 2016. Cetacean Palaeobiology. Topics in Paleobiology, Wiley-Blackwell, 336 pp.

Marx, F.G., Uhen, M.D., 2010. Climate, Critters, and Cetaceans: Cenozoic drivers of the evolution of modern whales. Science 327, 993-996.

Melanotte-Rizzoli, P., Artale, V., Borzelli-Eusebi, G.L., Brenner, S., Crise, A., et al., 2014. Physical forcing and physical/biochemical variability of the Mediterranean Sea: a review of unresolved issues and directions for future research. Ocean Science 10, 281-322.

Meyer, C.A., 2011. Amazing graze – Grazing traces of sea urchins on turtles – An example from the Late Jurassic of Switzerland: Annalen des Naturhistorisches Museum in Wien 113, 555-565.

Miyamoto, N., Yoshida, M.A., Koga, H., Fujiwara, Y., 2017. Genetic mechanisms of bone digestion and nutrient absorption in the bone-eating worm *Osedax japonicus* inferred from transcriptome and gene expression analyses. BMC Evolutionary Biology 17: 17, doi: 10.1186/s12862-016-0844-4.

- Motani, R., Jiang, D.-Y., Tintori, A., Sun, Y.-L., S., Hao, W.C., Boyd, A., Hinic-Frlog, S., Schmitz, L., Shin, J.-Y. and Sun Z.-Y. 2008. Horizons and assemblages of Middle Triassic marine reptiles from Panxian, Guizhou, China. *Journal of Vertebrate Paleontology* 28, 900-903.
- Motani, R., Ji, C., Tomita, T., Kelley, N., Maxwell, E., Jiang, D.-Y., Sanders, M., 2013. Absence of suction feeding ichthyosaurs and its implications for Triassic mesopelagic paleoecology. *PLoS ONE* 8(12): e66075. doi:10.1371/journal.pone.0066075.
- Motani, R., Jiang, D.-Y., Tintori, A., Rieppel, O., Chen, G.-B., 2014. Terrestrial origin of viviparity in Mesozoic marine reptiles indicated by Early Triassic embryonic fossils. *PLoS ONE* 9(2): e88640. doi:10.1371/journal.pone.0088640.
- Motani, R., Chen, X., Jiang, D., Cheng, L., Tintori, A., Rieppel, O., 2015. Lunge feeding in early marine reptiles and fast evolution of marine tetrapod feeding guilds. *Scientific Reports* 5, 8900. doi:10.1038/srep08900.
- Moulins, A., Rosso, M., Nani, B., Würtz, M., 2007. Aspects of the distribution of Cuvier's beaked whale (*Ziphius cavirostris*) in relation to topographic features in the Pelagos Sanctuary (north-western Mediterranean Sea). *Journal of the Marine Biological Association of the United Kingdom* 87, 177-186, doi: 10.1017/S0025315407055002.
- Murakami, M., Shimada, C., Hikida, Y., Soeda, Y., Hirano, H. 2014. *Eodelphis kabatensis*, a new name for the oldest true dolphin *Stenella kabatensis* Horikawa, 1977 (Cetacea, Odontoceti, Delphinidae), from the upper Miocene of Japan, and the phylogeny and

paleobiogeography of Delphinoidea, *Journal of Vertebrate Paleontology* 34, 491-511, doi: 10.1080/02724634.2013.816720.

Mutti, E., Davoli, G., Mora, S., Sgavetti, M., 1994. The eastern sector of the south-central folded Pyrenean foreland: criteria for stratigraphic analysis and excursion notes. 2nd International conference on high-resolution sequence stratigraphy, Guidebook. Istituto di Geologia, Università di Parma, Italy.

Muttoni G., Lanci L., Argnani A., Hirt A.M., Cibin U., Abrahamsen N., Lowrie W., 2001. Paleomagnetic evidence for a Neogene two-phase counterclockwise tectonic rotation in the Northern Apennines (Italy). *Tectonophysics* 326, 241-253-

Nalin R., Ghinassi M., Basso D., 2010. Onset of temperate carbonate sedimentation during transgression in a low-energy siliciclastic embayment (Pliocene of the Val d'Orcia Basin, Tuscany, Italy). *Facies* 56, 353-368.

Nalin, R., Ghinassi, M., Foresi, L., Dallanave, E., 2016. Carbonate deposition in restricted basins: a Pliocene case study from the central Mediterranean (Northwestern Apennines). *Journal of Sedimentary Research* 86, 1-32.

Navarro, J., López, L., Coll, M., Barría, C., Sáez-Liante, R., 2014. Short- and long-term importance of small sharks in the diet of the rare deep-sea shark *Dalatias licha*. *Mar Biol* 161, 1697-1707, doi: 10.1007/s00227-014-2454-2.

Noakes, S.E., Pyenson, N.D., McFall, G., 2013. Late Pleistocene gray whales (*Eschrichtius robustus*) offshore Georgia, U.S.A., and the antiquity of gray whale migration in the North Atlantic Ocean. *Palaeogeography, Palaeoclimatology, Palaeoecology* 392, 502-509.

Notarbartolo di Sciara, G., 1987. Killer whale, *Orcinus orca*, in the Mediterranean Sea. *Marine Mammal Science* 3, 356-360.

Notarbartolo di Sciara, G., Zanardelli, M., Jahoda, M., Panigada, S., Airoldi, S. 2003. The fin whale *Balaenoptera physalus* (L. 1758), in the Mediterranean Sea. *Mammal Review* 33, 105-150.

Notarbartolo di Sciara, G., Agardy, T., Hyrenbach, D., Scovazzi, T., Van Klaveren, P., 2008. The Pelagos sanctuary for Mediterranean marine mammals. *Aquatic Conservation: Marine and Freshwater Ecosystems* 18, 367-391.

Notarbartolo di Sciara, G., Podestà, M., Curry, B.E. (Eds.), 2016a, Mediterranean marine mammals ecology and conservation. *Advances in Marine Biology* 75, 458 pp.

Notarbartolo di Sciara, G., Castellote, M., Druon, J.-N., Panigada, S., 2016b. Fin whales, *Balaenoptera physalus*: at home in a changing Mediterranean Sea?. In: G. Notarbartolo Di Sciara, M. Podestà, B.E. Curry, eds., *Advances in Marine Biology* 75, 75-101, doi: 10.1016/bs.amb.2016.08.002.

Panigada, S., Notarbartolo di Sciara, G., 2012. *Balaenoptera physalus* (Mediterranean subpopulation). The IUCN Red List of Threatened Species 2012, e.T16208224A17549588, doi: 10.2305/IUCN.UK.2012.RLTS.T16208224A17549588.en.

- Parrish, F.A., 2009. Do monk seals exert top-down pressure in subphotic ecosystems? *Mar. Mamm. Sci.* 25, 91-106, doi: 10.1111/j.1748-7692.2008.00245.x.
- Pascucci, V., Costantini, A., Martini, I.P., Dringoli, R., 2006. Tectono-sedimentary analysis of a complex, extensional, Neogene basin formed on thrust-faulted, Northern Apennines hinterland: Radicofani Basin, Italy. *Sedimentary Geology*, 183, 71-97.
- Patzkowsky, M.E., Holland S.M., 2012. *Stratigraphic paleobiology. Understanding the distribution of fossil taxa in time and space.* Chicago: University of Chicago Press 256 pp.
- Pauly, D., Trites, A.W., Capuli, E., Christensen, V., 1998. Diet composition and trophic levels of marine mammals. *ICES Journal of Marine Science*, 55, 467-481.
- Pilleri, G., 1987. The cetacea of the Italian Pliocene with a descriptive catalogue of the species in the Florence Museum of Paleontology. Ostermundigen: Brain Anatomy Institute, University of Berne, Berne, 160 p.
- Pimiento, C., Griffin, J.N., Clements, C.F., Silvestro, D., Varela, S., Uhen, M.D., Jaramillo, C., 2017. The Pliocene marine megafauna extinction and its impact on functional diversity. *Nature Ecology & Evolution*, doi: 10.1038/s41559-017-0223-6.
- Pirotta, E., Matthiopoulos, J., MacKenzie, M., Scott-Hayward L., Rendell, L., 2011. Modelling sperm whale habitat preference: a novel approach combining transect and follow data. *Marine Ecology Progress Series* 436, 257-272, do: 10.3354/meps09236.

Podestà, M., Azzellino, A., Cañadas, A., Frantzis, A., Moulins, A., Rosso, M., Tepsich, P., Lanfredi, C., 2016. Cuvier's beaked whale *Ziphius cavirostris*, distribution and occurrence in the Mediterranean Sea: high-use areas and conservation threats. In: Notarbartolo di Sciara, G., Podestà, M., Curry, B.E. (Eds.), Mediterranean Marine Mammals Ecology and Conservation. Advances in Marine Biology 75, 105–142, doi: 10.1016/bs.amb.2016.07.007.

Pokines, J.T., Symes, S.A., 2013. Manual of forensic taphonomy. CRC Press, Boca Raton, FL, 496 p.

Polis, G.A., Anderson, W.B., Holt, R.D., 1997. Toward an integration of landscape and food web ecology: The dynamics of spatially subsidized food webs. Annual Review of Ecology and Systematics 28, 289-316.

Polis, G.A., Myers, C.A., Holt, R.D., 1989. The ecology and evolution of intraguild predation: potential competitors that eat each other. Annual Review of Ecology and Systematics 20, 297-330, doi: 10.1146/annurev.es.20.110189.001501.

Praca, E., Gannier, A., 2008. Ecological niches of three teuthophageous odontocetes in the northwestern Mediterranean Sea. Ocean Science 4, 49–59. <http://dx.doi.org/10.5194/os-4-49-2008>.

Prista, G.A., Agostinho, R.J., Cachão, M.A., 2015. Observing the past to better understand the future: a synthesis of the Neogene climate in Europe and its perspectives on present climate change. Open Geoscience 7, 65-83.

- Pyenson, N.D., Haasl, D.M., 2007. Miocene whale-fall from California demonstrates that cetacean size did not determine the evolution of modern whale-fall communities. *Biology Letters (Palaeontology)* 3, 709-711, doi: 10.1098/rsbl.2007.0342.
- Pyenson, N.D., Lindberg, D.R., 2011. What happened to gray whales during the Pleistocene? The ecological impact of sea-level change on benthic feeding areas in the North Pacific Ocean. *PLoS ONE* 6(7): e21295. doi:10.1371/journal.pone.0021295
- Raffi, S., Stanley, S., Marasti, R., 1985. Biogeographic patterns and Plio-Pleistocene extinction of *Bivalvia* in the Mediterranean and southern North Sea. *Paleobiology* 11, 368-388.
- Raymo, M.E., Hearty, P., De Conto, R., O'Leary, M., Dowsett, H.J., Robinson, M.M., Mitrovica, J.X., 2009. PLIOMAX: Pliocene maximum sea level project. *PAGES News* 17(2), 58-59.
- Rendell, L., Frantzis, A. 2016. Mediterranean sperm whales, *Physeter macrocephalus*: the precarious state of a lost tribe. In: Notarbartolo di Sciara, G., Podestà, M., Curry, B.E. (Eds.), *Mediterranean Marine Mammals Ecology and Conservation. Advances in Marine Biology*, 37-74, doi: 10.1016/bs.amb.2016.08.001.
- Reolid, M., Santos, A., Mayoral, E., 2015. Grazing activity as taphonomic record of necrobiotic interaction: A case study of a sea turtle carapace from the Upper Jurassic of the Prebetic (south Spain). *Revista Mexicana de Ciencias Geológicas* 32(1), 21-28.

Roemer, G.W., Gompfer, M.E., Van Valkenburgh, B., 2009. The ecological role of the mammalian mesocarnivore. *BioScience* 59, 165-173, doi: 10.1525/bio.2009.59.2.9.

Roman, J., Estes, J.A., Morissette, L., Smith, C., Costa, D., McCarthy, J.B., Nation, S.N., Pershing, A., Smetacek, V., 2014. Whales as marine ecosystem engineers. *Frontiers in Ecology and the Environment* 12 (7), 15 pp.

Roman, J., McCarthy, J.J., 2010. The whale pump: marine mammals enhance primary productivity in a coastal basin. *PLoS ONE* 5: e13255.

Roveri, M., Flecker, R., Krijgsman, W., Lofi, J., Lugli, S., Manzi, V., Sierro, F.J., Bertini, A., Camarlenghi, A., De Lange, G., Govers, R., Hilgen, F.J., Hübscher, C., Meijer, P.T.H., Stoica, M., 2014. The Messinian Salinity Crisis: Past and future of a great challenge for marine sciences. *Marine Geology* 352, 25-58.

Sarti, C., Lanzetti, A., 2014. I cetacei fossili del museo geologico Giovanni Capellini dell'università di Bologna. *Museologia Scientifica Memorie*, 70-78.

Scarponi, D., Kowalewski, M., 2004. Stratigraphic paleoecology: bathymetric signatures and sequence overprint of mollusk associations from upper Quaternary sequences of the Po Plain, Italy. *Geology* 32, 989-992.

Scheyer, T.M., Romano, C., Jenks, J., Bucher, H., 2014. Early Triassic Marine Biotic Recovery: The Predators' Perspective. *PLoS ONE* 9(3): e88987. doi:10.1371/journal.pone.0088987.

- Schmeisser McKean, R.L., Gillette, D.D., 2015. Taphonomy of large marine vertebrates in the Upper Cretaceous (Cenomanian-Turonian) Tropic Shale of southern Utah. *Cretaceous Research* 56, 278-292.
- Sergio, F., Schmitz, O.J., Krebs, C.J., Holt, R.D., Heithaus, M.R., Wirsing, A.J., Ripple, W.J., Ritchie, E., Ainley, D., Oro, D., Jhala, Y., Hiraldo, F., Korpimäki, E., 2014. Towards a cohesive, holistic view of top predation: a definition, synthesis and perspective. *Oikos* 123, 1234-1243.
- Slater, G.J., Goldbogen, J.A., Pyenson, N.D., 2017. Independent evolution of baleen whale gigantism linked to Plio-Pleistocene ocean dynamics. *Proc. R. Soc. B* 284, 20170546. <http://dx.doi.org/10.1098/rspb.2017.0546>.
- Smith, C., Glover, A.G., Treude, T., Higgs, N.D., Amon, D.J. 2015. Whale-Fall Ecosystems: Recent Insights into Ecology, Paleoecology, and Evolution. *Annual Review of Marine Science* 7: 10.1-10.26.
- Smith, F.A., Payne, J.L., Heim, N.A., Balk, M.A., Finnegan, S., Kowaleski, M., Lyons, K., McClain, C.R., McShea, D.W., Novack-Gottshall, P.M., Spaeth Anich, P., Wang, S.C., 2016, Body Size Evolution Across the Geozoic. *Annu. Rev. Earth Planet. Sci.* 44, 523-53, doi: 10.1146/annurev-earth-060115-012147.
- Sorbi, S., Domning, D.P., Vaiani, S.C., Bianucci, G., 2012. *Metaxytherium subapenninum* (Bruno, 1839) (Mammalia, Dugongidae), the latest sirenian of the Mediterranean Basin. *Journal of Vertebrate Paleontology* 32, 686-707.

Sorbini, C., Bianucci, G., Baldassarri, M., 2014. La balena fossile di Montopoli in Val d'Arno (Pisa). *Museologia Scientifica, Memorie* 13, 90-92.

Stambler, N., 2014. The Mediterranean Sea – Primary productivity. In: Goffredo, S., Dubinsky, Z., eds., *The Mediterranean Sea: Its history and present challenges*. Springer-Verlag, Berlin, 113-121.

Stanley, S., 1975. Adaptive themes in the evolution of the Bivalvia (Mollusca). *Annual Review of Earth and Planetary Sciences* 3, 361-385.

Steeman, M., 2010. The extinct baleen whale fauna from the Miocene-Pliocene of Belgium and the diagnostic cetacean ear bones. *Journal of Systematic Palaeontology* 8, 63-80, doi: 10.1080/14772011003594961.

Steeman, M., Hebsgaard, M.B., Fordyce, R.E., Ho, S.Y.W., Rabosky, D.L., Nielsen, R., Rahbek, C., Glenner, H., Sørensen, M.V., Willerslev, E., 2009. Radiation of extant cetaceans driven by restructuring of the oceans. *Systematic Biology* 58, 573-585.

Stinnesbeck, W., Frey, E., Rivas, L., Pardo Pérez, J., Leppe Cartes, M., Salazar Soto, C., Zambrano Lobos, P., 2014. A Lower Cretaceous ichthyosaur graveyard in deep marine slope channel deposits at Torres del Paine National Park, southern Chile. *Geological Society of America Bulletin* doi: 10.1130/B30964.1

Taboada, S., Riesgo, A., Bas, M., Arnedo, M.A., Cristobo, J., Rouse, G.W., Avila, C., 2015. Bone-eating worms spread: insights into shallow-water *Osedax* (Annelida, Siboglinidae) from Antarctic, Subantarctic, and Mediterranean Waters. *PLoS One* 10(11): e0140341.

Tinelli, C., Ribolini, A., Bianucci, G., Bini, M., Landini, W., 2012. Ground penetrating radar and palaeontology: the detection of sirenian fossil bones under a sunflower field in Tuscany (Italy). *Competes Rendus Palevol* 11, 445-454.

Tinelli, C. (2013). *Marine vertebrates from Pliocene shell beds from Tuscany (Italy): prospecting, taphonomy, palaeoecology and systematic palaeontology*. Pisa: PhD Thesis, 164 pp.

Tomašových, A., Dominici, S., Zuschin, M., Merle, D., 2014. Onshore-offshore gradient in metacommunity turnover emerges only over macroevolutionary time-scales. *Proceedings of the Royal Society B* 281, 1-10.

Tresguerres, M., Katz, S., Rouse, G.W., 2013. How to get into bones: proton pump and carbonic anhydrase in *Osedax* boneworms. *Proceedings of the Royal Society B* 280: 20130625.

Ugolini, R., 1899. Lo *Steno bellardii* Portis del Pliocene di Orciano Pisano. *Atti della Società Toscana di Scienze Naturali, Memorie* 17, 132-143.

Ugolini, R., 1902. Il *Monachus albiventer* Bodd. del Pliocene di Orciano. *Palaeontographia Italica* 8, 1–20.

Ugolini, R., 1907. Resti di vertebrati marini del Pliocene di Orciano. In: *Atti del Congresso dei Naturalisti Italiani promosso dalla Società Italiana di Scienze Naturali 1906*. Tipografia

degli Operai, Milano, 195-207.

Uhen, M.D., 2008. New protocetid whales from Alabama and Mississippi, and a new cetacean clade, Pelagicti. *Journal of Vertebrate Paleontology*, 28, 589-593.

Volkenandt, M., O'Connor, I., Guarini, J.-M., Berrow, S., O'Donnell, C., 2015. Fine-scale spatial association between baleen whales and forage fish in the Celtic Sea. *Canadian Journal of Fisheries and Aquatic Sciences* 73, 197-204, doi: 10.1139/cjfas-2015-0073.

Young, M.T., Brusatte, S.L., Beatty, B.L., Brandalise De Andrade, M., Desojo, J.B., 2012. Tooth-on-tooth interlocking occlusion suggests macrophagy in the Mesozoic marine *Crocodylomorph Dakosaurus*. *The Anatomical Record* 295, 1147-1158.

Williams, R., Okey, T.,A., Wallace, S., Gallucci, V.F., 2010. Shark aggregation in coastal waters of British Columbia. *Marine Ecology Progress Series* 414, 249-256.

Woodley, T.H., Gaskin, D.E., 1996. Environmental characteristics of North Atlantic right and fin whale habitat in the lower Bay of Fundy. Canada. *Canadian Journal of Zoology* 74, 75-84.

Figure and table captions:

Fig. 1 - Location of sedimentary logs within the largest Pliocene basins of Tuscany. Fine Basin (FB): Pagliana (1), Pieve Vecchia (2) and Orciano Pisano (3). Volterra-Era Basin (VEB): Parlascio (4), Lajatico (5), Fabbrica (6) and Volterra (7). Era Basin (EB): San Lorenzo (8), La Serra (9), Poggio al lupo (10), San Maiano (11), Canneto (12), Casenuove (13),

Castelfiorentino (14), Fiano (15) and San Gimignano (16). Orcia-Ombrore Basin: Arcille (17) and Poggio alle Mura (18). Siena-Radicofani Basin (SRB): Siena (19), Monteaperto (20), Castelnuovo Berardenga (21), Radicofani (22) and Fastelli (23). Chiana Basin (CB): Sinalunga (24), Cetona (25) and Allerona (26).

Fig. 2 - Pliocene stratigraphy of Tuscany, see Fig. 1 for the location of numbered sections. All logs measured and described by the authors, except Volterra (log 7: Bianucci et al. 1998), Arcille (log 17: Tinelli et al., 2012; Tinelli, 2013), Siena (log 19: Bianucci et al., 2001), Monteaperto (log 20: Martini et al., 2011), Castelnuovo Berardenga (log 21: Martini et al., 2016) and Radicofani (log 22: Ghinassi and Lazzarotto, 2005). Chronostratigraphy of localities 25 (Poltriciano, Cetona: Capellini, 1883) and 26 (Allerona: Danise, 2010) are unknown. Correlation between sections is also shown as boundaries of planktonic foraminifera biozones (dashed lines), following the scheme of Sprovieri (1992), based on available biostratigraphic studies for each basin (see main text for relevant references).

Fig. 3 - Abundance of fossil MM records in major museums of Tuscany, distributed by locality of provenance. Each record ranges from a single fragment or tooth, to articulated, nearly complete skeleton. A: masticates and odontocetes; B: large sharks; C: sirenians — scuba diver for scale in each figure. Symbols for basins as in Fig. 1.

Fig. 4 - Detailed sedimentary logs measured at three localities, representing three different stratigraphic contexts for the large marine vertebrate fossil record of the Tuscan Pliocene. The succession at Orciano Pisano is included in synthems S3-S4 of FB, at Arcille-Poggio alle Mura-Camigliano in synthem S2 of OOB, at Sinalunga in synthem S6 of CB. See Figs.1-2 for the location of the numbered localities and references in the main text for facies

analysis and sequence stratigraphy of synthem S5. Arcille log from Tinelli et al., 2012, and Tinelli, 2013.

Fig. 5 - Taphonomy of large marine vertebrates at Poggio alle Mura (Figs. 5A-5D) and Arcille (Fig. 5E, see Fig. 4 for the sequence stratigraphic and sedimentary context; plan view of the sirenian skeleton is modified from Tinelli et al., 2012), synthem S2. A: Plan view of the Poggio alle Mura undetermined balaenopterid. Parts of the skeleton are quasi-articulated, others are scattered, but not far from the original position. B: Quasi-articulated vertebrae of the same specimen, lying on top of a *Haustator* shell bed. C: Side view of a vertebra on top of the densely-packed shell bed. D. Detail of the shell bed, in top view. The turritelline gastropod *Haustator vermicularis* is visible in the upper left, a large fragment of wood in the lower right, with the ichnofossil *Teredolites* produced by wood-dwelling teredinid bivalves, in the centre of the photograph. E: Plan view of one of the Arcille sirenian specimens of *Metaxitherium appenninicum*. Same scale as in 5A, the arrows points to the North.

Fig. 6 - Taphonomy of a 10m-long, undetermined balaenopterid at Orciano Pisano (see Fig. 4 for the sequence stratigraphic and sedimentary context), synthem 4. A: Planimetry of the quasi-articulated and nearly complete skeleton. B: Detail of the central part of the skeleton in the field. The cortex layer of vertebrae and flipper bones is badly consumed, whereas some of the costae are still pristine. C: Lateral view of a turritellid shell bed, below, and the surface where the whale skeleton lied (dashed line), about 15 cm above the shell bed. The sediment is a very fine-grained silty sand, completely bioturbated (large vertical burrows are visible). D: Top view detail of the turritellid shell bed. At the center a valve of *Yoldia nitida*, surrounded by a few specimens of the turritellid *Archimediella spirata*.

Fig. 7 - Taphonomy of an incomplete, undetermined mystecete at Castel San Gimignano, synthem 4, comprising articulated torso elements. A: Planimetry of the articulated elements. B: Detail of one of the limbs in the field (trowel for scale = 22 cm): humerus, radius and ulna are in anatomical relationship; the cortex layer is well preserved, suggesting quick burial of the carcass. C: Bones of the chest region; on the background the massive sandstone associated with the fossil whale. Articulated shoreface pectinid bivalves (*Pecten flabelliformis*) were interspersed in the sandstone (photographs by Fabio Cozzini, 1985).

Fig. 8 - Taphonomy of a 5m-long, undetermined balaenopterid at Sinalunga (see Fig. 4 for a tentative sequence stratigraphic interpretation), synthem 6. A: Oblique view of the fully articulated skeleton lying in a gravelly sandstone, stratified in the lower part, massive in the upper. Vertebrae are tightly connected as if in life. B: Plan view of the flipper, with carpals, metacarpals and phalanges in perfect anatomical connection. C: The gravelly sandstone lies above a bioturbated muddy sandstone, with vertical burrows (*Ophiomorpha*). D: The lower part of the unit with the whale skeleton is formed by three fining-upward beds. Each bed grades from gravel to medium-grained sand. Articulated and empty shoreface bivalves (e.g., *Callista chione*), not in life position, are interspersed with the coarse gravel.

Fig. 9 - Quantitative analysis of the facies type and sequence stratigraphic distribution of large marine vertebrates, Pliocene of Tuscany (N = 39, see Tab. 2). A: The vast majority of cases (77%) are associated with fine-grained muddy sediments of the shelf, a few are found in deltaic coarse-grained strata. B: Most MM (69%) is found in highstand deposits, a few in proximity of the maximum flooding interval, or in transgressive deposits.

Fig. 10 - NMDS ordination of bulk samples (N = 72, see Fig. 2 for their stratigraphic position), based on the distribution of standardised abundances of 329 mollusc species (further

4981 explanation on multivariate techniques in the main text). A: Samples are subdivided based
4982 on the associated sedimentary facies types (F2-F5 in Tab. 1). The main axis ordinales
4983 samples along a paleodepth gradient, from shallowest to deepest, moving from the left to
4984 the right side of the bivariate plot. B: Same ordination, with an estimate of absolute
4985 paleodepth of each sample based on score along the main axis and calibrated through the
4986 average modern depth distribution of 23 extant species characterising the Pliocene dataset
4987 (abundance > 0.15%, see text; regression logarithmic curve in the inset). Vertebrates
4988 recovered in proximity of some of the samples (N = 13) are plotted on the diagram,
4989 confirming that, on average, the MM fossil record is concentrated on the open shelf at an
4990 estimated depth of 30-300 m (M = sirenians; S = sharks; P = pinnipeds; W = whales).
4991
4992
4993
4994
4995
4996
4997
4998
4999
5000
5001
5002
5003
5004

5005 Fig. 11 - Occupancy of trophic levels by Pliocene marine mammals and sharks in the north-
5006 western Mediterranean, expressed by number of species per trophic level (see Tabs 3, 5 for
5007 explanation and references). This figure, summing up data for the whole epoch, spanning
5008 circa 2,8 My, closely matches the association found in one single synthem S4, of much
5009 shorter duration (mid-Piacenzian, 3-4 hundred thousand years).
5010
5011
5012
5013
5014
5015
5016
5017

5018 Tab. 1 - Sedimentary facies types.
5019
5020
5021

5022 Tab. 2 - Stratigraphic, taphonomic and paleoenvironmental framework for Pliocene marine
5023 mammals recovered in Tuscany, with abundance data (N = 64). A: Articulated and quasi-
5024 articulated skeleton; C: Complete and quasi-complete skeleton; C+PC: Cranial and post-
5025 cranial remains.
5026
5027
5028
5029
5030
5031
5032

5033 Tab. 3 - Paleoecology of Pliocene large mammals and ecology of modern NWMS
5034 analogues.
5035
5036
5037
5038
5039
5040

5041
5042
5043
5044 Tab. 4 - Geographic distribution of Pliocene large mammals in Tuscany.
5045
5046
5047

5048 Tab. 5 . Paleoecology of Pliocene sharks and ecology of modern NWMS analogues.
5049
5050
5051

5052 Tab. 6 - Geographic distribution of Pliocene sharks in Tuscany.
5053
5054
5055

5056
5057 **Supplement Material**
5058
5059
5060


5061 Fig. S1 - Dendrogram resulting from cluster analysis of a dataset of 336 species distributed
5062 in 72 samples (standardised abundance, square-root transformed, Bray-Curtis similarity).
5063
5064
5065
5066
5067


5068 Tab. S1 - Ranked total average of standardised abundance of Pliocene molluscs.
5069
5070
5071




5072 Tab. S2 - Score on main axis of NMDS ordination (NMDS1) of Pliocene mollusc species
5073 (N=329).
5074
5075
5076
5077


5078 Tab. S3 - Average depth of extant mollusc species with >1,5% abundance in the Pliocene
5079 dataset (N=23).
5080
5081
5082
5083
5084


5085 Tab. S4 - Estimated depth of Pliocene samples (N=72) based on NMDS1.
5086
5087
5088
5089
5090
5091
5092
5093
5094
5095
5096
5097
5098
5099
5100
































Explanation

- ▲ Intertidal
- ▽ Shoreface
- Offshore transition
- ◆ Shelf
- Upper slope



2D Stress: 0,17





Explanation

